

The Escamilla Case In Court

The issue of U.S. Criminal jurisdiction to try Mario J. Escamilla for the alleged slaying of Bennie Lightsey, station manager on ice island T-3 (Fletcher's Ice Island) on 16 July 1970 was argued before U.S. Federal District Judge Oren R. Lewis in Alexandria, Virginia, on 5 May 1971.

The judge found sufficient grounds for taking jurisdiction and proceeded with the trial. Escamilla was then convicted of involuntary manslaughter. He has appealed.

Following are some observations on the jurisdictional aspect of the case.

1) It was evident throughout the hearings that the presiding judge wished, if possible, to decide the jurisdictional issue exclusively on the basis of U.S. domestic law, and that he was not disposed to address himself to abstruse questions of international law, once the Canadian waiver was on the record. At no point, therefore, did he express firm views on such matters as the international legal significance of T-3's origin, the possibility of permanent "possession" of such an island, the implications of a "ship" analogy as such for ice islands as a class, distinctions if any between ice islands and occupiable ice floes, and a host of other theoretical questions of universal, or at least arctic, applicability that might be considered of interest to an international lawyer.

2) In view of the very real, long continued, and unchallenged activities of U.S. Government agencies and Government grantees on T-3, the factual close connection between that particular ice platform and U.S. interests was so clear that the finding of proper criminal jurisdiction in the Escamilla case should probably not be given too broad a theoretical application.

3) One may argue that the case illustrated the extreme narrowness of the exclusively territorial basis for criminal jurisdiction (exceptions were noted) and that the judge was so troubled at the consequences of applying its pure logic in the Escamilla case that he chose instead to proceed on the basis of practical common sense and social responsibility, leaving it to others to split hairs over the meaning of commas in Section 7 of Title 18 of the U.S. Code or the analogies of ice islands with ships, guano islands, or commercial aircraft flying over the high seas.

Andreas G. Ronhovde
AINA Washington Office

A List of Vascular Plants from Polaris Bay, Northwest Greenland

The flora of the northwest coast of Greenland, north of 80° is not as well known as that of the northeast coast^{1,2}. Wulff³ collected about 70 species at various locations along the northwest coast between 81°25' and 83°06'N., and Simmons⁴ reviewed the earlier collections largely made at points south of 81°N. Two collections are reported from the vicinity of Polaris Bay, on the east side of Hall Basin at about 81°36'-40'N. Bessels⁵ lists 22 species collected there in 1872, and Hart⁶ lists 22 species collected by himself in May 1876, and in July and August 1876 by Dr. Copping of the same expedition. No recent reports are known of collections made in the Polaris Bay area.

On 17 August 1958, I had the opportunity to spend a few hours collecting vascular plants near the shores of Polaris Bay (81°36'N., 61°26'W.). Inland from the Bay a lowland gravel and clay plain extends eastwards towards Newman Bay. Along the coast of the Bay are dry, rocky-gravel and sandy terraces, with no prominent rock outcrops. A major river with its associated gravel and clay terraces and sand bars extends on to the coastal flats. Only a few coastal marshes exist above the intertidal zone which is very unstable owing to the action of large ice floes which are driven ashore. Generally the vegetation is sparse and is largely restricted to local areas favoured by more moisture. Collecting was restricted to the shores of the Bay for a distance of about one kilometre north of the river delta and for a distance of about 400 metres inland, onto the terraces of the lowland plain at a height of about 50 metres above sea level.

ANNOTATED LIST OF VASCULAR PLANTS

In the following list, the nomenclature of Porsild⁷ is used throughout. Specimens of all species collected are deposited in the National Herbarium, Ottawa, Canada. A few species (4) were noted but not collected and have been included in the list for completeness. Distribution records are based on the maps of Porsild⁷ and other literature reports. *Alopecurus alpinus* L. 410. Common on wet clay of old estuarine and littoral areas. *Phippsia algida* (Sol.) R. Br. 418. On moist clay flats of old river estuary. *Deschampsia brevifolia* R. Br. 413. Moist littoral clay flats.

- Poa alpigena* (Fr.) Lindm. var. *colpodea* (Fr.) Schol. 416. On dried up clay flats close to sea shore.
- Poa abbreviata* R. Br. 407, 417. Common on clay flats and along small streams draining to sea.
- Poa hartzii* Gand. 424, 426. Common on moist clay and sandy littoral areas.
- Pleuropogon sabinei* R. Br. 415. Growing in small coastal marsh.
- Colpodium vahlianum* (Liebm.) Nevski 398a, 425, 427. On sandy-clay littoral flats.
- Puccinellia angustata* (R. Br.) Rand & Redf. 408, 425. On wet littoral clay flats.
- Puccinellia andersonii* Swallen 398, 399, 412. Common on dry sandy-clay littoral flats and on sandy gravel bar.
- Eriophorum scheuchzeri* Hoppe 431. Around edge of littoral marsh.
- Carex ursina* Dew. 397. One stand on wet sandy bank of stream near shoreline.
- Carex maritima* Gunn. 421. On littoral sandy-clay soils.
- Carex stans* Drej. 414. Edge of small littoral marsh.
- Juncus biglumis* L. 406. On clay estuarine flats and in littoral zone.
- Salix arctica* Pall. Observed in littoral zone and up to about 30 m. above the shoreline.
- Oxyria digyna* (L.) Hill 409. In wet clay pocket on top of old gravel river terrace. 40 m.
- Polygonum viviparum* L. 424a. Moist clay flats.
- Stellaria crassipes* Hult. 411. Wet clay littoral areas.
- Stellaria monantha* Hult. 423a. Sandy-clay areas.
- Cerastium alpinum* L. var. (*C. arcticum* auct. non Lange) 403, 404. On clay flats and gravel slopes of old river terraces.
- Arenaria rubella* (Wahlenb.) Sm. 402. On sandy-clay flats below old river terrace.
- Melandrium apetalum* (L.) Fenzl. ssp. *arcticum* (Fr.) Hult. 401. Wet clay flats below old river terraces.
- Papaver radiculatum* Rottb. sensu lat. 400. Moist old river gravel terrace slopes.
- Cochlearia officinalis* L. ssp. *groenlandica* (L.) Porsild 420. Very common on moist sandy-clay littoral flats.
- Lesquerella arctica* (Wormskj.) Wats. Observed on dry gravel slopes of old river terraces.
- Draba bellii* Holm. 405. On rocky-gravel slopes down to shoreline.
- Braya purpurascens* (R. Br.) Bunge 419. On moist sandy-clay flats below old river terraces.
- Braya thorild-wulffii* Ostenf. 423. Sandy-clay flats below old river terraces.
- Saxifraga cernua* L. 422a. Moist littoral clay areas.
- Saxifraga nivalis* L. 420a. Moist littoral gravel slopes.
- Saxifraga oppositifolia* L. 422. Sandy-clay flats below old terraces.
- Saxifraga rivularis* L. 421a. In mosses at edge of small littoral marsh.
- Potentilla pulchella* Pursh 428, 429, 430. Common on sandy-clay and gravel soils.
- Dryas integrifolia* M. Vahl. Observed on dry gravel terrace slopes.
- Cassiope tetragona* (L.) D. Don 425a. On rocky gravel slope.
- Taraxacum* sp. ? *pumilum* Dahlst. Observed on moist clay flats. Recorded as *T. pumilum*.

The most significant range extension is for the littoral species *Carex ursina*, which had not previously been recorded north of 76° on the east coast and to 73° on the west coast of Greenland⁷. This species was recently collected from 81°25'N. on the west coast of Ellesmere Island⁸, but the present collection is the northernmost record of this species in the world. The collection of *Stellaria monantha* is a considerable range extension for Greenland, not having previously been recorded north of 78°, but is known from farther north in Canada⁷. Several other collections — *Poa alpigena* var. *colpodea*, *P. hartzii*, *Puccinellia andersonii*, and *Saxifraga rivularis* — are important range extensions in northwest Greenland, but are known from higher latitudes in northeast Greenland¹. *Puccinellia andersonii* has not been collected as far north as this in Canada, but the other 3 species are known from higher latitudes⁷. The collections of *Carex maritima* and *Draba bellii* are small northern range extensions in northwest Greenland. Several other species — *Deschampsia brevifolia*, *Carex stans* and *Arenaria rubella* would appear to be range extensions from Porsild's maps, but Ostenfeld³ reported them collected by Wulff at more northern latitudes in northwest Greenland.

The foregoing list comprises 37 species. Bessels⁵ and Hart⁶ both list 22 species from the vicinity of Polaris Bay. Simmons⁴ considered that some of the records of Bessels and Hall were wrongly determined, or could not be upheld as separate species, and produced a table recognizing only 25 species. From both lists can be added *Draba alpina* L. and *D. rupestris* R. Br., the latter probably being *D. subcapitata* Simm., and *Potentilla nivea* L. s. lat., which was not included by Simmons⁴. From Bessels' list⁵ can be added *DuPontia fisheri* R. Br. (*D. psilosantha* Rupr.), *Poa arctica* R. Br., *Ranunculus niva-*

lis L., and a *Pedicularis* sp. (not acknowledged by Simmons from this area, although he includes a doubtful collection under the Bessels Bay area), while the assignment of *Taraxacum palustre* DC. is in doubt. The record of *Ranunculus nivalis* var. is doubtful, although it has been recorded from the east coast of Ellesmere Island. It is probably a specimen of *R. sulphureus* Sol. which has been recorded elsewhere in northern Greenland. Polunin⁹ doubted the report of *Du-pontia fisheri* from this area, however the recent finding of this species by Brassard and Beschel⁸ at a similar latitude on the west coast of Ellesmere Island further supports this record. Bessels⁵ recorded the presence of *Carex dioica* L., which is usually considered as a synonym of *C. gynocrates* Wormskj., but the known distribution of this species in Greenland casts doubt on the identification of Bessels' specimen, as was indicated by Simmons⁴, Porsild¹⁰ and Polunin⁹. Hart⁶ includes a further *Draba* (*D. hirta* L.), *Melandrium* (*Lychnis*) *affine* (J. Vahl) Hartm., *Saxifraga caespitosa* L., *Luzula confusa* Lindebl. (*L. campestris* Sm.), *Festuca brachyphylla* (*F. ovina* L. var. *brevifolia* (R. Br.) Hart), and *Taraxacum Dens-leonis* Desf. (probably *T. phymatocarpum* J. Vahl). He also records *Potentilla frigida* Vill. and *Poa flexuosa* Wahl., but without the specimens it is difficult to assign these even tentatively. Ruling out any species for which the identification is in doubt, we can add 13 species from the lists of Bessels and Hart, giving a combined total of 50 species for the Polaris Bay area. This total is poor in number of species when compared with other areas of the High Arctic^{1,8,11}. Additional collecting will undoubtedly add other species, especially if habitats occurring at higher altitudes on Polaris Promontory, or further inland are included.

ACKNOWLEDGEMENTS

The Defence Research Board of Canada provided logistic support for my stay in the arctic as a member of "Operation Hazen". Lt. Cdr. J. P. Croal (RCN), liaison officer with "Operation Hazen" kindly arranged with Cdr. W. Reinhardt (USN), for my helicopter flights to the shores of Polaris Bay and back to the ice breaker U.S.S. Atka. Dr. A. E. Porsild, then Chief Botanist, National Museum of Canada, kindly verified or revised the determinations of all the collections.

John M. Powell
Canadian Forestry Service
Edmonton, Alberta
Canada

REFERENCES

- ¹Fredskild, B. 1966. Contributions to the flora of Peary Land, North Greenland. *Meddelelser om Grønland*, 178 (2), 1-23.
- ²Holmen, K. 1957. The vascular plants of Peary Land, North Greenland. *Meddelelser om Grønland*, 124 (9): 1-149.
- ³Ostenfeld, C. H. 1923. The vegetation of the north-coast of Greenland based upon the late Dr. Th. Wulff's collections and observations. *Meddelelser om Grønland*, 64 (9): 221-68.
- ⁴Simmons, H. G. 1909. Flowering plants and ferns from Northwest Greenland. *Report of the Second Norwegian Arctic Expedition in the "Fram" 1898-1902*. No. 16. 110 pp.
- ⁵Bessels, E. 1879. *Die Amerikanische Nordpol-Expedition*. Leipzig: Verlag von Wilhelm Engelmann. 647 pp.
- ⁶Hart, H. C. 1880. On the botany of the British Polar Expedition of 1875-6. *Journal of Botany*, N. S. 9: 52-6, 70-9, 111-5, 141-5, 177-82, 204-8, 235-42, 303-6.
- ⁷Porsild, A. E. 1964. Illustrated flora of the Canadian Arctic Archipelago. *National Museum of Canada, Bulletin* 146, 2nd. ed. revised. 218 pp.
- ⁸Brassard, G. R. and R. E. Beschel. 1968. The vascular flora of Tanquary Fiord, Northern Ellesmere Island, N.W.T. *Canadian Field-Naturalist*, 82: 103-13.
- ⁹Polunin, N. 1940. Botany of the Canadian Eastern Arctic. Part 1. Pteridophyta and spermatophyta. *National Museum of Canada, Bulletin* 92. 408 pp.
- ¹⁰Porsild, M. P. 1920. The flora of Disko Island and the adjacent coast of West Greenland from 66°-71° N. lat. with remarks on phytogeography, ecology, flowering, fructification and hibernation. *Meddelelser om Grønland*, 58: 1-156.
- ¹¹Powell, J. M. 1961. The vegetation and micro-climate of the Lake Hazen area, northern Ellesmere Island, N.W.T. *Defence Research Board, Canada, D. Phys R (G) Hazen* 14. 112 pp.

Winter Predation of *Mustela Erminea* in Northern Canada

Weasels are the most widely distributed mammalian predators in North America.¹ This paper reports the results of a study of short-tail weasel predation in northern Alberta and the Northwest Territories during the winter of 1964-65.