

Introduction

Health and health care provision continue to be top priorities of both individual Canadians and the Canadian government. In September of 2000, the First Ministers met to define goals and commitments regarding health in Canada. Their vision was that Canadians would have publicly funded health services that would provide quality health care and promote the health and well-being of Canadians in a cost-effective and fair manner. Accountability was a key part of the realization of that vision. Understanding this, the First Ministers agreed to direct Health Ministers in each province and territory to regularly report to the public on a number of health and health care system indicators. It was believed that clear public reporting, with appropriate, independent, third party verification would enhance the performance of health care in Canada.

A framework of 14 health indicators with common methods of measurement and reporting was developed through collaboration and consultation with health care professionals and other experts across Canada. The result is the following report, which gives the people of Nunavut a variety of comparable indicators measuring health status (e.g. life expectancy), health outcome (e.g., reduced burden of disease and illness) and quality of service (e.g. patient satisfaction). This report's primary objectives are:

- to allow the people of Nunavut and Canadians as a whole to see how we are doing in attaining our goals and objectives
- to assist individuals, governments, and health care providers in making more informed choices
- to promote the identification and sharing of best practices within Nunavut and across Canada, contributing to continuous service improvement
- to help Nunavut residents understand how their publicly funded health services are being delivered

We are pleased to be able to report on the majority of the 14 health indicators for Nunavut and are committed to providing the people of Nunavut with comprehensive and regular reporting on the health programs and services that we deliver. See the Note to the Reader below for those indicators that are excluded. It is our hope that this report will stimulate and guide further discussion and serve as a framework for collaborative action toward a healthier Nunavut.


List of Figures

Life Expectancy	1a.1	At birth in Canada and Nunavut, 1991-1999
	1a.2	At birth across Canada, 1999
	1a.3	At age 65 in Canada and Nunavut, 1991-1999
	1a.4	At age 65 across Canada, 1999
	1b.1	Disability-free at age 65 across Canada, 1996
	1b.2	Disability-free at age 65 in Canada and Nunavut, 1996
Infant Mortality	2.1	In Canada and Nunavut 1991-1999
	2.2	Across Canada, 1999
Low Birth Weight	3.1	Across Canada and Nunavut, 1991-1999
	3.2	Across Canada, 1999
Self-Rated Health	4.1	Across Canada, 2000-2001
	4.2	By age in Canada and Nunavut, 2000-2001
Mortality Rates	5a.1	Lung cancer in Canada and Nunavut, 1994-1999
	5a.2	Lung cancer across Canada, 1999
	5a.3	Breast cancer across Canada, 1999
	5a.4	Acute myocardial infarction in Canada and Nunavut, 1991-1999
Total Knee Replacement Rate	6b.1	Across Canada, 1999-2000
Cancer Incidence	7a.1	All cancer across Canada, 1997
	7a.2	All cancer in Nunavut, 1994-2000
	7a.3	Lung cancer in Nunavut, 1994-2000
	7a.4	Lung cancer across Canada, 1997
	7a.5	Breast cancer in Nunavut, 1994-2000
	7a.6	Breast cancer across Canada, 1997
Potential Years of Life Lost (PYLL)	7b.1	PYLL due to selected causes in Canada and Nunavut, 1999
	7b.2	PYLL due to lung cancer in Canada and Nunavut, 1994-1999
	7b.3	PYLL due to lung cancer across Canada, 1999
	7b.4	PYLL due to breast cancer in Canada and Nunavut, 1994-1999
	7b.5	PYLL due to breast cancer across Canada, 1999
	7b.6	PYLL due to AMI in Canada and Nunavut, 1991-1999
	7b.7	PYLL due to AMI across Canada, 1999
	7b.8	PYLL due to unintentional injuries in Canada and Nunavut, 1991-1999
	7b.9	PYLL due to unintentional injuries across Canada, 1999
	7b.10	PYLL due to suicides in Canada and Nunavut, 1991-1999
	7b.11	PYLL due to suicides across Canada, 1999
Vaccine Preventable Diseases	7c.1	Meningococcal disease in Nunavut, 1991-2000
	7c.2	Measles in Nunavut, 1989-2001
	7c.3	Haemophilus influenzae b (invasive) disease in Nunavut, 1990-2000


Indicator 1: Life Expectancy

Indicator 1a: Life Expectancy in Years at birth and at age 65

Life expectancy at birth is the number of years a person can expect to live, starting at birth, if current age-specific mortality rates continue to apply throughout his or her lifetime.



Since 1991, there has been a net increase in life expectancy at birth for Nunavut residents by 1.1 years, similar to the net increase for Canadians in general.




Life expectancy differs by gender. In all provinces and territories, females have longer life expectancies than males. Nunavut females, with a life expectancy of 70.2 years, exceed Nunavut males by 2.5 years.


For Nunavut residents, no gender difference can be seen in life expectancy at age 65. In 1999, Nunavut women could expect to live 7 years less than their Canadian counterparts, and Nunavut men could expect to live 3.2 years less.

Indicator 1b: Disability-Free Life Expectancy in Years at age 65

Disability-free life expectancy is the number of years an average individual would be expected to live free of moderate or severe disability, starting at age 65. Disability-free life expectancy emphasizes quality of life. It is used to distinguish between years of life free of any activity limitation and years experienced with at least one activity limitation. To that end, disability-free life expectancy establishes a threshold based on the nature of such limitations. Years of life lived in conditions above this threshold are counted in full. Those lived in conditions below the threshold are not counted.




On average, a 65 year old Nunavut resident can expect to live 9.2 years disability-free, two and a half years less than the Canadian average. Nunavut men and women share roughly the same disability-free life expectancy.




In comparison to Nunavut, Canadian females can expect to live 3.2 more years disability-free, and Canadian males can live 1.6 more years disability-free.

Indicator 3: Low Birth Weight

Low birth weight is defined as live births with a birth weight less than 2,500 grams, expressed as a percentage of all live births with known birth weights. It has been adjusted for borderline viable births by excluding all birth weights under 500 grams.




Trends in low birth weight appear to have changed very little between 1991 and 1999 in both Nunavut and the whole of Canada.



Low birth weight tends to show a gender difference, with females in all jurisdictions experiencing a greater percentage of low birth weight. Low birth weight babies are more common in Nunavut than Canada as a whole. In 1999, about 35% more infants were born underweight in Nunavut compared to the rest of Canada.

Figure 4.2 Self-Reported Health by Age in Canada and Nunavut
(reported as very good or excellent)
2000-2001




Source: Canadian Community Health Survey - Cycle 1.1, 2000/01.
*Numbers for Nunavut residents age 65+ are too small to be reported with confidence. See note 7.

Self-reported health differs by age group. Nunavut residents in the 35-44 year age bracket tend to consider their health the best, while 45-64 year olds consider their health the worst. This is slightly different from the national average, where the greatest percentage of people reporting very good or excellent health lies in the 20-34 year age bracket.

Female Breast Cancer Mortality Rates


Because there have been few cases of breast cancer in Nunavut relative to Canada as a whole, time trend data for breast cancer mortality is too unreliable to be published. It is very difficult in a small population to assess real trends because small increases or decreases become exaggerated.



Roughly half as many women in Nunavut die of breast cancer when compared to Canada. This may partially be explained by the higher breastfeeding and fertility rates in Nunavut, both of which have some protective effect against breast cancer.

Acute Myocardial Infarction Mortality Rates

Between 1991 and 1999, AMI mortality rates for Nunavut are lower than for Canada; however, the Nunavut rates are now approaching Canadian rates. As Canadian rates seem to steadily decline, Nunavut rates appear to be growing closer to the national average. This may reflect the changes in lifestyle in the past ten years among Nunavut residents that are bringing them closer to their counterparts across the country.




Indicator 7: Reduced Burden of Disease, Illness and Injury


Indicator 7a: Incidence Rates for all cancers and lung and breast cancers

All Cancer Incidence

All cancer incidence is defined as the number of newly diagnosed primary cancer cases of all types in a given year, per 100,000 population. Rates have been age standardized according to the 1991 Canadian population.




In 1997, unlike most other Canadian jurisdictions, women in Nunavut were more likely to be diagnosed with any cancer than men. The rate for Nunavut males was lower than the national average, but the rate for women was higher.




Between 1994 and 2000, incidence rates of all cancers in Nunavut seem to have declined slightly for males while increasing for females. 35% of all cancer diagnosed in women in Nunavut between 1991 and 1996 was cervical cancer.

Female Breast Cancer Incidence

Female breast cancer incidence is defined as the number of newly diagnosed primary breast cancer cases in a given year, per 100,000 population.




There appears to be an upward trend in breast cancer rates in Nunavut between 1994 and 2000.




However, in 1997, breast cancer rates were still much lower than the national average. In fact, women in Nunavut were almost three times less likely to be diagnosed with breast cancer than Canadian women as a whole. This may partially be explained by the higher breastfeeding and fertility rates in Nunavut, both of which have some protective effect against breast cancer.

Potential Years of Life Lost due to Lung Cancer




Between 1994 and 1999, potential years of life lost due to lung cancer appears stable for Canada, but increasing for Nunavut.




In 1999, Nunavut had high rates for PYLL due to lung cancer when compared with the national average. This applies to both men and women: rates are 1.3 times higher for men and 2.2 times higher for women. Unlike most of the provinces, PYLL due to lung cancer is higher for women than men in Nunavut.

Potential Years of Life Lost due to Acute Myocardial Infarction (AMI)




Between 1991 and 1999, it appears that there was a steady decline in PYLL due to acute myocardial infarction in Canada. Rates in Nunavut appear fairly stable in this time period, but they are considerably lower than the national average.




PYLL due to AMI for Canadian males were almost nine times higher than rates for men in Nunavut in 1999. There were no cases of AMI among Nunavut women in 1999. Low rates among Nunavut residents are partially due to the traditional Inuit diet, which consists of large amounts of marine foods rich in n-3 fatty acids. Studies have shown that n-3 fatty acids have beneficial effects on key risk factors for cardiovascular disease.

Potential Years of Life Lost due to Suicides




PYLL due to suicides were consistently higher in Nunavut than in Canada between 1991 and 1999, and while Canadian rates are stable, rates in Nunavut appear to be rising.




As in all provinces and territories, more men in Nunavut commit suicide than women. In 1999, this resulted in PYLL due to suicides for men more than eight times that of women. As well, Nunavut men had PYLL due to suicides roughly eight and a half times higher than Canadian men in general. In the same year, Nunavut women had an approximately four times greater rate of PYLL than their national counterparts.

High PYLL due to suicide reflects the high suicide rate in Nunavut. Most suicides are committed by young single Inuit males by hanging. Many factors come into play, among them acculturation, alcohol abuse, depression, and impulsiveness.


Indicator 7c (iii): Haemophilus Influenzae b (invasive) Disease Incidence



Haemophilus influenzae b (Hib) disease is expressed as the rate of new cases reported by year in children under 5, per 100,000 population. Invasive disease includes meningitis, bacteraemia, epiglottitis, pneumonia, pericarditis, septic arthritis, or emphysema. Hib was the most common cause of bacterial meningitis and a leading cause of other serious invasive infections in children prior to the introduction of the Hib vaccine in 1988.¹⁵ As an infectious disease, Hib tends to appear in outbreaks followed by periods of low incidence. When reported per 100,000, the rates look dramatic, however, they represent a small number of cases.



While the majority of Canadians are satisfied with the service provided by their family physicians and other doctors, satisfaction among Nunavut residents is lower than anywhere else. The difference is likely due to the fact that the model of health care delivery in Nunavut has primary care provided by the Community Health Nurse with physicians assisting. Few communities have a resident doctor.




Across Canada as well as in Nunavut, the majority of patients surveyed were satisfied with the community-based care they received. Nurses in a community care setting provide most primary health care in Nunavut, and residents have indicated satisfaction with their care.


Indicator 13: Public Health Surveillance and Protection

Indicator 13a: Tuberculosis

Tuberculosis is reported as the rate of new active and relapsed infectious cases reported by calendar year, per 100,000 population.




Tuberculosis rates in Nunavut appear to be climbing since 1991, reaching a peak of nearly 200 cases per 100,000 in 2000.




In 1999, Nunavut's tuberculosis incidence rate was about 17 times higher than the Canadian average. The high rate is partly due to a large reservoir of latent tuberculosis from earlier cases that reactivate as Nunavut residents age.

Indicator 13c: Verotoxigenic *E. coli*

Verotoxigenic *E. coli* is expressed as the rate of new cases reported by year, per 100,000 population.



Over the past ten years, reported cases of *E. coli* infection are relatively rare in Nunavut. An outbreak occurred in 1991 and 2000. In Nunavut, verotoxigenic *E. coli* is mainly spread through contaminated hamburger meat, appearing in occasional outbreaks.





The graph above depicts the occurrence of *E. coli* cases in Canada in 2001. The rate is calculated per capita and although Nunavut is almost two times higher than the national average, this actually only represents one case in Nunavut.

Indicator 14: Health Promotion and Disease Prevention

Indicator 14a: Youth Smoking Rates

Smoking rates are expressed as the percentage of the population aged 12 and over who are current or daily smokers. Current smokers are those who reported smoking either on a daily or occasional basis when interviewed for the Canadian Community Health Survey. Daily smoking refers to smoking at least one cigarette per day for each of the 30 days preceding the survey. Occasional smoking refers to smoking at least one cigarette during the past 30 days preceding the survey, but not every day.



The percentage of teenagers in Nunavut who currently smoke is nearly two and a half times the national average. In both Canada and Nunavut, teenage girls are more likely to smoke than teenage boys. The high smoking rate probably contributes to the lung cancer statistics for Nunavut. Nunavut has more new cases of lung cancer per year, more potential years of life lost to lung cancer, and higher lung cancer mortality rates than Canada as a whole.

