



D003641

**WORKSHOP ON PERMAFROST DATA
RESCUE AND ACCESS**


GLOBAL GEOCRYOLOGICAL DATABASE

DATA APPLICATIONS ONE DATABASE — MANY USES

USER REQUIREMENTS

RESEARCH

OPERATIONAL



**ENVIRONMENTAL
PROTECTION**

An illustration of the diverse user requirements of the Global Geocryological Database (GGD).

GLACIOLOGICAL **DATA**

REPORT GD-28

WORKSHOP ON PERMAFROST DATA RESCUE AND ACCESS 3-5 November 1994, Oslo, Norway

by

Roger G. Barry
NSIDC
University of Colorado
Campus Box 449
Boulder, Colorado 80309-0449 U.S.A.

J. Alan Heginbottom
Geological Survey of Canada
Department of Energy, Mines and Resources
Ottawa, K1A 0E8
Canada

Jerry Brown
International Permafrost Association
Arlington, Virginia 22219-0200
U.S.A.

Published by:

WORLD DATA CENTER FOR GLACIOLOGY
[SNOW AND ICE]
Cooperative Institute for Research in Environmental Sciences
University of Colorado
Boulder, Colorado 80309-0449 U.S.A.

WDC operated for:
U.S. Department of Commerce
National Oceanic and Atmospheric Administration
National Environmental Satellite, Data, and Information Service
Boulder, Colorado 80303 U.S.A.

June 1995

DESCRIPTION OF THE WORLD DATA CENTER SYSTEM¹

The World Data Centers (WDCs) were established in 1957 to provide archives for the observational data resulting from the International Geophysical Year (IGY). In 1958 the WDCs were invoked to deal with the data resulting from the International Geophysical Cooperation 1959, the one-year extension of the IGY. In 1960, the International Council of Scientific Unions (ICSU) Comité International de Géophysique (CIG) invited the scientific community to continue to send to the WDCs similar kinds of data from observations in 1960 and following years, and undertook to provide a revised *Guide to International Data Exchange* for that purpose. In parallel the CIG inquired of the IGY WDCs whether they were willing to treat the post-IGY data; with few exceptions, the WDCs agreed to do so. Thus the WDCs have been serving the scientific community continuously since the IGY, and many of them archive data for earlier periods.

In November 1987 the International Council of Scientific Unions (ICSU) Panel on World Data Centers prepared a new version of the *Guide to International Data Exchange*, originally published in 1957, and revised in 1963, 1973 and 1979. The new publication, *Guide to the World Data Center System, Part 1, The World Data Centers (General Principles, Locations and Services)*, was issued by the Secretariat of the ICSU Panel on World Data Centers. This new version of the *Guide* contains descriptions of each of the twenty-seven currently operating disciplinary centers, with address, telephone, telex, and contact persons listed. The reader is referred to the new *Guide* for descriptions of the responsibilities of the WDCs, the exchange of data between them, contribution of data to WDCs, and the dissemination of data by them. The WDCs for Glaciology are listed below.

World Data Center A for Glaciology [Snow and Ice]

Address:	WDC-A for Glaciology CIRES, Campus Box 449 University of Colorado Boulder, Colorado 80309-0449 USA
Telephone:	(303)492-5171
Fax:	(303)492-2468
Network Address:	nsidc@kryos.colorado.edu
Director:	Dr. R. G. Barry

World Data Center B1

Address:	World Data Center B1 for Glaciology Molodezhnaya 3 Moscow 117296 USSR
Telephone:	130-05-87
Telex:	411478 SGC SU

World Data Center C for Glaciology

Address: WDC-C for Glaciology
Scott Polar Research Institute
Lensfield Road
Cambridge CB2 1ER
UNITED KINGDOM
Telephone: (0223)336556
Telex: 81240 CAMSPL G
Network Address: ojm21@cus.cam.ac.uk
Manager: Oliver J. Merrington

World Data Center D for Glaciology [Snow and Ice] and Geocryology

Address: Lanzhou Institute of Glaciology and Geocryology
Chinese Academy of Sciences
Lanzhou 730000, China
Telephone: (86)0931-082-2815
Fax: (86) 931-888-5241
Director: Professor Cheng Guodong

The following organization provides international data services including data analyses and preparation of specialized data products. It merges the previous activity of the Permanent Service on the Fluctuations of Glaciers and the Temporary Technical Secretariat for World Glacier Inventory. These activities are not part of the WDC system but the center cooperates with WDCs in the discipline. Users wishing assistance in seeking data or services from this group may contact an appropriate WDC.

World Glacier Monitoring Service (WGMS)

Dr. W. Haeberli
Section of Glaciology
VAW/ETH, ETH Zentrum
CH-8092 Zurich
SWITZERLAND
Telephone: +41-1-532-4093
Fax: +41-1-632-1192
Internet: haeberli@vaw.ethz.ch

¹Adapted from *Guide to the World Data Center System. Part 1. The World Data Centers (General Principles, Locations and Services)*. International Council of Scientific Unions. Panel on World Data Centers, November 1987, 91pp.

FOREWORD

This issue of *Glaciological Data* highlights international efforts to improve the accessibility of data on permafrost, ground ice and seasonally frozen ground and to rescue data sets that may be at risk. A small workshop on the prioritization of permafrost data and on improved access to data in general was organized under the auspices of the International Permafrost Association, through its Executive Committee and the Working Group on Permafrost Data and Information. The meeting took place at the Norwegian Geotechnical Institute (NGI) in Oslo, Norway, 3-5 November 1994. We wish to thank Dr. Susan Lacasse, Director of the NGI, for hosting the meeting, and especially Dr. Odd Gregerson, NGI, and Dr. Kaare Flaate, Norwegian Directorate of Public Roads, for their generous assistance with the local arrangements.

The attendance of Russian and Chinese delegates at the meeting and partial assistance to some other non-government scientists was made possible by funds from NOAA's Earth System Data and Information (ESDIM) program for data rescue and the National Science Foundation Arctic System Science data management program support to NSIDC. The assistance of Claire Hanson, Paul Farley and Cindy Brekke, NSIDC, in the coordination of the travel arrangements was indispensable. Claire Hanson also prepared the Data Base Inventory format included as Appendix 1. Ann Brennan has assisted greatly with the editing and Lyn Ryder and Carol Pedigo with word processing.

Roger G. Barry
Professor of Geography and
Director, WDC-A for Glaciology/NSIDC
University of Colorado
Boulder, CO 80309-0449

CONTENTS

FOREWORD	vii
Workshop on Permafrost Data Rescue and Access, Norwegian Geotechnical Institute, Oslo, Norway, 3-5 November 1994	
1. Background	i
1.1. The International Permafrost Association	i
1.2. NOAA Earth System Data and Information Management Program	i
1.3. Arctic System Science Program of the US National Science Foundation	2
2. Permafrost Data and Applications	2
3. Workshop Objectives	5
3.1. Prioritizing Data Sets	5
3.2. Identification of Candidate Data Sets	5
3.3. The Global Geocryological Database (GGD) Concept	5
3.4. Data Base Structure	5
3.5. The GGD Process	6
4. Priorities for Data Rescue	6
4.1. Demands for Permafrost Data	7
4.1.1. Validation of Earth System Models	7
4.1.1.1. Permafrost and Ground Ice	7
4.1.1.2. Snow Cover	8
4.1.3. Process Studies/Process Model Development and Verification	9
4.2. Guidelines for Prioritizing Data Sets	9
4.3. High Priority Data Sets	10
4.4. General Recommendations	12
5. Next Steps	12
6. References	14
7. Agenda	15
8. Participants	17
Appendix 1	IPA Global Geocryological Database Inventory
Appendix 2	Global Geocryological Database: Suggestion for Data Structure
Appendix 3	Russian Contribution to the Global Geocryological Database
Appendix 4	Draft Protocols: a Possible Model for the GGD
Appendix 5	Russian Permafrost Map Inventory
	Russian Permafrost Map Inventory Index by Scale
Appendix 6	List of Institutions in Russia and the Commonwealth of Independent States Involved in Studies of Permafrost and Seasonally-frozen Ground
Appendix 7	International Arctic Environmental Data Directory
Acronyms	131
Book Notes	133

Workshop on Permafrost Data Rescue and Access

Norwegian Geotechnical Institute
Oslo, Norway
3-5 November 1994

1. Background

The origins of this workshop and its purpose arose from several parallel and converging initiatives within the science and data programs of the International Permafrost Association, the National Oceanic and Atmospheric Administration's (NOAA) Earth System Data and Information Management (ESDIM) program and the National Science Foundation (NSF) Arctic System Science (ARCSS) program. These three strands are summarized below.

1.1. The International Permafrost Association

At the ninth council meeting of the International Permafrost Association (IPA) in Beijing, China, 8 July 1993, a resolution was adopted to "seek a more active role in International Geosphere-Biosphere Program (IGBP) core projects" and for "relevant IPA working groups [to] give particular attention to global climate change." The IPA Working Group on Data and Information seeks to improve and standardize the collection, archiving, documentation and dissemination of permafrost and ground ice data (*Frozen Ground*, No. 14, December 1993, p. 8 and p. 10).

The report of a workshop on permafrost data and information, held at the Fifth International Conference on Permafrost in Trondheim, Norway, on 7 August 1988 (Brennan and Barry, 1989) recommended that the IPA, through its working groups, should seek to ensure the security of key historical information and data records and to facilitate the development of user-friendly data systems, including data directories. Some preliminary steps were reported by Barry and Brennan (1993). As an outcome of those initiatives, during the winter of 1993-94, the IPA Working Group on Data and Information, in conjunction with the IPA Executive Committee, developed a draft prospectus for a Global Geocryological Database (GGD). This was circulated for comment and finalized at a two-day workshop held at the GeoData Institute, University of Southampton, UK, on 30 June-1 July 1994. The IPA Executive Committee, at its meeting on 2-3 July, endorsed the project and the proposed course of action. Accordingly, a workshop was organized in Oslo, Norway, by the Working Group on Data and Information to establish priorities for data rescue, to develop an implementation plan and procedures for data recovery, storage and dissemination, and to review the results of pilot projects underway in Russia and the UK.

1.2. NOAA Earth System Data and Information Management Program

The National Oceanic and Atmospheric Administration has instituted a program of Earth System Data and Information Management to coordinate data and information management activities on an agency-wide basis. The specific objectives of the program are to:

- Rescue critical NOAA environmental data currently at risk of being lost.
- Improve access to NOAA environmental data and information for scientists and administrators.
- Modernize and interconnect environmental data systems throughout NOAA to increase their capability and responsiveness.
- Assist in developing standards for data documentation, data quality, and network connectivity.
- Provide agency-wide-guidelines on developing policies related to environmental data management.
- Build a top-level consensus within NOAA on data management issues, and formulate a vision of the agency's data and information management strategy for the 1990s and beyond.

The final objective is addressed in the ESDIM plan and involves the construction of a strategic approach to data management and information that can be applied agency-wide. The early focus of the ESDIM implementation plan is on the rescue of critical NOAA environmental data, with the cryosphere being one of the first topics addressed. The first Workshop on Cryospheric Data Rescue and Access was held in May 1993 (Crane, 1993). The Permafrost Data Rescue and Access Workshop built on the approach developed at the earlier meeting; it addressed specific problems of data rescue and improved access to data. Data rescue in this context refers to saving data sets that are critical for scientific research. This may involve: copying data from existing magnetic tapes to new tapes or to other media; transcribing disintegrating or otherwise inaccessible historical paper records to digital, analog, or micro-form; or the compilation of new data sets from highly varied, original sources with different media types. The concept of data rescue, in the context of this workshop, focused on data records relating to permafrost, ground ice and seasonally frozen ground, and to the consideration of potential future data sources. Although the primary objective of the workshop was data rescue, the discussion included plans to make data readily accessible to the user community. Data access focuses on providing, or enhancing, the ability of researchers to access existing or rescued data sources.


1.3. Arctic System Science Program of the US National Science Foundation

Since September 1994 the National Snow and Ice Data Center (NSIDC), University of Colorado at Boulder, has been funded as the ARCSS Data Coordination Center for all components of the United States Arctic System Science (ARCSS) Program; ARCSS represents NSF's global change program for the Arctic. This unified data project follows separate grants for ARCSS data management pilot projects relating to the Ocean-Atmosphere-Ice Interactions (OAI) and Land-Atmosphere-Ice Interactions (LAI), and the Greenland Ice Sheet Project Two (GISP2) components of ARCSS. Current NSIDC efforts focus on identifying ways to integrate the ARCSS component communities, and on providing access to existing, unarchived data of interest for ARCSS research. In parallel, the planning for archiving of ARCSS-funded data sets continues in concert with the OAI, LAI, GISP2 and PALE (Paleoclimates of Arctic Lakes and Estuaries) Science Management Offices. Close contact is maintained with the emerging Human Dimensions component, the Surface Heat Budget of the Arctic Ocean (SHEBA) experiment planning process, and the developing Synthesis, Integration and Modeling component. The concept of "System Science", or integration, depends heavily on the accessibility and sharing of data and results among all those involved. NSIDC is seeking to develop the ways and means to ensure that accessibility.

As part of the LAI component and in keeping with the underlying ARCSS principles of integration and data accessibility, NSIDC is collaborating with the IPA Working Group on Data and Information in its efforts to identify and archive permafrost and other active-layer data for the IPA Global Geocryological Database. As an example, J. Brown and F. Nelson have provided active layer temperature and thickness, soil moisture, and snow depth data from the former US Army Cold Regions Research and Engineering Laboratory (CRREL) site and the new ARCSS site at Barrow for the first ARCSS/LAI CD-ROM product, "North Slope Alaska Data Sampler" (NSIDC, 1994). The CD-ROM also contains climate, soils, vegetation, and river runoff data. Such information is identified as vital for documenting the climate signal, for assessing changes in hydrological regimes, biologic processes and the Arctic ecosystem (McCauley and Meier, 1991).

2. Permafrost Data and Applications

The cryosphere, representing the solid phase of the hydrosphere, occupies a unique place in the global water cycle. Ice in the atmosphere plays a vital role in the precipitation process and, and at the surface of the oceans, it drastically modifies the ocean-atmospheric exchanges of heat and momentum. When ice occurs on land, it represents a major source of fresh water for societal use, and acts as a significant agent for geomorphic activity. The widespread distribution of permafrost in the Northern Hemisphere is shown in Figure 1. Past changes in climate have led to variations in the extent of the cryosphere, and the effects of this can still be seen in the geomorphic and isostatic history of large parts of the middle and high latitudes. Changes in the extent and distribution of the cryosphere itself



KEY TO PERMAFROST AND GROUND ICE REGIONS			
NORTH AMERICA		NORTH ATLANTIC	
Canada		12	Greenland
1	Queen Elizabeth Islands	13	Iceland
2	Western Arctic Lowlands	14	Svalbard
3	Interior Plains	15	Fennoscandia
4	Eastern Arctic and Canadian Shield	ASIA	
5	Basins within Canadian Shield	Russia	
6	Cordillera	16	Arctic Islands and Arctic Ocean Continental Shelf
7	Areas outside the permafrost region	17	European North
USA		18	Urals
8	Alaska	19	West Siberia
9	Conterminous USA and Mexico	20	East Siberia
		21	Northeast Siberia
		22	Far East
		23	Southern Siberia and Trans-Baikalia
		24	Mongolia
		China	
		25	Northeast China
		26	Tibet Plateau
		27	West China and other Mountain Permafrost Areas
		28	Japan and Korea
		29	Central Asia
		30	Southwestern Asia
		EUROPE	
		31	Central and Southern Europe

Figure 1. The distribution of permafrost in the Northern Hemisphere. (See Brown *et al.*, in press; Heginbottom *et al.*, 1993)

have a positive feedback on climate, and thus the magnitudes of past climatic changes are likely to be linked, in part, to the expansion and contraction of the snow and ice cover. Similarly, global climate model predictions of future global warming through enhanced greenhouse effects are strongly dependent on the effectiveness of the cryospheric response.

The cryosphere interacts with the earth system in a variety of ways, the most important of which is its interaction with climate. While climate controls on the cryosphere are readily apparent (in general outline, if not in detail), what may be less obvious is the way in which changes in the permafrost can feed back to influence the climate system. The presence of permafrost has a strong influence on soil water and runoff and therefore on the energy and moisture balance in cold regions. Thawing of thin permafrost, or a deepening of the active layer would greatly modify the runoff regimes, both water and sediments, as well as the soil moisture balance. Frozen organic materials also sequester substantial amounts of radiatively active trace gases (CO_2 , CH_4 , etc.) both within and beneath the permafrost body; their potential release to the atmosphere involves an important and little understood biogeochemical/climate link. Permafrost contains large quantities of ground ice and, once thawed, the ground becomes unstable, promoting dramatic increases in sediment yields.

Feedbacks between the cryosphere and the rest of the earth system have led to the suggestion that the permafrost is not only an important component of the system; possibly instrumental in mediating global change, but it may also be a sensitive indicator of such global change processes (see Koster et al., 1994, for an annotated bibliography). The attention focused on the cryosphere has been prompted by the important role that the snow and ice plays in General Circulation Models' (GCM) climate change experiments, together with the recognition that the cryosphere has undergone large variations in the past that correspond to periods of extensive changes in global climate. It is important to note, however, that many processes involving perennially or seasonally frozen ground are not well understood and are very poorly simulated by current numerical climate models. Permafrost data are thus essential for validating Earth System Models and for improving model physics, as well as for monitoring environmental change and variability. These two (overlapping) requirements are central components of the IGBP and many national global change programs, and they contributed to themes for the data rescue discussions in the present workshop. Additionally engineering design and environmental impacts were also considered.

The cryosphere is an appropriate target for ESDIM's early data rescue efforts not only because of the apparent importance of these data, but also because of the nature of the data themselves. Permafrost-related data encompass a wide range of parameters including, for example, depth and distribution of permafrost and seasonally frozen soils, as well as snow cover extent, depth, and mass balance, periglacial features, chemistry and temperature structure from boreholes in permafrost. In addition to their contribution to studies of snow cover-vegetation-ground ice interactions and trace gas fluxes, these data have operational applications in hydrology, engineering, shipping/fisheries, and off-shore development. Scientific applications, as noted above, tend to focus on cryosphere-climate interactions, but again this implies a wide range of possibilities that would include, in the case of permafrost, ground ice and seasonally-frozen ground, hydrological processes, and paleoclimatic reconstructions. The importance of these permafrost conditions is reported in the most recent impacts assessment of the Intergovernmental Panel on Climate Change, Working Group II; the report containing a chapter on the cryosphere is in preparation and final review.

Apart from the wide range of possible data sets and applications, several other factors complicate decision making with regard to cryospheric data management. For example, data sets are not application specific, and where time series of active-layer thickness and its physical and chemical properties may be of use to scientists interested in modeling frozen ground processes, they may also be of interest to engineers concerned with construction of pipelines and transportation facilities. The importance of any given data set will, therefore, vary according to the application concerned, which must obviously be reflected in the guidelines developed for prioritizing data sets, i.e., prioritizing of data sets must also involve some prioritizing of scientific objectives.

Permafrost data have been collected by both governmental and non-governmental agencies from many different countries. This raises problems of data acquisition and highlights the problem encountered when integrating data collected over varying temporal and spatial scales that exist on different media with a wide range of formats. Some further indication of the nature of the problem is shown in earlier data surveys/inventories conducted by the World Data Center-A for Glaciology (e.g. Barry, 1988; Crane, 1993).

3. Workshop Objectives

3.1. Prioritizing Data Sets

A first objective of the workshop was to derive a set of guidelines for data set selection that will facilitate the successful implementation of a rescue and access program. Given the limited resources initially available for the project, the volume of data involved, and the wide range of data sets/applications noted above, it was recognized that guidelines must be established early in the program.

3.2. Identification of Candidate Data Sets

A second objective was to identify high-priority data sets that satisfy the criteria recognized as a result of the first objective. The purpose was to begin the process of data set selection by identifying data sets with which to begin the data rescue activity.

3.3. The Global Geocryological Database (GGD) Concept

The specific objectives of the GGD project are to help identify, acquire and disseminate data on permafrost and frozen ground to serve several important purposes:

- To advance the scientific understanding of permafrost, with specific reference to relationships among climate, process, material and morphology; definition of paleo-permafrost conditions; and specification of future long-term environmental monitoring programs.
- To improve the basis of engineering design in cold regions, for both contemporary and predictive purposes.
- To aid in understanding and predicting global and regional climatic change, and specifically to support the verification of general circulation models and trace gas cycles.
- To offer a basis for detecting environmental change at a range of temporal and spatial scales, particularly through establishing and managing long-term, wide-area monitoring programs.
- To enhance the basis for developing environmental scenarios and assessing environmental impact, including pollution and the socioeconomic implications of environmental change in cold regions, for planning and environmental protection.

The meeting endorsed the proposition of M.J. Clark (see Appendix 2) of "one database, many uses" and the proposed organization of the Russian National Geocryological Database (NGD), (see Appendix 3).

3.4. Data Base Structure

As currently proposed, the GGD will consist of an internationally distributed system of linked data centers or nodes. Information from regions of perennially and seasonally frozen ground will be assembled in National or Regional Geocryological Databases (NGDs, RGDs) and/or selected World Data Centers. The information will be made available to the scientific, engineering, environmental and policy communities.

The general issue of data structure was specifically addressed by M.J. Clark and J. Branson; their report is included in its entirety in Appendix 2. The group made no specific recommendations, however, as to a preferred structure.

3.5. The GGD Process

The GGD will operate by identifying existing data sets, current and historic; rescuing those that are at risk of being lost; managing the acquired data; and making data available to the scientific and engineering communities either in raw form or processed into specific usable forms of information. Standard data descriptions will be held in national and international directories, and users will gain access to the data through a variety of modes and media. The IPA is working with user communities to identify priorities for data rescue, acquisition and monitoring. Initially, the emphasis is being placed on retrieving data and time-series that are in danger of being lost.

Once identified and described, data sets will be organized into standard file structures and accessioned by an appropriate National Geocryological Database or regional node. Currently, the designated nodes are:

- Federal Center for Geocological Systems, Moscow, Russia
- GeoData Institute, Southampton, UK
- World Data Center A for Glaciology, Boulder, Colorado, USA
- World Data Center D for Glaciology and Geocryology, Lanzhou, China

The GGD nodes are using their existing in-house facilities, but are also developing additional funding to support NGD/GGD data rescue and management activities, as well as to promote information generation and dissemination through analysis and modeling.

Further working links have been established with other organizations pursuing similar goals for data rescue, monitoring, management and dissemination.

These include:

- Global Resources Information Database (UNEP/GRID), Arendal, Norway; Director: Svein Treitdal
- International Arctic Science Committee (IASC), Oslo, Norway (see IASC, 1994); President: Magnús Magnússon
- Scientific Committee on Antarctic Research (SCAR); Chair: A.C. Clarke
- Council of Managers of National Antarctic Programs (COMNAP), Ad Hoc Planning Group on Antarctic Data Management, Cambridge, UK; Co-Chair: M.R. Thorley
- World Conservation Monitoring Centre (UNEP/WCMC), Cambridge, UK
- World Glacier Monitoring Service, Zurich, Switzerland; Director: W. Haeberli.

The topic of access to the GGD was raised by N.N. Romanovskii and M. Liebman (Appendix 3). A proposed protocol for use of the database was presented by J. Branson and is included as Appendix 4.

4. Priorities for Data Rescue

The results of the workshop discussions are presented under four headings:

- **Demands for permafrost data.** Priority should be given to data sets for which there is a high demand or that are important for a critical research goal. As resources are limited and as we cannot anticipate all future demands for data, priority is given to research areas considered to be important today, as well as to monitoring impact assessment and engineering needs. Three of these areas fall within the current US Global Change program:

data sets identified as being important for the validation of general circulation of global climate or earth system models, for system monitoring, and for process studies. Specific cryospheric parameters are assessed in the light of all these areas of application, and each is prioritized as discussed below.

- **Guidelines for prioritizing data sets.** A set of guidelines was presented for assessing the relative importance of prospective data sets and their priority in the data rescue effort.
- **High priority data sets.** Two groups of data sets are identified – one group having high priority and worthy of immediate attention, and one group that should be considered in the data rescue effort, but have a lower priority or require more information before priority can be fully assessed.
- **General recommendations.** A set of recommendations is presented for the near-term implementation of the permafrost data rescue effort.

4.1. Demands for Permafrost Data

As noted above, the workshop discussion focused on emphasizing data sets useful for parameterizing or validating large-scale GCMs or earth system models, data sets that could be used for monitoring climate change and variability, and data for developing or validating empirical or numerical models of system processes. It is also recognized that data are required to support specific international and national programs, e.g., Global Energy and Water Experiment (GEWEX), Arctic Climate System Study (ACSYS), Arctic System Science (ARCSS), International Tundra Experiment (ITEX), etc.

Although not considered specifically in this workshop, it should be noted that there are important linkages between the distribution and erosion of permafrost terrain and sea level (which is a concern over time scales of 100 years), and between the cryosphere and hydrology (important in terms of future water resources and the timing and quantity of runoff). Both of these are important research questions that may require data not discussed here.

4.1.1. Validation of Earth System Models

The results from the present generation of climate models suggest that the model climate and its sensitivity to greenhouse gas-induced climate change are greatly affected by the cryosphere, particularly by the distribution of sea ice and snow cover. This is exemplified by the fact that most model results show their greatest greenhouse-induced warming at high latitudes, and at least one study has shown that about one third of the temperature feedback is due to albedo changes in high latitudes.

4.1.1.1. Permafrost and Ground Ice

The primary importance of ground ice and permafrost is the way in which they modify the surface thermal and moisture regimes and how they respond to these surface modifications (thermokarst, etc.). In this regard, the most important parameters are ground ice extent, active layer depth and moisture content. Given this, we can establish that the highest priority data sets will be those that verify a model's ground ice extent and ground temperature above the depth of zero annual amplitude (DZAA). Most of the present generation of climate models include only a simple treatment of ground processes (Nelson et al., 1993). However, temperature and moisture are treated in newer models of surface processes. For example, a permafrost subcomponent has been developed by the UK Meteorological Office (UKMO) Hadley Centre and implemented in a single column model. For each of four soil layers, the variables of soil temperature, liquid and frozen water content are calculated based on the hydraulic and thermal properties of the soil. Dr. H. Cattle, of the UKMO (personal communication, 1994), confirms that the current data requirements include the seasonal variations in soil temperature and moisture profiles, together with information on the annual variations in thaw depth. Currently, research projects in several countries are obtaining soil moisture and temperature data, as reported at the 1994 Fall Meeting of the American Geophysical Union. Maps of these variables and of the current distribution of permafrost would be useful model validation data. In

addition, seasonal patterns of freeze-thaw may be important in the context of modelling and predicting trace gas fluxes.

For Earth System Models, highest priority will also go to those data sets that are global in coverage and have sufficient duration to derive climatological statistics (mean and variance). Although monthly data are adequate for most purposes, original data must be collected at sufficient temporal resolution to derive representative monthly averages. Several soil models utilize daily and monthly input data (see Waelbroeck, 1993). A possible future development could be a move towards the use of statistical methods to derive the spatial variability of the temperature-moisture ground-ice parameters within a grid cell. This would require the collection of data at a higher spatial resolution than the surface grid of the model (i.e., 2 degrees or better) in order to derive the spatial relationship for use in the model.

4.1.1.2. Snow Cover

Snow cover was not specifically discussed at the workshop. However, there is a consensus among permafrost scientists and engineers that snow cover is a key control in mediating ground temperature and must be considered in all models. It is worth noting that state-of-the-art Earth System Models predict the fractional snow cover, snow depth and water equivalent, and layer temperature. Little work has been done on the sensitivity of climate models to snow cover, the importance of which appears to be due to its albedo (and the related masking effect on the vegetation canopy), and to its effect on the surface hydrology and active layer temperature and moisture regimes. The required data sets for model validation are snow cover extent, snow water equivalent, and snow depth (for land surface models that allow vegetation masking by snow accumulation). Again the data should be global and of sufficient duration (> 10 years) to derive climatological statistics. Both conventional and remotely sensed snow data are required. Recent work by Hölze (1994) in the upper Engadine Valley, Switzerland, shows that the bottom temperature of winter snow cover is a good indicator of permafrost distribution if combined with digital elevation model data. This would potentially be an additional parameter for inclusion in the list shown in Table 1, for change detection.

Table 1. Results of priority setting for key permafrost variables.

Parameters	APPLICATIONS				
	Process understanding	Engineering design	Model validation	Change detection	Impact evaluation
Geometry					
Permafrost extent	M [*]	H	H	H	M
Permafrost thickness	M	H	M	H	M
Active layer thickness	H	H	M	H	H
Ground ice extent	H	H	M	M	H
Thermal State					
Temperature<DZAA [†]	H	H	H	H	H
Temperature>DZAA	M	M	M	H	L
Thermal conductivity	H	H	M	L	L
Composition and Properties					
Moisture content	H	H	M	H	H
Chemical composition					
Soil	M	H	L	M	M
Water or ice	M	M	L	M	M
Trace gases	M	L	M	M	M

* H, M, L = High, Moderate, and Low priority.

† Depth of zero annual amplitude.

4.1.2. System Monitoring and Change Detection

Permafrost data can be used for Earth System monitoring and change detection in several ways: the data may be used alone (e.g., measuring the areal extent of permafrost and ground ice content, or development of thermokarst features); they may be used as an integrator of various climate parameters (e.g., active layer thickness); and they may be used to support trends noted in the data sets (e.g., changes in shallow ground temperatures may be used to support trends noted in regional meteorological data).

For monitoring purposes the most important consideration is the length of record of the data set and its internal consistency. As with any meteorological data set these require a length of record sufficient to extract long-term trends from the short-term variability, and they require sufficient metadata to assess the data set history and to separate natural changes from artifacts introduced by the data collection process. While global data are less critical for change detection than they are for model validation, data sets have to be available from enough regions to demonstrate that observed changes are globally significant.

Interactions and feedbacks between permafrost and climate (mainly via greenhouse reinforcing as a consequence of thawing of frozen organic matter) involve very long time scales, but alterations due to continued or even accelerated warming could be dramatic over vast areas (affecting land subsidence, coastal erosion, drainage patterns, slope stability, etc.). Borehole temperatures in ice-supersaturated and, hence, impermeable permafrost provide extremely clear signals of secular warming trends (Lachenbruch *et al.*, 1988) and recently accelerated temperature increases. Commercial boreholes have been drilled in connection with Arctic oil exploration, but these boreholes are not always available for long-term temperature measurements. A few research boreholes of limited depth have recently been drilled at high altitude/low latitude sites in Canada, China, and the European Alps, and their records should become part of the proposed GGD.

4.1.3. Process Studies/Process Model Development and Verification

Data priorities for process studies and models are less easily defined than they are for the Earth System Models or for climate monitoring. Virtually any data at any temporal or spatial resolution are potentially useful for empirical analysis, as inputs for process models, for model parameterization, or for model validation. In this case, the greatest priority would go to those data sets that comprise a suite of co-located measurements of several related parameters.

4.2. Guidelines for Prioritizing Data Sets

The guidelines reported by Crane (1993) have been modified slightly, but essentially adopted for the present purpose.

- 1) The five areas of application – process understanding, engineering design, model validation, change detection, and impact evaluation – have equal priority. Other applications should not be excluded but they would receive a lower priority level.
- 2) Only data sets with a certain minimum level of accompanying metadata should be considered. Metadata do not have to be complete, but enough information is needed to determine the exact nature of the data collected, the location and period of coverage, and to evaluate the data reliability. It should be recognized, however, that the reliability may not be immediately obvious. The evaluation of reliability often comes from the research process and through comparison with other data sets. What defines a minimum level of metadata will vary from application to application.
- 3) Data in danger of being lost should get a higher priority for data rescue. Priority is determined by length of time before the data set is likely to be lost. Higher priority should also go to data that, while in no danger of being lost, are presently inaccessible to the user community.

- 4) Data quality is an important consideration, but it is difficult to quantify. Requirements for data accuracy and reliability will vary from application to application. A lower quality may be more acceptable for unique data sets compared to those for which other alternatives are available. The decision on an acceptable data quality (as with the decision on what constitutes sufficient metadata) will rest with the Data Center manager.
- 5) Having satisfied the previous criteria, data sets may be prioritized according to data set attributes such as coverage, duration, frequency, and cost.

The results of the prioritization exercise conducted at the workshop for five key applications are summarized in Table 1 (p.8). As the table illustrates, the requirements in these categories vary for each application. Comments on these rankings are welcomed as we anticipate that the priorities will be reassessed as the GGD process develops.

It was noted that data would be especially useful for model validation if they coincided with the 10-year (1979 to 1988) Atmospheric Model Intercomparison Project (AMIP) period used for GCM intercomparisons. High priority would also go to data sets having a spatial coverage and duration that match other extensive data collection/analysis programs such as GEWEX, ACSYS, ITEX, etc.

Further discussion following this priority-setting exercise led to the conclusion that permafrost thickness should also have been divided into shallow and deep (or thin and thick), perhaps using the same depth division as used for temperature. Greater emphasis might also have been given to trace gas composition, especially methane. Site descriptions (location, geology, vegetation, ownership), and metadata (sampling techniques, equipment used, precision, post-processing) are included in the data description and were not prioritized separately.

Ultimately, decisions about which data sets to rescue will depend on the cost of the rescue operation as well as scientific importance. A cost-benefit analysis should be part of any decision making process – where costs are high and the importance of the data set relatively low, the data set would automatically receive a low priority. Low costs, on the other hand, should not automatically raise the priority of the data set. It is likely that decisions regarding costs can only be made by considering the data rescue in terms of opportunity cost (i.e., the loss of other data sets that might alternatively have been rescued). Again, this is a decision that would appear to rest with the data center. Recognizing the limited resources available for data retrieval tasks, the workshop also developed a number of recommendations for guiding data rescue, archiving, and information management. Other, more technical and procedural results of the workshop included the discussion of formats for data set information and the preparation of drafts of a statement of protocol for data management, acquisition, and dissemination (see Appendix 4).

4.3. High Priority Data Sets

The meeting participants discussed several candidate data sets that should receive priority treatment, following the presentations by the Russian specialists (see Appendix 3) and brief status reports by other national representatives. It was decided that the following data should be targeted for early attention in the Global Geocryological Database:

- digital point values of permafrost thickness, ground temperature and ground ice content that were used in preparing the 1:10 million Circumarctic Map of Permafrost and Ground Ice Conditions (Heginbottom et al., 1993; Brown et al., in press); data from some 100 boreholes in Russia also incorporated in the same map;
- data on temperatures and moisture content in soil layers at standard depths at selected stations in Russia (Table 2);
- metadata on national and regional maps of permafrost published in Russia (see Appendix 5);
- a directory of institutions and individuals in the Former Soviet Union holding major collections of permafrost data (See Appendix 6).

**Table 2. SELECTED RUSSIAN METEOROLOGICAL STATIONS.
SOIL TEMPERATURE DATA (PILOT PROJECT).
STATIONS WITH OBSERVATIONS AT DEPTHS > 2.4 m**

STATIONS	LOCATION	PERIOD OF RECORD
Aleisk	Not Available	1947 - 1965
Eleckay	Not Available	1955 - 1965
Erbogachon	61°16' N - 108°01' E	1953 - 1970
Irkutsk	52°16' N - 104°21' E	1958 - 1970
Ishim	56°06' N - 69°26' E	1947 - 1965
Kazach'ye	70°75' N - 136°26' E	1952 - 1965
Khantynansiysk	61°00' N - 69°10' E	1961 - 1965
Komsomolskiy	69°17' N - 172°70' E	1963 - 1965
Markovo	64°41' N - 170°25' E	1948 - 1965
Olëkminsk	60°36' N - 120°36' E	1948 - 1965
Oimykon	63°47' N - 142°50' E	1949 - 1965
Ostrovnoy	58°58' N - 163°57' E	1951 - 1965
Russkaya Polyana	53°48' N - 73°54' E	1954 - 1965
Salekhard	66°55' N - 66°67' E	1949 - 1958
Seimchan	62°55' N - 152°25' E	1948 - 1965
Sidorovsk	66°35' N - 82°18' E	1953 - 1965
Skovorodino	54°60' N - 123°53' E	1953 - 1965
Srednekolymsk	62°27' N - 153°35' E	1931 - 1965
Syktyvkar	61°42' N - 50°45' E	1949 - 1965
Tarko-Sale	64°55' N - 77°50' E	1949 - 1965
Tobolsk	58°20' N - 68°23' E	1947 - 1965
Troitsko-Pechorsk	62°70' N - 56°22' E	1949 - 1965
Ugut	60°31' N - 74°07' E	1947 - 1965
Ust-usa	66°00' N - 57°00' E	1947 - 1965
Verhoyansk	67°58' N - 133°50' E	1932 - 1965
Vitim	59°26' N - 112°35' E	1931 - 1965
Yakutsk	62°02' N - 129°72' E	1949 - 1965
Zhigansk	66°75' N - 123°33' E	1948 - 1965

4.4. General Recommendations

Given limited resources we recommend that:

- 1) Immediate efforts should be made to begin, or to assist with, the rescue of data sets pertaining to the high priority variables listed in Table 1.
- 2) The IPA should consider a follow-up meeting in mid-1995 that would review the data rescue procedures and focus on identifying other high priority data sets for the next stage of the rescue effort.
- 3) This workshop focused almost entirely on the recent time period. We should not lose sight of the fact that the best verification of an Earth System Model's ability to predict climate change is to run the model for past climate states. Paleoclimatic data are vital for this approach, and a focused effort should address the question of permafrost data rescue for paleoclimate applications in this context. This can be done in cooperation with other programs such as the International Geosphere-Biosphere Program's Past Global Changes Project (PAGES), the UNESCO-International Union of Geological Sciences CLIMEX (Climatic Extremes of the Past), mapping project, and the ARCSS PALE program (PALE Steering Committee, 1993).
- 4) The order of processing for rescuing data sets should not automatically start with the oldest records and work forward (unless the oldest records are deteriorating at a rate that makes this approach necessary), nor should it start at the present and work backwards. As was noted in the criteria for data set prioritization, the most useful data, such as those that coincide temporally with other large-scale experiments or overlap with the introduction of new satellite systems, should receive early attention.
- 5) The question of data rescue, in general, and permafrost data rescue in particular, should be included on the agenda of the meetings of the IPA Executive Committee and Council and special sessions at the five-yearly international conferences.
- 6) Data rescue should also be placed on the agenda of the International Commission of Snow and Ice, and other international organizations with common interests.

5. Next Steps

Several useful data sets have been compiled and released on CD-ROM; an example is the Alaska North Slope Data Sampler containing soil and permafrost data distributed by NSIDC (1994). It would be useful if a general cryospheric data set that included permafrost data could be made available in this form. This could include gridded data contained in the permafrost map of the northern hemisphere, selected borehole temperature data from Russia and Canada, and measurements of active layer characteristics for circum-Arctic sites.

The Data and Information Working Group anticipates that the IPA, through the Council and other Working Groups, will encourage the implementation of the GGD in the following ways:

- Develop an inventory of relevant national data sets (see Appendix 5).
- Compile approved data set descriptions into master directories, accessible to users. The International Arctic Environmental Data Directory (ADD) project is a possible means of facilitating user access to GGD (Appendix 7).
- Retrieve priority data sets, and archive them in standard formats at National Database Centers or GGD nodes.
- Make the databases available to users through appropriate distribution media (diskettes, CD-ROM, hard copy) or on-line via FTP.
- Promote user-oriented analytical, modeling and mapping information products, based on NGD or GGD data sets.
- Concurrently develop funding proposals to national and international, governmental and non-governmental funding sources.

The Data and Information WG invites all IPA member countries, working groups and interested individuals, wherever they are, to participate in and contribute to the GGD project. As a first step this requires completion of a data set description on the form being distributed with the December 1994 issue of *Frozen Ground* to over 2,000 readers (see Appendix 1). Readers of *Frozen Ground*, are encouraged to complete and return the form to the WDC-A for Glaciology or their National Representative at their earliest convenience. The WG plans to convene a three-day meeting and workshop in Potsdam, Germany, before the XIV International Congress of the International Union for Quaternary Research, in Berlin. It will report on the status and results of GGD activities to the IPA Council at its 4-5 August, 1995 meeting.

The participants recommended the following strategic targets and time table for GGD implementation.

A. RESEARCH AND DEVELOPMENT

- A1** Review and refine the GGD Parameter Priorities List (through *Frozen Ground*) (6 months)
- A2** Refine the GGD Data Structure
(coordinate through the groups in Southampton and Moscow) (6 months)
- A3** Converge on agreed GGD DIF (in consultation with USGS) (9 months)
- A4** Combine A1-A3 into a first Draft of IPA "Guidelines for Geocryological Data Management" with metadata guidance (12 months)

B. DATA SETS

- B1** Compile first version of Inventory of Candidate Data Sets (6 months)
- B2** Using refined List A1, develop from B1 a List of Rescue Priorities for IPA (9 months)
- B3** Define technology. (Scan v. Automated digitization v. Manual digitization) (6 months)

C. INFORMATION SYSTEM

- C1** Continue work on IPA GGD database pilot, initially using Barrow and CPM data (6to12months)
- C2** Develop proposals for pilot GGD Dissemination System (media and management) (6to12months)
- C3** Develop proposals for high level derivative products (9to15 months)
 - Derivative data
 - Analysis and monitoring
 - GGM input data
 - Permafrost bulletin
 - Inputs to the Intergovernmental Panel on Climate Change
 - Flagship data sets

6. References

- Barry, R.G. 1988. Permafrost data and information: status and needs. In: *Fifth International Permafrost Conference Proceedings*, Vol. 1. K. Senneset, ed., Tapir Publishers, Trondheim, pp. 119-122.
- Barry, R.G. and A.M. Brennan. 1993. Towards a permafrost information system. In: *Permafrost, Sixth International Conference Proceedings*, Vol. 1. South China University of Technology Press, Wushan, Guangzhou, pp. 23-26.
- Brennan, A.M. and R.G. Barry. (eds.) 1989. Permafrost Data Workshop. In: *Glaciological Data. Report GD-23*, World Data Center-A for Glaciology, Boulder, Colorado, pp. 107-127.
- Brown, J., O.J. Ferrians, Jr., J.A. Heginbottom and E.S. Mel'nikov, (eds.). in press. *Circumarctic Map of Permafrost and Ground Ice Conditions*. (1:10 million) International Permafrost Association.
- Crane, R.G. (Ed.) 1993. Workshop on Cryospheric Data Rescue and Access. In: *Glaciological Data. Report GD-25*, World Data Center-A for Glaciology, Boulder, Colorado, pp. 274-294.
- Heginbottom, J.A., J. Brown, E.S. Mel'nikov and O.J. Ferrians, Jr. 1993. Circumarctic map of permafrost and ground ice conditions. In: *Permafrost: Sixth International Conference Proceedings*. Vol. 2, South China University of Technology Press, Wushan, Guangzhou, pp. 1132-1136.
- Hölze, M. 1994. Permafrost und Gletscher im Oberengadin. *Mitteilungen VAW/ETH* 132, ETH, Zürich, 121 pp.
- IASC. 1994. *The 1994 IASC Meeting Report*. International Arctic Science Committee, Oslo, Norway, 35 pp. and Appendices.
- Koster, E.A., M.E. Nieuwenhuijzen and A.S. Judge. 1994. Permafrost and Climatic Change: An Annotated Bibliography. *Glaciological Data. Report GD-27*, World Data Center-A for Glaciology, Boulder, Colorado, 94 pp.
- Lachenbruch, A.H., T.T. Cladouhos and R.W. Saltus. 1988. Permafrost temperature and the changing climate. In: *Fifth International Conference on Permafrost Proceedings*. Vol. 3. Trondheim, Norway, Tapir Publishing. pp. 9-18.
- McCauley, L.L. and M.F. Meier (eds). 1991. *Arctic System Science: Land / Atmosphere / Ice Interactions. A Plan for Action*. ARCUS, Fairbanks, Alaska, 48 pp.
- Nelson, F.E., A.H. Lachenbruch, M.-K. Woo, E.A. Koster, T.E. Osterkamp and M. Gavrilov. 1993. Permafrost and changing climate. In: *Permafrost, Sixth International Conference Proceedings*. Vol. 2, South China University of Technology Press, Wushan, Guangzhou, pp. 987-1005.
- NSIDC. 1994. *North Slope Alaska Data Sampler*. ARCSS LAII, 1 CD-ROM, National Snow and Ice Data Center, University of Colorado, Boulder.
- PALE Steering Committee. 1993. Paleoclimates of Arctic Lakes and Estuaries. Research Protocols.
- Waelbroeck, C. 1993. Climate-soil processes in the presence of permafrost: a systems modeling approach. *Ecological Modelling* 69(3-4), pp. 185-225.

7. AGENDA

Objectives

1. Establish priorities for the recovery of permafrost and frozen ground data
2. Develop the implementation plan and procedures for the data recovery, storage and dissemination
3. Review results of pilot projects and approaches to database structure and utilization

Agenda – Day 1, Thursday, 3 November 1994

- | | | |
|-------|---------------------------------------------------------------------------------------------------------------------------------------------|----------------------|
| 09:00 | Welcome and Introductions | (Flaate, Barry) |
| 09:15 | Review Purpose and Program of Meeting, Accept Agenda | (Barry, Heginbottom) |
| 09:30 | GGD project – Status Report, Summary of Southampton Meeting | |
| 10:00 | Break | |
| 10:15 | Criteria for data prioritization
(See Crane, 1993) | (Barry, Brown) |
| 13:00 | Lunch | |
| 14:00 | Data set identification and description, Data availability
(existing Directory Interchange Format [DIF] and Master Directory activities) | (Participants) |
| 16:00 | Break | |
| 16:15 | Presentations of on-going GGD-NGD activities
(Russia, US, Canada, China, Fennoscandia, Alpine Europe, Japan, etc.) | (Participants) |
| 17:30 | Adjourn | |

Day 2, Friday, 4 November 1994

- | | | |
|-------|-------------------------------------------------------------------------------|------------------|
| 08:30 | Models and structures:
GeoData results of pilot projects including GLOCOPH | (Clark, Branson) |
| 10:00 | Break | |
| 10:15 | Working sessions: Priorities and options | (Participants) |
| 12:30 | Lunch | |
| 13:30 | Working session: Data distribution and archives | (Participants) |
| 15:00 | Break | |

- 15:15 Working session: Activities, options, schedules, funding, responsibilities (Participants)
- 17:30 Adjourn

Day 3, Saturday, 5 November 1994

- 08:30 Working session: Report preparation
- 10:00 Break
- 10:15 Plenary session: Review progress, Next meeting
- 12:00 Closure
- 12:30 Lunch
- 13:30 IPA Data Working Group Business Meeting
- 16:30 Closure

8. PARTICIPANTS

Dr. Roger G. Barry

WDC-A for Glaciology
University of Colorado
Boulder, Colorado 80309-0449 USA
Phone: 1 (303) 492-5488
FAX: 1 (303) 492-2468
e-mail: rbarry@kryos.colorado.edu

Dr. Julia Branson

Geodata Institute
Department of Geography
University of Southampton
Southampton SO17 1BJ, United Kingdom
Phone: +44 (1703) 594488
FAX: +44 (1703) 592849
geodata@soton.ac.uk
e-mail: j.branson@soton.ac.uk

Dr. Jerry Brown

International Permafrost Association
Post Office Box 9200
Arlington, Virginia 22219-0200 USA
Phone: 1 (703) 525-3136
FAX: 1 (703) 525-3136
e-mail: jerrybrown@igc.apc.org

Professor Chen Xianzhang

WDC-D for Glaciology
Lanzhou Institute of Glaciology and
Cryopedology
Chinese Academy of Sciences
Lanzhou 730 000, China
Phone: +86 (931) 8826725
FAX: +86 (931) 8885241
Telex: 72008 IGGAS CN
e-mail: ggcheng@bepc2.ihep.ac.cn

Dr. Michael J. Clark

Geodata Institute
Department of Geography
University of Southampton
Southampton SO17 1BJ, United Kingdom
Phone: +44 (1703) 594488
FAX: +44 (1703) 592849
geodata@soton.ac.uk
e-mail: mjc@soton.ac.uk

Dr. Kaare Flaate

Norwegian Directorate of Public Roads
Post Office Box 8142 DEP
N-0033 Oslo, Norway
Phone: +47 (22) 073 900
FAX: +47 (22) 073 444

Dr. Masami Fukuda

Institute for Low Temperature Sciences
Sapporo N19W8 Japan
Phone: +81 (11) 706-5492
FAX: +81 (11) 706-7142
e-mail:
masami.fukuda@LT.hines.hokudai.ac.jp
mfukudatj@LT.hines.hokudai.ac.jp

Dr. Odd Gregersen

Norwegian Geotechnical Institute
Sognsveien 72
3930 Ulleval Hageby
N-0806 Oslo, Norway
Phone: +47 (22) 02 3000
FAX: +47 (22) 23 0448

Dr. Wilfried Haeberli

VAW/Glaciology
Gloriastrasse 37/39
ETH-Zentrum
CH-8092 Zurich, Switzerland
Phone: +41 (1) 632-4093
FAX: +41 (1) 632-1192
e-mail: haeberli@vaw.ethz.ch

Mr. J. Alan Heginbottom

Geological Survey of Canada
601 Booth Street
Ottawa
Ontario, Canada K1A 0E8
Phone: +1 (613) 992-7813
FAX: +1 (613) 992-2468
e-mail: heginbottom@gsc.emr.ca

Dr. Arne Instanes

SINTEF
N-7034 Trondheim, Norway
Phone: +47 (73) 59 3000
+47 (73) 58 8135
FAX: +47 (73) 59 5340
e-mail: arne.instanen@geotek.sintef.no

Dr. Rostislav M. Kamensky

Permafrost Institute
Russian Academy of Sciences
677 018 Yakutsk, Russia
FAX: +7 (095) 230-2919
e-mail: lans@im22an.infonet.yakutia.su

Dr. Suzanne Lacasse

Director
Norwegian Geotechnical Institute
Sognsveien 72
3930 Ullevål Hageby
N-0806 Oslo, Norway
Phone: +47 (22) 02 3000
FAX: +47 (22) 23 0448

Dr. Marina O. Liebman

Federal Center for Geoecological Systems
Ministry of Environmental Protection and
Natural Resources
Zvezdny Boulevard 5/2-28
129085 Moscow, Russia
Phone: +7 (095) 215-8666
FAX: +7 (095) 818-1254
e-mail: nromanovsky@glas.apc.org

Dr. Evgenii S. Mel'nikov

VSEGINGEO
142425 Zeleny Village
Noginsk District
Moscow Region, Russia
Phone: +7 (095) 529-1101 (office)
+7 (095) 529-5101 (ext. 2586)
+7 (095) 287-4307 (home)
e-mail: emelnikov@glas.apc.org

Dr. Frederick E. Nelson

Department of Geography and Planning
State University of New York
Social Science 147
Albany, New York 12222 USA
Phone: +1 (518) 442-4469
FAX: +1 (518) 442-4867
e-mail: fnelson@cnsvox.albany.edu

Mr. Douglas Posson

U.S. Geological Survey
Box 25046 (Mail Stop 801)
Denver Federal Center
Denver, Colorado 80225-0046 USA
Phone: +1 (303) 236-4944
FAX: +1 (303) 236-8888
e-mail: dposson@isdminl.wr.usgs.gov

Professor Nikolai N. Romanovskii

Geocryology Department
Faculty of Geology
Moscow State University
119899 Moscow, Russia
Phone: +7 (095) 939-1937 (office)
+7 (095) 133-2668 (home)
FAX: +7 (095) 932-8889
+7 (095) 318-1254
e-mail: nromanovsky@glas.apc.org

Professor J.L. Sollid

University of Oslo
Department of Physical Geography
Post Office Box 1042 Blindern
N-0316 Oslo, Norway
Phone: +47 (22) 85 5936
FAX: +47 (22) 85 7230
e-mail: j.l.sollid@geografi.uio.no

APPENDIX 1

IPA GLOBAL GEOCRYOLOGICAL DATABASE INVENTORY

The International Permafrost Association is conducting a survey of available data on past and current investigations of permafrost, seasonally frozen ground and periglacial conditions and related laboratory studies. An electronic mail form is also available. You may submit more than one form if data types differ substantially. Results of this survey will be reported in *Frozen Ground* and other relevant publications. Please complete this form and return it to your IPA national representative with a copy to the World Data Center-A for Glaciology, Attn: Claire Hanson, Campus Box 449, University of Colorado, Boulder, Colorado 80309-0449, U.S.A. **Forms and instructions are also available electronically from: hanson@kryos.colorado.edu.**

Name of data set			
Principal Investigator			
Name			
Address			
Tel	Fax	E-mail	
Data compiler/author			
Name			
Address			
Tel	Fax	E-mail	
Coverage			
Study location (region/country)			
Latitude (south to north)		to	
Longitude (east to west)		to	
Period of investigation			

Summary: Give brief details of site, collection technique, measurement frequency, and quality. Indicate what ancillary data on site, climate, data processing and metadata are available or can be reliably obtained (with sources).

(Attach additional sheets if necessary)

Current storage medium: CD-ROM / Spreadsheet / Word processor / Database / Data centre (give name) / Paper

Are your data at risk of being lost? YES / NO

Bibliography (Published and unpublished reports about this data set; attach additional sheets if necessary)

Key Words: (Maximum of 10 such as Active layer, Permafrost thickness, Temperature, Moisture content, Ice content, Chemistry, etc.)

APPENDIX 2

GLOBAL GEOCRYOLOGICAL DATABASE: SUGGESTION FOR DATA STRUCTURE

Julia Branson and Mike Clark
GeoData Institute
University of Southampton
Southampton, United Kingdom

Database Prototype

A prototype database has been prepared to illustrate how the development of the GGD database could be approached. The development of the prototype has assumed that the end product should be a tool to facilitate data manipulation and analysis (see figure 1) rather than simply a method of data storage. The prototype as presented is a developers' version and has no user front end and only limited functionality compared to a full production system. Two different types of data set have been entered into the database: 1) the Barrow data set comprising soil moisture, temperature, etc., taken at multiple points and plots over a number of years and 2) the data used to compile the Canadian permafrost map (See Heginbottom *et al.*, 1993).

This report outlines issues that should be addressed during the development of the production database, describes the software used and discusses three different possible forms of data structure.

Software

For the prototype the data have been stored in a relational database system. A relational database is one in which the data are kept in several related tables. Each table contains data from a particular aspect of the data set. Tables are linked to each other using unique keys: for example, in a library data set the borrowers and book detail tables could be linked by a unique book number.

Advantages of relational structure over storing data in individual flat files include:

- 1) data can be queried to obtain items in the database which select data fulfilling a requirement either from many data sets or just particular items from one data set. Data can be output from queries as a text report or as an ASCII file that could be transferred into a spreadsheet or other package;
- 2) data redundancy (duplication) is reduced and thus storage requirements are reduced.

A disadvantage, however, is that some data manipulation may be required to input the data and fit it into the table format, particularly if it is presently held in a digital format.

The system used for the prototype is Paradox by the Borland Corporation. This package has been used by many other international projects, such as the North American and European pollen databases and the global palaeoflood database, and is relatively inexpensive (not more than about \$200). The advantage of this system over ones such as ORACLE is that graphical images can be easily incorporated into the product and there are also facilities for producing simple graphs within the database interface. A disadvantage, however, is that Paradox does not support structured query language (generally regarded as the industry standard language for relational databases), although this may not affect the end-user. For the production database it may be appropriate to hold the data on a central robust large database system and to transfer it to a more user-friendly system for distribution.

GLOBAL GEOCRYOLOGICAL DATABASE DATA APPLICATIONS
ONE DATABASE — MANY USES

USER REQUIREMENTS




Figure 1. An illustration of the diverse user requirements of the Global Geocryological Database (GGD): research versus operational and science versus engineering design. The requirements range from access to archived 'raw' data to standardized/transformed data.

Access to the database

It is envisaged that access to the database could be either on-line (via INTERNET), thus allowing the user to perform interactive queries on the main database, or distributed on a CD-ROM that contains simple querying and visualization software. (The disadvantage of the latter is that the data that the user accesses will not necessarily be the most up-to-date version as available from the main database.) Either distribution method would allow users to download data to their own software.

The majority of database systems have the facility to allow restricted access to the database. Thus users may be requested to register to use their data on-line. Database security will restrict the users to only viewing and downloading the data and they will be unable to change the data. A sample of a database protocol is attached as Appendix 4.

Data structure

A possible data structure for the GGD is shown in Figure 2. This structure is most suitable for the representation of point data. For the storage of non-point data (either linear or area) it may be appropriate to link the database to a Geographical Information System.

The tables relate to the following:

The data in the database have been initially classified by data set and further details are given in the **Data set** table. This table contains similar information to other Directory Interchange Format (DIF) flat files and should be designed when the DIF for the GGD has been finalized and would include details of spatial and temporal coverage, keywords and a brief description. This is linked by publication number to the **Bibliography** table that contains references to the reports/papers from which the data were taken or which refer to the data and other relevant articles. It is not suggested that a complete permafrost bibliography should be maintained as this is available elsewhere.

The data set table also connects to the **Researchers** table which gives the name, address, e-mail, etc. of the person/people who submitted the data to the database and who should be contacted regarding the data.

The data are held in a hierarchical structure. Initially they have been split into sites (in the case of the Barrow data set a site is an individual plot). The **Site** table gives details of the site name and a unique site key. Locational information (latitude, longitude, elevation) is described in the **Location** table (if data are referenced to a smaller scale then they can be recorded at the Record level (see below)), and a longer text description of the site is in the **Site comments** table.

The data are then classified by what has been termed a **Record**, in the case of the Barrow data a record is an individual core (or point). The **Record** table gives the name and type of each record.

Representation of data

For ease of use of the database, to ensure that different data sets can be easily compared, and to avoid misrepresentation, it may be appropriate to translate all data to a common mensuration (probably the International Standard). It is important that a copy of the raw data is also held in the database, however, although it is recommended that this should only be available to privileged users.

Storage of spatially and temporally variable data at the record level

Storage of data that have been collected at different spatial and temporal intervals is a complex matter and three different alternatives have been explored here (see Figures 2, 3, and 4). The following tables are common to the three types of structure above: Data Set, Site, and Record tables.




Figure 2. Candidate data structure for GGD: Alternative 1. This is especially suitable for point data.




Figure 3. Candidate data structure for the GGD: Alternative 2.




Figure 4. Candidate data structure for the GGD: Alternative 3.

Alternative 1) The data are held in just one main table; within this the different depths within a core have unique sequence numbers. The table contains the following:

Type of data (e.g., soil moisture content, snow depth, permafrost extent)

Value — this can be either numerical (e.g., 50%) or text (e.g. continuous)

Depth at which the sample is taken

Date — this is stored in a year field, a day/month field and an AD/BP date field.

The disadvantage behind this method is that there is a considerable amount of data redundancy, as the depth, dates, etc., have to be repeated for each sequence in each record. The advantage, however, is that queries of the data are relatively simple and quick to construct and, also, only a limited amount of manipulation is usually required to transfer the data from an original digital flat file to the database table.

The other two alternatives are more fully relational, and thus reduce the amount of data redundancy (and the amount of storage required). A considerable amount of data manipulation would be required, however, to transfer the digital data into the table structure. Queries are also complex to construct.

Alternative 2) The location and the date values are kept in individual tables so that many sequence numbers can refer to them, and the tables linked with key numbers (Figure 3).

Alternative 3) Each data type is stored in an individual table (Figure 4). This solution is not recommended as a large number of tables would be required to cover the wide range of possible data. Querying would also be very complex.

Other database tables

If the date has been determined by a dating method (e.g., radiocarbon) or has been inferred by stratigraphic inference then further details are given in the **Date information** table. This gives the errors, sample and laboratory numbers, dating method, etc., associated with the date.

If various statistics have been collected for a particular item of data (e.g. maximum, minimum temperature) then the details are given in the **Statistics** table which is linked to the **Data** table.

APPENDIX 3

RUSSIAN CONTRIBUTION TO THE GLOBAL GEOCRYOLOGICAL DATABASE

N.N. Romanovskii and M. Leibman
Moscow State University
Moscow, Russia

According to the goals and objectives announced in the letter distributed by the IPA Data Working Group on September 26, 1994, Russian participants in the meeting suggest that the approaches developed for the Russian National Geocryological Database (NGD) be applied to the Global Geocryological Database (GGD). Geocryological databases represent one component of the much broader cryospheric database and should be compatible with the latter. At the same time, permafrost and geocryological conditions differ fundamentally from other components of the cryosphere, such as snow and ice. As a geological feature, permafrost is much more inert to external influences. That is why it is more sensitive to medium (10s of years) and long-term (100s-1000s of years) climate fluctuations than to short-term (interannual years) and seasonal ones.

The main manifestations of permafrost changes are concentrated at and just below the earth's surface where external climatic and human-induced factors are interacting with permafrost features. This interaction results in specific cryogenic (periglacial) processes and phenomena. Nevertheless permafrost is not only a geological but a climatic feature as well, and hence several processes having a seasonally variable character are inherent in the geocryological conditions of the permafrost/atmosphere/hydrosphere interfaces.

Another specific feature of permafrost in comparison with surficial features is that some methods of observation are not applicable to the study of the permafrost temperature regime, such as repeated aerial and satellite photography. This method gives little information on the upper layers of permafrost and virtually none for the deep layers of permafrost. Only those features that have a geological origin but surficial expression, such as icings, can be studied like glacial features. Data obtained when studying permafrost thus differ from that obtained for snow and ice, and the data processing and generalization depend upon the existence of stability in the medium. The main method of collecting data and of mapping permafrost in Russia is linked to the landscape type. This involves two main procedures for acquiring data, depending on the degree of generalization. The first is archiving maps with generalized information on each category (the maximal degree of generalization). The second is archiving of the coordinates of the points (boreholes, etc.) with complete information layer by layer on lithology, cryogenic features, and laboratory tests (the minimal degree of generalization, but still varying according to the depth and goals of drilling, or description).

The problem of choosing the degree of generalization also concerns data collection about the permafrost regime. At meteorological and agricultural stations, climatic and active layer records are averaged for daily, weekly, monthly, annual and multi-year intervals. Different subjects need different degrees of generalization. Some cryogenic processes and active layer studies need daily information. Such processes as thermoerosion and thermoabrasion need seasonal information, frost heave and thermokarst need long-term information. There are several hundred weather and agricultural stations in the Russian permafrost zone and in the zone of the deep seasonal freezing (for example, a seasonally frozen layer of about 3-5 m deep is characteristic of the southern part of West Siberia). At least one hundred of them measure the depth of thawing or freezing and the temperature of active layer at 8 levels in the profile (0, 20, 40, 80, 120, 160, 240, and 320 cm). There is a 60-year long period of observations at one of the stations. In this case we have 35,000 readings just for the temperature of the active layer; moisture content is also measured there. Should all these records be digitized or should we choose a selection based on some criteria?

The data obtained at the meteorological stations, agricultural stations, and steady-state permafrost stations during investigations for construction have different content, different techniques

of monitoring, laboratory testing and generalization. This makes it difficult to work out unique criteria for data selection.

To summarize:

- Permafrost as a geological feature has parameters that are stable or changing very slowly with time. They are important to the fundamental understanding of permafrost and its relationships with "... climate, process, material and morphology ..."
- At the same time permafrost as a climatic feature is characterized by short-term fluctuations reflecting the changes in climate and environments and in its turn influencing them.
- This duality results in heterogeneity of the database structure. We suggest that GGD should consist of several blocks with different internal structures depending on their subject. Neither present geological nor geographical database structures are acceptable. The structure of the geocryological database combines several blocks of features and parameters on one side and regime observations on the other side. Four main criteria and several specific criteria are proposed.

We suggest the following main criteria for data prioritization:

1. **Relevance** — the use of relations between the geocryological data and important problems such as global climatic change, environmental protection, and development of useful mineral deposits in the Arctic.
2. **Availability** — the possibility of obtaining data with minimal time and use of funds.
3. **Completeness** — the maximal set of information at one point.
4. **Spatial distribution** — the acquisition of a uniform distribution of the points throughout the territory.

The last two points are, to some extent, contradictory, so an optimal combination should be found.

Specific criteria should be attached to specific data sets. Some permafrost elements, such as icings, have features of a stochastic character (analogous to glaciers) and icing data can be generalized only for situations with adequate local data. Hence, the fourth criterion cannot be used in cases where statistical analyses are to be performed.

Most permafrost data are concentrated in regions of active economic development where they are often not accompanied by the collection of weather data. In this case, high quality analyses of permafrost data are impossible. In this situation weather records play a more important role than the number of permafrost parameters monitored and the third criterion is not considered.

The most numerous data on the active layer are concentrated at agricultural stations but they are situated mostly to the south of the southern limit of the permafrost zone. Here the main criterion should be the existence of paired data (on hills and in valleys) and at an intermediate latitude; the latter refers to a location on some transect from the southernmost to the northernmost stations within different geographical regions.

Most data on ground temperature refer to the depth of zero annual amplitude. Often there are single measurements in each borehole, but sometimes there are repeated measurements although not necessarily on the permafrost regime. Those cases are most important in considering the stability of the permafrost thermal regime. Here the criterion of the time of recording is less important. The main criteria for monitoring borehole temperature are data extending below the base of permafrost, or at

least exceeding of the depth of zero annual amplitude, measuring the temperature gradient in permafrost, and monitoring processes in the same location (again data at paired elevations).

In order to unravel the problem of dividing the criteria, we suggest a list of informational blocks. Each block in the list would have a different structure and criteria for data acquisition.

- Block I —** Geocryological maps.
- Block II —** Geocryological conditions existing in boreholes, excavations, and description points.
- Block III —** Cryogenic (periglacial) processes.
- Block IV —** Observations on permafrost regime.
- Block V —** Inventory of metadata sets.

The inventory of metadata sets includes:

- Set 1 —** A list of permanent weather stations, agricultural stations and permafrost stations.
- Set 2 —** A list of investigations, design institutions and construction companies involved in Arctic and subarctic development.
- Set 3 —** A list of published and unpublished sources of information.
- Set 4 —** A list of existing digital or hard copy local databases (with a specific standard description of each database).
- Set 5 —** A list of important unpublished geocryological maps.

We have started to develop a Russian National Geocryological Database (NGD) taking account of the following:

- the NGD should be compatible with the GGD, and
- the NGD should be compatible with existing local databases. This is why we started by planning the NGD structure. An initial version was reported at the summer meeting 1994 in Southampton. Now we are presenting the version that has been accepted for the NGD use for the GGD.

The NGD is based on a GIS-program and other compatible programs of data archiving. Four separate data blocks, referred to above, have been prepared for this meeting containing different type of information:

- I —** Bibliographical description of geocryological maps both published and unpublished, including maps of permafrost conditions, cryohydrogeological maps, terrain engineering maps, ecological maps, and maps of cryogenic (periglacial) processes and phenomena.
- II —** Numerical and text information on geocryological conditions obtained in boreholes, excavations, and at description points,
- III —** Numerical and text information on cryogenic (periglacial) processes,
- IV —** Numerical information on observations of permafrost regime.

The structure of the first block is rather obvious – it is a list including information on the authors, publisher, date and place of publication, scale and legend of the map. As a completed example we present a database of published maps of the FSU, Russia and regions of Russia (Appendix 5).

The second block includes practically all the information obtained in the boreholes and other points having coordinates on digitized maps. All the information is accumulated in several subblocks and can be retrieved layer by layer, or in a combination of layers according to the given legend.

The third block provides information on the extent, characteristics and rates of cryogenic (periglacial) processes and the morphometry of the phenomena referred to the category or to the area.

The fourth block is a collection of tables including measurement data and readings at the observation points of permanent research stations. Regime information can be obtained also from the published data of meteorological and agricultural stations (up to 1978) and unpublished, but available at a reimbursable cost, after 1978.

The fifth block gives information on the existence and content of geocryological data at the regional and local levels, format, archive address, owners' names and contact telephones. The structure of the block is not yet worked out but the contacts with the owners have begun.

The main sources of the data are held at:

- Permafrost institutions and research stations of the Russian Academy of Sciences (Yakutsk, Chersky, Tiksi, Syktyvkar),
- Research stations of other institutions (All-Russian Geological Institutes – Yamal, Gydan)
- Geological Survey of Russia (throughout Russia),
- Design companies (Hydroproject – Transbaikial region, Kolyma region; Mosgyprotrans, Lengyprotrans, Sibgyprotrans – Transbaikial region, West Siberia including Yamal, South Yakutia, Fundamentproekt – West Siberia, and many others).

It is a long-term task to complete the entire database, but a start should be made so that it can be of use in the future when the main emphases may change.

Conclusions

1. We support the goals and objectives of the GGD Project of IPA. We especially appreciate reference to permafrost conditions as a basis for environmental scenario development, impact assessment, and engineering design, together with the reference to global climatic change.
2. We agree that the status of GGD should be as a non-commercial, internationally accessible database. The only reason to encourage the owners of the data to make their data accessible to the general public should be that of free access to the database. Nevertheless, we suggest including a list of owners (block V, set 4) who are ready to release their data only on a commercial basis. That will at least help to provide a list of data available from the companies who are making temperature measurements in course of extensive drilling in Russia.
3. We would like to know how the existing system will work, what will be the ways of using the international database. In what way is its public character achieved?
4. We consider it important that the placing of data in the GGD is recognized as a publication. If we can gain such an acceptance, it would help in submitting proposals to different science foundations.

We would like to stress the additional problem of translation from Russian into English while developing the Russian part of GGD. This takes extra time and funds.

The system of GGD development is currently at the starting point. To make it work we need equipment, software, and funds. The minimal amount required for equipment and software is detailed in a separate report.

APPENDIX 4

DRAFT PROTOCOLS: A POSSIBLE MODEL FOR THE GGD

The participants discussed the issue of protocols for data management and dissemination. Draft protocols prepared for the ARCSS PALE program (PALE Steering Committee, 1993) and for the Global Continental Paleohydrology Project (GLOCOPH) database were discussed. GLOCOPH is organized by a commission of the International Union for Quaternary Research (INQUA).

The GLOCOPH draft protocol, presented by J. Branson, is summarized below.

Data

- 1) Data should include both original field measurements (raw data) and the paleoenvironmental reconstructions.
- 2) Only data that are available for unrestricted use will be accepted for the database.
- 3) GLOCOPH is not liable for ensuring the accuracy of the data - this is the responsibility of the data contributor.
- 4) Data must include a time-frame, even if it is very approximate.

Data contributors

- 1) Data should be made available to the database free of charge.
- 2) A printout of the data can be sent to the data contributor, if requested, so that the entry of the data may be verified.
- 3) Contributors can request to be informed if their data are being used, for what purpose and by whom.
- 4) Data can be removed from the database at any time at the request of the contributor.

Data users

- 1) Users must be registered to use the database.
- 2) The database can be used free of charge.
- 3) The database should be referenced when referring to data obtained from it and publications by the data contributor which discuss the original data should be cited.
- 4) Users should acknowledge the contributor if unpublished data are used.
- 5) Users should send contributors reprints of publications which use their data.
- 6) Data from the database should not be passed onto a third party; all data should be taken directly from the central database.
- 7) The data can be used by non-profit-making organizations for research purposes only. Profit-making organizations may use the data only if written consent has been given by the originator of the data and the database production group.

PALE Steering Committee. 1993. *Research Protocols for PALE. Paleoclimates of Arctic Lakes and Estuaries*. PAGES Workshop Report-Series 94-1. Bern, Switzerland, pp. 35-39.

APPENDIX 5

RUSSIAN PERMAFROST MAP INVENTORY

The list of published maps includes both maps printed as separate items and those found in papers and monographs as illustrations. In the latter, references are given to the publication and page where the map is found. There are 375 maps listed; this is less than 20% of those published. Only the period 1950 to 1980s is covered by the list. The compilation is being continued and will include later publications. Irina D. Streletskaia, Ph.D. (Industrial and Research Institute for Investigations of Construction, Moscow) and Marina O. Leibman, Ph.D., Federal Center for Geoecological Systems, Moscow) are the editors. The listing is alphabetical by first author and the entries are numbered consecutively. The map scale index which follows p. 114, refers to that number.

- AUTHOR Afanasenko, V.E., Goncharov, S.V., Zaitsev, V.N.
NAME: Geocryological-hydrogeological regionalization of the territory
PUB: Moscow, Moscow University Publishers, 1981
SCALE: 1:3 500 000
SOURCE: "Natural conditions Transbaikal railway industrialization zone", p.18. In the article by Afanasenko, V.E., Goncharov, S.V., Zaitsev, V.N. p.4-20
REGION: Transbaikal
LEGEND Cryohydrogeological massif of deep discontinuous freezing, cryoartesian basins of continuous and discontinuous shallow freezing
LAT/LONG: 54°00-57°00/120°00-127°00
INSTITUTE: Moscow State University
NUMBER 00001
- AUTHOR Are, F.E.
NAME: Soviet Arctic regionalization scheme referring to thermoabrasion in shelf cryolithozone
PUB: Moscow, Nauka, 1983
SCALE: 1:37 500 000
SOURCE: "Geocryology problems"/edited by P.I. Mel'nikov, p.197. In the article by Are, F.E., p.195-201
REGION: North of the USSR
LEGEND 5 areas (characteristic in article)
LAT/LONG: 68°00-82°00/20°00-170°00
INSTITUTE: Permafrost Institute
NUMBER 00002
- AUTHOR Arkhipov, S.A., Astakhov, V.I., Volkov, I.A.
NAME: Paleogeography of West Siberian plain (sketch-map) at maximum of Late Zyriansk Glaciation (22-17 thousand years)
PUB: Novosibirsk: Nauka, 1980
SCALE: 1:15 000 000
SOURCE: Inset-map in monograph "Paleogeography of West Siberian plain at the maximum of Late Zyriansk Glaciation"
REGION: West Siberia
LEGEND Northern limit of permafrost deep thawing
LAT/LONG: 48°00-80°00/70°00-96°00
NUMBER 00003
- AUTHOR Badu, I.U.B., Vasilchuk, I.U.K., Kashperiuk, P.I., Trofimov, V.T.
NAME: Extent and contemporary tendency for development of ice wedges in West-Siberian platform
PUB: Moscow, Moscow University Publishers, 1986
SCALE: 1:20 000 000

SOURCE: West-Siberian platform exodynamic (spatial-temporal regularities)/ edited by V.T. Trofimov, p.104
REGION: West Siberia
LEGEND Old and contemporary developing and static syngenetic and epigenetic wedge ice in different soils on different topography, soil wedges spread, degradation ice places
LAT/LONG: 50°00'-72°00'/60°00'-95°00'
INSTITUTE: Moscow State University
NUMBER 00004

AUTHOR Badu, I.U.B., Trofimov, V.T.
NAME: Extent of wedge ice and injective ice in Yamal Peninsula
PUB: Moscow, Moscow University Publishers, 1974
SCALE: 1:2 500 000
SOURCE: "Problems of cryolithology", issue 4 / edited by A.I. Popov, p.132
REGION: West Siberia, Yamal Peninsula
LEGEND Epigenetic and syngenetic wedge ice of active, nonactive and ice melting stages in ground and organic soils, volumetric macro ice content, polygonal-wedge relict two-floor ice, injective ice
LAT/LONG: 68°00'-74°00'/66°00'-74°00'
INSTITUTE: Moscow State University
NUMBER 00005

AUTHOR Badu, I.U.B.
NAME: General features of paleogeography of Gydan Peninsula to the end of Kazan period
PUB: Moscow, Moscow University Publishers, 1986
SCALE: 1:3 000 000
SOURCE: "Engineering-geological conditions of Gydan Peninsula", p.14
REGION: West Siberia, Gydan Peninsula
LEGEND Regions of formation of syngenetic and epigenetic wedge ice, permafrost of Salekhard strata, permafrost near surface with epigenetic wedge ice
LAT/LONG: 67°30'-74°00'/73°00' -84°00'
INSTITUTE: Moscow State University
NUMBER 00006

AUTHOR Badu, I.U.B.
NAME: General features of paleogeography of Gydan Peninsula to the end of Zyryan period
PUB: Moscow, Moscow University Publishers, 1986
SCALE: 1:3 000 000
SOURCE: "Engineering-geological conditions of Gydan Peninsula", p.17
REGION: West Siberia, Gydan Peninsula
LEGEND Divided plain (salekhrdskaya) with permafrost in upper part of section, coastal plain (kazanksevskaya) with permafrost in upper part of section, epigenetic and syngenetic wedge ice
LAT/LONG: 67°30'-74°00'/73°00' -84°00'
INSTITUTE: Moscow State University
NUMBER 00007

AUTHOR Badu, I.U.B.
NAME: General features of paleogeography of Gydan Peninsula to the end of Late-Middle Quaternary maximum (Yamal transgression)
PUB: Moscow, Moscow University Publishers, 1986
SCALE: 1:3 000 000
SOURCE: "Engineering-geological conditions of Gydan Peninsula", p.9
REGION: West Siberia, Gydan Peninsula
LEGEND Lowlands with sporadic permafrost, places of shelf with syngenetic wedge ice, freezing in the end of Middle Pleistocene, areas of formation of syngenetic and epigenetic wedge

ice
LAT/LONG: 67°30'-74°00'/73°00'-74°00'
INSTITUTE: Moscow State University
NUMBER 00008

AUTHOR NAME: Badu, I.U.B.
Map of ground ice genetic types and macro- ice content in 10m upper of deposit section in West-Siberian platform
PUB: Moscow, Moscow University Publishers, 1980
SCALE: 1:7 500 000
SOURCE: Trofimov, V.T., Badu, I.U.B., Dubikov, G.I. Cryogenic structure and ice content of permafrost in West-Siberian platform, p.84-85
REGION: West Siberia
LEGEND: Syngenetic and epigenetic polygonal wedge ice in different evolution states, volumetric macro ice content, injected ice, sheet ice, injected-segregated ice in different genetic complexes of deposits, buried ice

LAT/LONG: 64°00'-74°00'/65°00'-85°00'
INSTITUTE: Moscow State University
NUMBER 00009

AUTHOR NAME: Badu, I.U.B.
Map of the potential thaw settlement for the upper 10 m of permafrost section in West-Siberia platform
PUB: Moscow, Moscow University Publishers, 1980
SCALE: 1:7 500 000
SOURCE: Trofimov, V.T., Badu I.U.B., Dubikov, G.I. Cryogenic structure and ice content of permafrost in West-Siberian platform, p.216-217
REGION: West Siberia
LEGEND: Summary of potential settlement in the different genesis sediments, composition and ice content ground, catastrophic settlement in areas with injected ice

LAT/LONG: 64°00'-74°00'/65°00'-85°00'
INSTITUTE: Moscow State University
NUMBER 00010

AUTHOR NAME: Badu, I.U.B.
Primary features of paleogeography of Gydan Peninsula in Sartan epoch
PUB: Moscow, Moscow University Publishers, 1986
SCALE: 1:3 000 000
SOURCE: "Engineering-geological conditions of Gydan Peninsula", p.21
REGION: West Siberia, Gydan Peninsula
LEGEND: Permafrost extent on terrace plains, deep lakes and lake depression on thawing sheet ice places, polygonal-wedge ice erosion

LAT/LONG: 67°30'-74°00'/73°00'-84°00'
INSTITUTE: Moscow State University
NUMBER 00011

AUTHOR NAME: Badu, I.U.B., Trofimov, V.T.
Scheme of genetic types and ice content extent of upper part (10 m) permafrost in Yamal Peninsula
PUB: Moscow, Moscow University Publishers, 1974
SCALE: 1:2 500 000
SOURCE: "Problems of cryolithology", issue 4 / edited by A.I. Popov, p.126
REGION: West Siberia, Yamal Peninsula
LEGEND: Type of freezing, ice content, age of permafrost, boundaries of various age permafrost, Paleozoic bedrock

LAT/LONG: 68°00'-74°00'/66°00'-74°00'

INSTITUTE: Moscow State University
 NUMBER 00012

AUTHOR Badu, I.U.B., Kudryashov V.G., Lurie I.S., Trofimov, V.T., Firsov N.G.
 NAME: Scheme of permafrost average annual temperature in Yamal peninsula
 PUB: Moscow, Moscow University Publishers, 1977
 SCALE: 1:2 500 000
 SOURCE: Trofimov, V.T. "Regularities of spatial variability for engineering-geological conditions in West Siberian platform", p.67
 REGION: West Siberia
 LEGEND Prevailing temperature
 LAT/LONG: 73°00-67°00/66°00-74°00
 INSTITUTE: Moscow State University
 NUMBER 00013

AUTHOR Badu, I.U.B., Gruzlov, A.V., Gusev, A.B.
 NAME: Scheme of territories having different forecast erosional ground resistance
 PUB: Moscow, Moscow University Publishers, 1986
 SCALE: 1:5 000 000
 SOURCE: West-Siberian platform exodynamic (spatial-temporal regularities)/ edited by V.T. Trofimov, p.. 230-231
 REGION: West Siberia
 LEGEND Forecast resistance types depending on macro ice content and wedge ice in the different lithological composition places
 LAT/LONG: 56°00-72°00/60°00-85°00
 INSTITUTE: Moscow State University
 NUMBER 00014

AUTHOR Badu, I.U.B., Vasilchuk, I.U.K., Kashperiuk, P.I., Trofimov, V.T.
 NAME: Sketch map of massive ice extent in the Northern part of West-Siberian platform
 PUB: Moscow, Moscow University Publishers, 1986
 SCALE: 1:10 000 000
 SOURCE: West-Siberian platform exodynamic (spatial-temporal regularities) / edited by V.T. Trofimov, p.131
 REGION: West-North Siberia
 LEGEND Injected-segregated, injected, buried ice in different genesis and age sediments
 LAT/LONG: 65°00-85°00/68°00-72°00
 INSTITUTE: Moscow State University
 NUMBER 00015

AUTHOR Badu, I.U.B., Vasilchuk, I.U.K., Kashperiuk, P.I.
 NAME: Types of seasonal freezing referring to lithological composition and soil moisture content
 PUB: Moscow, Moscow University Publishers, 1986
 SCALE: 1:7 500 000
 SOURCE: West-Siberian platform exodynamic (spatial-temporal regularities)/ edited by V.T. Trofimov, P.. 76
 REGION: West Siberia
 LEGEND Main seasonal freezing and thawing types and their expansion boundaries
 LAT/LONG: 50°00-72°00/60°00-95°00
 INSTITUTE: Moscow State University
 NUMBER 00016

AUTHOR Badu, I.U.B., Vasilchuk, I.U.K., Kashperiuk, P.I.
 NAME: Types of seasonal freezing and thawing referring to ground average annual temperature and temperature amplitudes on surface ground massif
 PUB: Moscow, Moscow University Publishers, 1986

SCALE: 1:7 500 000
 SOURCE: West-Siberian platform exodynamic (spatial-temporal regularities) / edited by V.T. Trofimov, p.75
 REGION: West Siberia
 LEGEND: Types of seasonal freezing, boundaries of seasonal freezing and thawing types
 LAT/LONG: 50°00-72°00/60°00-95°00
 INSTITUTE: Moscow State University
 NUMBER: 00017

AUTHOR: Badu, I.U.D., Trofimov, V.T.
 NAME: Map of genetic types and ice content in upper permafrost (10 m) of section of West-Siberia platform
 PUB: Moscow, Moscow University Publishers, 1980
 SCALE: 1:7 500 000
 SOURCE: Trofimov, V.T., Badu, I.U.B., Dubikov G.I. Cryogenic structure and ice content of permafrost in West-Siberian platform, p.60-61
 REGION: West Siberia
 LEGEND: Genetic types, ice content
 LAT/LONG: 64°00-74°00/65°00-85°00
 INSTITUTE: Moscow State University
 NUMBER: 00018

AUTHOR: Baranov, I.J.
 NAME: Geocryology map of the USSR
 PUB: Moscow, GUGK, 1977
 SCALE: 1:5 000 000
 REGION: USSR
 LEGEND: Genetic type of permafrost, expansion, thickness, temperature, depth of seasonal freezing and thawing, frozen ground features and hillocky terrain conditions
 LAT/LONG: 38°00-82°00/30°00-170°00
 INSTITUTE: PNIIS, Gosstrois USSR, Glavpromstroyproect, Permafrost Institute, Academy of Science USSR
 NUMBER: 00019

AUTHOR: Baranov, I.J.
 NAME: Map of permafrost regions of the Earth
 PUB: Moscow, Moscow University Publishers, 1978
 SCALE: 1:50 000 000
 SOURCE: General permafrost (geocryology)/ edited by V.A. Kudriavtsev, p.16-17
 REGION: The Globe
 LEGEND: Permafrost regions and glacial caps, regular and irregular repetition freezing of soil-zones
 INSTITUTE: Obruchev Permafrost Institute
 NUMBER: 00020

AUTHOR: Baranov, L.J.
 NAME: Geocryological map of USSR
 PUB: Moscow, 1956
 SCALE: 1:10 000 000
 REGION: USSR
 LEGEND: Expansion, temperature, thickness, cryogenic processes and relief
 LAT/LONG: 38°00-80°00/20°00-170°00
 INSTITUTE: Obruchev Permafrost Institute
 NUMBER: 00021

AUTHOR: Baranova, I.U.P.
 NAME: General view and interpretation "lunar landscape"

PUB: Magadan, Knizh. izd-vo, 1972
SCALE: Large
SOURCE: Tomirdiario, S.V. "Perennial frost and industrialization of mountain countries and lowlands, the Magadan area and Yakutia", p.61
REGION: North-East
LEGEND: Alas depressions, thermodenudational terraces, baydzherakhs, thermokarst depression, places of original surface
LAT/LONG: 63°00-69°00/158°00-170°00
NUMBER: 00022

AUTHOR: Baulin, V.V.
NAME: Average annual temperature of the ground for districts with different conditions (3 maps)
PUB: Moscow, Nedra, 1985
SCALE: 1:30 000 000
SOURCE: Baulin, V.V. "Permafrost in the oil-gas containing regions of the USSR", p.24-25
REGION: West Siberia
LEGEND: Average annual temperature of ground in the places with the maximum and minimum snow cover, various lithological composition and thickness of the seasonal thaw layer
LAT/LONG: 55°00-74°00/60°00-86°00
INSTITUTE: PNIIS, Gosstro, USSR
NUMBER: 00023

AUTHOR: Baulin, V.V.
NAME: General geocryological areas of the Siberian platform
PUB: Moscow, Nedra, 1985
SCALE: 1:40 000 000
SOURCE: Baulin, V.V. "Permafrost in oil-gas containing regions of the USSR", p.92
REGION: North-East
LEGEND: Geocryological areas, zones, boundaries of zones, subzones and areas
LAT/LONG: 53°00-75°00/90°00-130°00
INSTITUTE: PNIIS, Gosstro, USSR
NUMBER: 00024

AUTHOR: Baulin, V.V., Chekhovskii A.L., Gruzlov, A.V.
NAME: Map of permafrost thickness in west Siberian plain
PUB: Moscow, Stroiizdat, 1976
SCALE: 1:5 000 000
SOURCE: Inset-map in monograph "Transactions of Industrial and Research Institute for Engineering Investigations of Construction", issue 49
REGION: West Siberia
LEGEND: Bedding near surface permafrost thickness, the depth of relict permafrost table, southern limit of relict permafrost, places of intensive contemporary freezing of ground, places of deeply bedding permafrost table
LAT/LONG: 60°00-71°00/60°00-87°00
INSTITUTE: PNIIS, Gosstro, USSR
NUMBER: 00025

AUTHOR: Baulin, V.V., Danilova, N.S., Kondratieva, K.A.
NAME: Map of permafrost expansion in Holocene climatic optimum
PUB: Moscow, Nauka, 1988
SCALE: 1:50 000 000
SOURCE: Geocryology of the USSR. European territory of the USSR / edited by E.D. Ershov, p.90
REGION: USSR
LEGEND: Permafrost expansion
LAT/LONG: 38°00-82°00/30°00-170°00
INSTITUTE: PNIIS, Gosstro USSR, Moscow State University

NUMBER 00026
AUTHOR Baulin, V.V., Chekhovskii, A.L.
NAME: Map of permafrost thickness in West Siberia plain
PUB: Novosibirsk, Nauka, 1990
SCALE: 1:3 000 000
SOURCE: Inset-map in monograph "Geocryological investigations history in West Siberia"/ edited by Nekrasov
REGION: West Siberia
LEGEND Thickness, depth of relict permafrost table, isolines of depth of permafrost base, areas of intense contemporary freezing, areas of deep permafrost table
LAT/LONG: 62°00'-74°00'/60°00'-90°00'
INSTITUTE: PNIIS, Gostroi USSR
NUMBER 00027

AUTHOR Baulin, V.V.
NAME: Map of taliks under lakes with different depth
PUB: Moscow, Nedra, 1985
SCALE: 1:25 000 000
SOURCE: Baulin, V.V. "Permafrost in the oil-gas containing regions of the USSR", p.49
REGION: West Siberia
LEGEND Southern limit of lake taliks (with snow cover and without snow cover, different depth of lakes)
LAT/LONG: 62°00'-74°00'/60°00'-86°00'
INSTITUTE: PNIIS, Gosstro, USSR
NUMBER 00028

AUTHOR Baulin, V.V.
NAME: Permafrost base in Urengoi gas field
PUB: Moscow, Nedra, 1985
SCALE: Large
SOURCE: Baulin, V.V. "Permafrost in oil-gas containing regions of the USSR", p.121
REGION: West Siberia
LEGEND Isolines (m) of permafrost base
LAT/LONG: 66°30'/77°00'
INSTITUTE: PNIIS, Gosstro, USSR
NUMBER 00029

AUTHOR Baulin, V.V., Danilova, N.S., Sukhodolskaia, L.A.
NAME: Permafrost map of the Holocene climatic optimum
PUB: Moscow, Nauka, 1981
SCALE: 1:40 000 000
SOURCE: "History of permafrost development in Eurasia", p.28. In the article by Baulin, V.V., Danilova, N.S., Sukhodolskaia, L.A. p.24-40
REGION: USSR
LEGEND Zone of deep layer permafrost; seasonal frost thickness,m; thermokarst, permafrost spread, average annual temperature geoisotherms
LAT/LONG: 30°00'-180°00'/48°00'-80°00'
INSTITUTE: PNIIS, Gosstro, USSR
NUMBER 00030

AUTHOR Baulin, V.V., Danilova, N.S., Sukhodolskaia, L.A.
NAME: Permafrost map of Late Pleistocene
PUB: Moscow, Nauka, 1981
SCALE: 1:40 000 000
SOURCE: "History of permafrost development in Eurasia", p.26. In the article by Baulin, V.V.,

REGION: Danilova, N.S., Sukhodolskaia, L.A. p.24-40
 LEGEND USSR
 Permafrost spread, depth of permafrost base, average annual temperature geoisotherms, southern limit of permafrost, frost cracking, cryogenic weathering, frost heaving
 LAT/LONG: 30°00-180°00/48°00-80°00
 INSTITUTE: PNIIS, Gosstro, USSR
 NUMBER 00031

AUTHOR Baulin, V.V.
 NAME: Permafrost zoning map of West Siberian plain (upper part of permafrost)
 PUB: GUGK, Moscow, 1985
 SCALE: 1:1 500 000
 REGION: West Siberia
 LEGEND Expansion, temperature, ice content, ground, peat, geomorphological levels
 LAT/LONG: 50°00-70°30/60°00-90°30
 INSTITUTE: Gosstro USSR, Geological Ministry, Moscow State University
 NUMBER 00032

AUTHOR Baulin, V.V., Chekhovskii, A.L.
 NAME: Permafrost zoning of West Siberian plain referring to thickness of permafrost and cryogenic structure
 PUB: Moscow, Gosstro of the USSR, 1985
 SCALE: 1:2 500 000
 REGION: West Siberia
 LEGEND Geomorphology, thickness, structure and condition of permafrost, depth of relict permafrost table
 LAT/LONG: 50°00-70°00/60°00-90°00
 INSTITUTE: PNIIS, Gosstro USSR
 NUMBER 00033

AUTHOR Baulin, V.V.
 NAME: Regionalization scheme of West Siberian plain (for the permafrost thickness map)
 PUB: Moscow, Nedra, 1985
 SCALE: 1:25 000 000
 SOURCE: Baulin, V.V. "Permafrost in the oil-gas containing regions of the USSR", p.66
 REGION: West Siberia
 LEGEND Permafrost regionalization, boundaries: permafrost zones, provinces and areas
 LAT/LONG: 55°00-74°00/60°00-86°00
 INSTITUTE: PNIIS, Gosstro, USSR
 NUMBER 00034

AUTHOR Baulin, V.V.
 NAME: Scheme of contemporary thermokarst extent in West Siberia platform
 PUB: Moscow, Moscow University Publishers, 1977
 SCALE: 1:10 000 000
 SOURCE: Trofimov, V.T. "Regularities of spatial variability for engineering-geological conditions in West Siberia platform", p.210
 REGION: West Siberia
 LEGEND Dynamic thermokarst zones, thermokarst on peatlands, boundaries of thermokarst on polygonal wedge ice, segregated and sheet ice
 LAT/LONG: 62°00-74°00/60°00-90°00
 INSTITUTE: PNIIS, Gosstro USSR
 NUMBER 00035

AUTHOR Baulin, V.V.
 NAME: Scheme of long standing ground heave phenomena extent range in West Siberian

- platform
 PUB: Moscow, Moscow University Publishers, 1977
 SCALE: 1:10 000 000
 SOURCE: Trofimov, V.T. "Regularities of spatial variability for engineering- geological conditions in West Siberian platform", p.206
 REGION: West Siberia
 LEGEND: Hillocky peatlands expansion range zones, areas of hydrolaccoliths and frost mounds
 LAT/LONG: 62°00-72°00/60°00-90°00
 INSTITUTE: PNIIS, Gosstroï USSR
 NUMBER: 00036
- AUTHOR: Baulin, V.V.
 NAME: Scheme of permafrost extent in Holocene Climatic Optimum . The third formation stage
 PUB: Moscow, Nedra, 1985
 SCALE: 1:25 000 000
 SOURCE: Baulin, V.V. "Permafrost in oil-gas containing regions of the USSR", p.147
 REGION: West Siberia
 LEGEND: Zones of continuous, discontinuous and deep seated permafrost, the depth of relict permafrost base and table, isotherms of average annual temperature, southern limit of permafrost
 LAT/LONG: 55°00-75°00/50°00-90°00
 INSTITUTE: PNIIS, Gosstroï, USSR
 NUMBER: 00037
- AUTHOR: Baulin, V.V.
 NAME: Scheme of permafrost extent in Demyanskoe Glacial epoch (Eopleistocene)
 PUB: Moscow, Nedra, 1985
 SCALE: 1:25 000 000
 SOURCE: Baulin, V.V. "Permafrost in oil-gas containing regions of the USSR", p.133
 REGION: West Siberia
 LEGEND: Subareal, subsea permafrost, permafrost under retaining basins, ice caps, southern limit of permafrost
 LAT/LONG: 55°00-75°00/50°00-90°00
 INSTITUTE: PNIIS, Gosstroï, USSR
 NUMBER: 00038
- AUTHOR: Baulin, V.V., Bykov, I. IU., Sadchikov, P.B.
 NAME: Scheme of permafrost expansion in the north-east of European part of the USSR
 PUB: Moscow, Stroizdat, 1984
 SCALE: 1:20 000 000
 SOURCE: Geocryological conditions and their change forecast in the primary development regions of the North, p.187
 REGION: North European part of the USSR and West Siberia
 LEGEND: Permafrost expansion, vertical structure, southern limit of relict permafrost
 LAT/LONG: 60°00-70°00/50°00-80°00
 INSTITUTE: PNIIS, Gosstroï USSR
 NUMBER: 00039
- AUTHOR: Baulin, V.V., Danilova, N.S., Pavlova, O.P.
 NAME: Scheme of permafrost extent for Medvezhie gas field region
 PUB: Moscow, Stroizdat, 1984
 SCALE: Large
 SOURCE: Geocryological conditions and their change forecast in primary development regions of the North, p.4
 REGION: West Siberia
 LEGEND: Expansion, depth of permafrost base, depth of taliks

LAT/LONG: 65°00-68°00/72°00-78°00
INSTITUTE: PNIIS, Gosstroj USSR
NUMBER 00040

AUTHOR Baulin, V.V.
NAME: Scheme of permafrost extent in Sartan Glaciation epoch (the second half of Neopleistocene). The second formation stage
PUB: Moscow, Nedra, 1985
SCALE: 1:25 000 000
SOURCE: Baulin, V.V. "Permafrost in oil-gas containing regions of the USSR", p.146
REGION: West Siberia
LEGEND Off shore permafrost, permafrost under ice caps, permafrost in areas of Kazan sea, average annual ground temperature
LAT/LONG: 55°00-75°00/50°00-90°00
INSTITUTE: PNIIS, Gosstroj, USSR
NUMBER 00041

AUTHOR Baulin, V.V.
NAME: Scheme of permafrost extent in Yamal transgressive epoch (the second part of Mesopleistocene). The first formation stage
PUB: Moscow, Nedra, 1985
SCALE: 1:25 000 000
SOURCE: Baulin, V.V. "Permafrost in oil-gas containing regions of the USSR", p.143
REGION: West Siberia
LEGEND Subsea permafrost, permafrost in regions with change glacial and sea conditions, permafrost under ice caps and subaeral, southern limit of permafrost (54-55° of n.l.)
LAT/LONG: 55°00-75°00/50°00-90°00
INSTITUTE: PNIIS, Gosstroj, USSR
NUMBER 00042

AUTHOR Baulin, V.V.
NAME: Scheme of permafrost extent in Tobolskoe Interglacial epoch (the first part of Mesopleistocene)
PUB: Moscow, Nedra, 1985
SCALE: 1:25 000 000
SOURCE: Baulin, V.V. "Permafrost in oil-gas containing regions of the USSR", p.142
REGION: West Siberia
LEGEND Subareal permafrost, places of subsea degradation permafrost, zone of subareal degradation permafrost, zone of permafrost, unfrozen ground
LAT/LONG: 55°00-75°00/50°00-90°00
INSTITUTE: PNIIS, Gosstroj, USSR
NUMBER 00043

AUTHOR Baulin, V.V., Trofimov, V.T.
NAME: Scheme of seasonal frozen ground and permafrost spreading
PUB: Moscow, Moscow University Publishers, 1977
SCALE: 1:10 000 000
SOURCE: Trofimov, V.T. "Regularities of spatial variability for engineering- geological conditions in West Siberian platform", p.64
REGION: West Siberia
LEGEND Seasonal frozen ground and permafrost expansion zones
LAT/LONG: 62°00-74°00/60°00-90°00
INSTITUTE: Moscow State University, PNIIS, Gosstroj USSR
NUMBER 00044

AUTHOR Baulin, V.V.

NAME: Sketch-map of permafrost spreading, thickness and structure in West Siberian plain
 PUB: Moscow, Nedra, 1985
 SCALE: 1:15 000 000
 SOURCE: Baulin, V.V. "Permafrost in the oil-gas containing regions of the USSR", p.69
 REGION: West Siberia
 LEGEND: Permafrost spreading, thickness, genetic complexes and lithological composition
 LAT/LONG: 60°00-73°00/60°00-86°00
 INSTITUTE: PNIIS, Gosstro, USSR
 NUMBER: 00045

AUTHOR: Baulin, V.V.
 NAME: Sketch-map of ice content in West-Siberia plain
 PUB: Moscow, Nedra, 1985
 SCALE: 1:15 000 000
 SOURCE: Baulin, V.V. "Permafrost in the oil-gas containing regions of the USSR", p.41
 REGION: West Siberia
 LEGEND: Ice content (3 ranges), boundaries: southern limit of before Holocene and Holocene syngenetical permafrost, epigenetic wedge ice, permafrost spreading, sheet ice
 LAT/LONG: 55°00-72°00/60°00-80°00
 INSTITUTE: PNIIS, Gosstro, USSR
 NUMBER: 00046

AUTHOR: Baulin, V.V.
 NAME: Sketch-map of relict permafrost
 PUB: Moscow, Nedra, 1985
 SCALE: 1:3 500 000
 SOURCE: Baulin, V.V. "Permafrost in the oil-gas containing regions of the USSR", p.73
 REGION: West Siberia
 LEGEND: Large blocks of relict permafrost, small islands of relict permafrost, geomorphological levels
 LAT/LONG: 58°00-63°00/69°00-78°00
 INSTITUTE: PNIIS, Gosstro, USSR
 NUMBER: 00047

AUTHOR: Baulin, V.V., Efimova I.V., Timofeev V.G.
 NAME: Sketch-map of relict permafrost
 PUB: Moscow, Moscow University Publishers, 1972
 SCALE: 1:2 000 000
 SOURCE: Permafrost Studies, XII, p.144
 REGION: West Siberia
 LEGEND: Relict permafrost in large massifs and islands, geomorphological level
 LAT/LONG: 59°00-63°00/69°00-78°00
 INSTITUTE: PNIIS, Gosstro USSR
 NUMBER: 00048

AUTHOR: Baulin, V.V., Belopukhova, E.B., Dubikov, G.I.
 NAME: West Siberian Geocryological Map
 PUB: Moscow, Academy of Science Publisher, 1968
 SCALE: 1:5 000 000
 SOURCE: Inset-map in paper by Baulin, V.V., Belopukhova, E.B., Dubikov, G.I. "Permafrost geographical features in West Siberia". Proceedings of Academy of Science USSR, Geography, p.64-70
 REGION: West Siberia
 LEGEND: Type of freezing, spreading, temperature, active layer depth, ice wedges and massive ice, recent and fossil thermokarst, perennial frost-heave mounds, southern permafrost limit
 LAT/LONG: 60°00-74°00/64°00-87°00

INSTITUTE: PNIIS, Gosstroj USSR
 NUMBER 00049

AUTHOR Baulin, V.V.
 NAME: West Siberian regionalization scheme for expansion and average annual temperature permafrost study
 PUB: Moscow, Nedra, 1985
 SCALE: 1:30 000 000
 SOURCE: Baulin, V.V. "Permafrost in oil-gas containing regions of the USSR", p.22
 REGION: West Siberia
 LEGEND Southern limit of permafrost, potential expansion permafrost zones
 LAT/LONG: 55°00-75°00/60°00-90°00
 INSTITUTE: PNIIS, Gosstroj USSR
 NUMBER 00050

AUTHOR Belopukhova, E.B., Tikhomirova, N.A., Sukhov A.G.
 NAME: Permafrost expansion in Yagenetta river head (Nadym-Pur interfluve)
 PUB: Moscow, Stroiizdat, 1984
 SCALE: Large
 SOURCE: Geocryological conditions and their change forecast in primary development regions of the North, p.167
 REGION: West Siberia
 LEGEND Expansion, geomorphological levels
 LAT/LONG: 64°00-68°00/72°00-78°00
 INSTITUTE: PNIIS, Gosstroj USSR
 NUMBER 00051

AUTHOR Belopukhova, E.B., Tikhomirova, N.A., Sukhov, A.G.
 NAME: Permafrost extent in Yamsovey down-stream (left shore Pur valley)
 PUB: Moscow, Stroiizdat, 1984
 SCALE: Large
 SOURCE: Geocryological conditions and their change forecast in primary development regions of the North, p.166
 REGION: West Siberia
 LEGEND Expansion, geomorphological levels
 LAT/LONG: 64°00-68°00/72°00-78°00
 INSTITUTE: PNIIS, Gosstroj USSR
 NUMBER 00052

AUTHOR Belopukhova, E.B.
 NAME: Permafrost extent map of central part of West Siberia
 PUB: Moscow, TSINIS, 1972
 SCALE: 1:5 000 000
 SOURCE: "Geocryological research for engineering investigations for construction"(Transactions of PNIIS, vol. XVIII), p.95
 REGION: West Siberia
 LEGEND Permafrost spreading
 LAT/LONG: 60°00-68°00/60°00-87°00
 INSTITUTE: PNIIS, Gosstroj, USSR
 NUMBER 00053

AUTHOR Belopukhova, E.B.
 NAME: Scheme of permafrost temperature zoning of Yamburg tectonic structure
 PUB: Moscow, Stroiizdat, 1984
 SCALE: Large
 SOURCE: Geocryological conditions and their change forecast in primary development of the North,

p.87

REGION: West Siberia
 LEGEND 4 types of places with different ground temperature ranges
 LAT/LONG: 68°00-69°00/74°00-77°00
 INSTITUTE: PNIIS, Gosstroj USSR
 NUMBER 00054

AUTHOR Belopukhova, E.B.
 NAME: Scheme of polygonal ground extent range in West Siberian platform
 PUB: Moscow, Moscow University Publishers, 1977
 SCALE: 1:10 000 000
 SOURCE: Trofimov, V.T. "Regularities of spatial variability of engineering and geological conditions of the West Siberian platform

REGION: West Siberia
 LEGEND Polygonal ground growth and dormant stages of hillocky terrain on poor and well drained places, young hillocky terrain relief
 LAT/LONG: 62°00-74°00/60°00-90°00
 INSTITUTE: PNIIS, Gosstroj USSR
 NUMBER 00055

AUTHOR Belopukhova, E.B.
 NAME: Sketch-map of polygonal relief
 PUB: Yakutsk, Yakutskoe knizh. izd., 1966
 SCALE: 1:20 000 000
 SOURCE: "Scientific report of VIII All-Union Conference on geocryology", no.6. Geomorphological section, p.124. In the report by Belopukhova, E.B. p.117-125

REGION: West Siberia
 LEGEND Area polygonal relief in stage of growth and dormant, hillocky terrain on bad and good drained places, on peatland and on soil, southern limit of continuous permafrost at present time
 LAT/LONG: 60°00-73°00/63°00-87°00
 INSTITUTE: PNIIS, Gosstroj, USSR
 NUMBER 00056

AUTHOR Belopukhova, E.B., Dubikov, G.I.
 NAME: West Siberia regionalization sketch-map referring to permafrost ice content
 PUB: Moscow, TSINIS, 1972
 SCALE: 1:10 000 000
 SOURCE: "Geocryological research for engineering investigations for construction"(Transactions of PNIIS, vol.XVIII), p.33

REGION: West Siberia
 LEGEND Boundaries of zones, subzones, districts
 LAT/LONG: 60°00-74°00/60°00-90°00
 INSTITUTE: PNIIS, Gosstroj, USSR
 NUMBER 00057

AUTHOR Bobov, N.G., Novoselskaiia, N.B.
 NAME: Kamchatka geocryological scheme
 PUB: Novosibirsk, Nauka, 1975
 SCALE: 1:10 000 000
 SOURCE: "Regional and special geocryological investigations", p.37

REGION: Kamchatka
 LEGEND Expansion (real and estimated), temperature, thermokarst lakes and depressions, frost mounds
 LAT/LONG: 53°00-61°00/155°00-165°00
 NUMBER 00058

AUTHOR Bobov, N.G., Molodykh, I.I.
NAME: Zoning of European part of the USSR referring to cryogenic processes and expansion
PUB: Moscow, Nauka, 1988
SCALE: 1:50 000 000
SOURCE: Geocryology of the USSR. European territory of the USSR / edited by E.D. Ershov, p.252
REGION: European part of the USSR
LEGEND Contemporary and late Valdai epoch permafrost expansion, processes
LAT/LONG: 40°00-70°00/20°00-70°00
INSTITUTE: VSEGINGEO
NUMBER 00059

AUTHOR Boyarskii, O.G., Mitt, K.L.
NAME: Regionalization sketch-map of Anabaro-Olenek North referring to thermokarst topography development
PUB: Moscow, Moscow University Publishers, 1964
SCALE: 1:2 500 000
SOURCE: Permafrost Studies, IV, p.152
REGION: North-East of the USSR
LEGEND Thermokarst development regions in different age and genesis deposits, polygonal ground expansion, classes, remanent lakes
LAT/LONG: 71°00-73°00/120°00-124°00
NUMBER 00060

AUTHOR Boyarskii, O.G., Maksimova L.N., Romanovskii N.N
NAME: Scheme of Patomski upland permafrost temperature
PUB: Moscow, Moscow University Publishers, 1968
SCALE: 1:2 500 000
SOURCE: Permafrost Studies, VIII, p.207
REGION: Baikal
LEGEND Number of area and region (the table lists the permafrost expansion, ground temperature, snow cover thickness, ground content and age, geomorphological levels)
LAT/LONG: 57°00-60°00/112°00-119°00
INSTITUTE: Moscow State University
NUMBER 00061

AUTHOR Boyarskii, O.G., Mitt, K.L.
NAME: Sketch-map of mounds of different genesis extent in Anabaro-Olenekskoi lowland
PUB: Moscow, Nauka, 1974
SCALE: 1:2 500 000
SOURCE: Fotiev, S.M., Danilova, N.S., Sheveleva, N.S. Permafrost conditions of Central Siberia, p. 46
REGION: Central Siberia
LEGEND Pingos, thermokarst topography, remanents of original rocks
LAT/LONG: 70°00-76°00/114°00-120°00
NUMBER 00062

AUTHOR Bubnov, V.M., Pokrovskii, N.S.
NAME: Permafrost thickness and tectonic structure in the Nizh. Tunguska drainage basin
PUB: Moscow, Nedra, 1985
SCALE: 1:9 000 000
SOURCE: Baulin, V.V. "Permafrost in oil-gas containing regions of the USSR", p.122
REGION: Central Siberia
LEGEND Permafrost thickness, zone of geothermal anomaly near deep-seated domes of tectonic structures
LAT/LONG: 64°00-77°00/95°00-110°00
NUMBER 00063

AUTHOR NAME: Buldovich, S.N., Melentiev, V.S., Naumov, M.S.
 Scheme of permafrost-hydrogeological conditions and fractured tectonics in Neryungi place
 PUB: Moscow, Moscow University Publishers, 1976
 SCALE: Large
 SOURCE: Permafrost Studies, XV, p.122
 REGION: Central Siberia
 LEGEND: Permafrost expansion, wells with underground water level, alluvial deposits
 LAT/LONG: 57°00-00°00/125°00-00°00
 INSTITUTE: Moscow State University
 NUMBER: 00064

AUTHOR NAME: Chekhovskii, A.L., Shamanova I.I.
 Map of lake depths (lakes, below which are possible taliks)
 PUB: Moscow, Stroiizdat, 1974
 SCALE: 1:10 000 000
 SOURCE: "Transactions of Industrial and Research Institute for Engineering Investigations of Construction", no.49, p.74
 REGION: West Siberia
 LEGEND: Depths of lake (7 ranges)
 LAT/LONG: 60°00-74°00/60°00-90°00
 INSTITUTE: PNIIS, Gosstro, U.S.S.R.
 NUMBER: 00065

AUTHOR NAME: Chekhovskii, A.L.
 Regionalization scheme of Kara Sea referring to benthic water layer temperature
 PUB: Moscow, TSINIS, 1972
 SCALE: 1:12 000 000
 SOURCE: "Geocryological research for engineering investigations for construction" (Transactions of PNIIS, vol. XVIII), p.105
 REGION: Kara Sea
 LEGEND: Benthic water layer average annual temperature (5 ranges), oceanic, suboceanic and subcontinental areas
 LAT/LONG: 70°00-90°00/54°00-114°00
 INSTITUTE: PNIIS, Gosstro, USSR
 NUMBER: 00066

AUTHOR NAME: Cherniadev, B.P.
 The changes of the position of southern limit of permafrost in West Siberia under disturbed natural conditions
 PUB: Moscow, TSINIS, 1971
 SCALE: 1:15 000 000
 SOURCE: "Geocryological and hydrogeological research for engineering investigations", Transactions of PNIIS, vol. 8, p.192
 REGION: West Siberia
 LEGEND: Southern limit: possible neogenesis of permafrost under cooling to maximum temperature; thawing potential of permafrost by moving of moss-lichen cover
 LAT/LONG: 52°00-72°00/60°00-90°00
 INSTITUTE: PNIIS, Gosstro, USSR
 NUMBER: 00067

AUTHOR NAME: Cherniadev, V.P.
 Map-scheme of upper limiting conditions and natural zones of West Siberia
 PUB: Moscow, Stroiizdat, 1987
 SCALE: 1:10 000 000
 SOURCE: "Recommendations to estimate change of permafrost conditions in industrialization

- territories of West Siberia/ PNIIS, Gosstroj USSR", p.9
- REGION:** West Siberia
- LEGEND:** Summary of mean monthly air temperature in warm and cold periods, summary mean monthly temperature of exposed soil surface, average snow cover, duration of summer period, natural zones
- LAT/LONG:** 60°00-72°00/60°00-85°00
- INSTITUTE:** PNIIS, Gosstroj, USSR
- NUMBER:** 00068
- AUTHOR:** Cherniadev, V.P.
- NAME:** Sketch-map of limiting conditions of perennial freezing
- PUB:** Moscow, TSINIS, 1971
- SCALE:** 1:20 000 000
- SOURCE:** "Geocryological and hydrogeological research for engineering investigations", Transactions of PNIIS, vol. 8, p.191
- REGION:** West Siberia
- LEGEND:** Summary of degree months in maximum cold period, summary of degree months in minimum and maximum warming-up periods
- LAT/LONG:** 52°00-72°00/60°00-90°00
- INSTITUTE:** PNIIS, Gosstroj, USSR
- NUMBER:** 00069
- AUTHOR:** Chizhova, N.I.
- NAME:** Map of Aldan-Timpton interfluvial relative formation of an icing(%)
- PUB:** Moscow, Nedra, 1989
- SCALE:** 1:2 000 000
- SOURCE:** Geocryology of the USSR. Central Siberia / edited by E.D. Ershov, p.309
- REGION:** Central Siberia
- LEGEND:** Relative formation of an icing (6 ranges), icings on the ground surface
- LAT/LONG:** 55°00-60°00/122°00-130°00
- INSTITUTE:** Moscow State University
- NUMBER:** 00070
- AUTHOR:** Chizhova, N.I.
- NAME:** Map of icings of Aldano-Timptonski interfluvial
- PUB:** Moscow, Moscow University Publishers, 1980
- SCALE:** 1:2 500 000
- SOURCE:** Ospennikov, E.N., Trush, N.I., Chizhov, A.B., Chizhova, N.I. Exogenetic geological processes and phenomena (South Yakutia), p.188
- REGION:** South Yakutia
- LEGEND:** Icing coefficient, icing on the ground surface
- LAT/LONG:** 57°00-69°00/123°00-128°00
- INSTITUTE:** Moscow State University
- NUMBER:** 00071
- AUTHOR:** Churinov, M.V., Tsipina, I.M., Lazareva, V.P.
- NAME:** Stratigraphic and genetic complexes and engineering- geological groups of sedimentary rocks
- PUB:** Moscow, GUGK, 1983
- SCALE:** 1:7 500 000
- SOURCE:** Atlas of hydrogeological and engineering-geological maps of the USSR
- REGION:** USSR
- LEGEND:** Expansion frozen and unfrozen Quaternary ground
- LAT/LONG:** 38°00-80°00/20°00-170°00
- INSTITUTE:** VSEGINGEO
- NUMBER:** 00072

AUTHOR: Danilova, N.S., Kondratieva, K.A.
 NAME: Central Siberia permafrost regionalization referring to cryogenic processes and frozen ground features development
 PUB: Moscow, Nedra, 1989
 SCALE: 1:20 000 000
 SOURCE: Geocryology of the USSR. Central Siberia/ edited by E.D. Ershov, p.125
 REGION: Central Siberia
 LEGEND: Latitudinal thermal zones, thermal and regional subzones, regions of prevalent Quaternary deposits
 LAT/LONG: 50°00-75°00/80°00-135°00
 INSTITUTE: Moscow State University
 NUMBER: 00073

AUTHOR: Danilova, N.S.
 NAME: Sketch-map of the structure of perennial frozen ground in Central Siberia
 PUB: Moscow, TSINIS, 1972
 SCALE: 1:10 000 000
 SOURCE: "Geocryological research for engineering investigations for construction" (Transactions of PNIIS vol. XVIII), p.166
 REGION: Central Siberia
 LEGEND: Types of freezing, permafrost thickness, ice content and cryogenic structure, genesis and composition of deposits, time of freezing, wedge ice (old and contemporary), cave and sheet ice
 LAT/LONG: 52°00-80°00/84°00-140°00
 INSTITUTE: PNIIS, Gosstroï, USSR
 NUMBER: 00074

AUTHOR: Danilova, N.S.
 NAME: Sketch-map of polygonal ice wedges of Central Siberia
 PUB: Moscow, Nedra, 1975
 SCALE: 1:20 000 000
 SOURCE: Fotiev, S.M., Danilova, N.S., Sheveleva, N.S. Permafrost conditions of Central Siberia, p.128
 REGION: Central Siberia
 LEGEND: Zone of occurrence of relict and contemporary permafrost
 LAT/LONG: 52°00-80°00/78°00-138°00
 INSTITUTE: PNIIS, Gosstroï USSR
 NUMBER: 00075

AUTHOR: Demidiuk, L.M.
 NAME: Geocryological scheme of Charanorskoi lowland
 PUB: Moscow, Nedra, 1989
 SCALE: Large
 SOURCE: Geocryology of the USSR. Mountain countries of the southern USSR / edited by E.D. Ershov, p.118
 REGION: Transbaikal
 LEGEND: Expansion, temperature, thickness, cryogenic processes and frozen ground features
 LAT/LONG: 50°00-52°00/114°00-119°00
 INSTITUTE: Moscow State University
 NUMBER: 00076

AUTHOR: Demidiuk, L.M., Shaumian, L.V.
 NAME: Sketch-map of jointing and permafrost extent of Talnakh field
 PUB: Moscow, Moscow University Publishers, 1969
 SCALE: Large
 SOURCE: Permafrost Studies, IX, P.35

REGION: Central Siberia
LEGEND Expansion, temperature, permafrost thickness, content and age deposits, jointing zones
LAT/LONG: 69°00-00°00/88°00-89°00
NUMBER 00077

AUTHOR Dubikov, G.I.
NAME: Districts with ice wedges and ground wedges in alluvial terraces
PUB: Moscow, Academy of Sciences, 1962
SCALE: 1:5 000 000
SOURCE: "Academy of Sciences Information, Geographical Serie"s, no. 6, p.81
REGION: West Siberia
LEGEND Wedge ice and soil wedges
LAT/LONG: 65°30-69°00/68°00-74°00
INSTITUTE: PNIIS, Gosstroj USSR
NUMBER 00078

AUTHOR Dubikov, G.I.
NAME: Geological sketch-map with massive ice outcropping
PUB: Moscow, Stroiizdat, 1983
SCALE: Large
SOURCE: Dubikov, G.I. Problems of Regional Engineering Geocryology, p.53
REGION: West Siberia, Yamal Peninsula
LEGEND Sites of massive ice outcrops; bore-holes with massive ice; deposits: genesis and age
LAT/LONG: 70°00/70°00
INSTITUTE: PNIIS, Gosstroj USSR
NUMBER 00079

AUTHOR Dubikov, G.I., Ivanova, N.V.
NAME: Map of saline permafrost in West Siberia
PUB: Moscow, Scientific Council of the Earth Cryology, 1990
SCALE: 1:12 000 000
SOURCE: "Saline permafrost as a foundation for construction. Collected scientific articles"/ Edited by Vyalov S.S., p.6
REGION: West Siberia
LEGEND Saline ground, iono-saline content, general genetic complexes, permafrost extent with different saline content, districts with bedding table of saline Paleogene ground to 50 m depth
LAT/LONG: 65°00-73°00/60°00-90°00
INSTITUTE: PNIIS, Gosstroj, USSR
NUMBER 00080

AUTHOR Dubikov, G.I., Ivanov, N.V.
NAME: Scheme of marine saline and non-saline permafrost in West Siberia
PUB: Moscow, Nauka, 1986
SCALE: 1:7 500 000
SOURCE: "Permafrost formation and forecast of cryogenic processes forecast", p.18. In the article by Dubikov, G.I. p.14-27
REGION: West Siberia
LEGEND Boundaries: Holocene freezing in upper layer permafrost, maximum advance of the sea in Pleistocene (by Lazukov), saline and non-saline ground, quantity of test
LAT/LONG: 46°00-73°00/64°00-84°00
INSTITUTE: PNIIS, Gosstroj, USSR
NUMBER 00081

AUTHOR Dubikov, G.I., Ivanova, N.V.
NAME: Scheme of saline permafrost extent in the USSR

PUB: Moscow, Scientific Council of the Earth Cryology, 1990
 SCALE: 1:35 000 000
 SOURCE: "Saline permafrost as a foundation for constructions. Collected scientific articles" / Edited by S.S. Vyalov, p.4
 REGION: The USSR
 LEGEND: Type of saltings, saline permafrost table, saline and non-saline permafrost spreading, boundary of saline permafrost, southern limit of permafrost
 LAT/LONG: 40°00-80°00/40°00-170°00
 INSTITUTE: PNIIS, Gosstroi, USSR
 NUMBER: 00082

AUTHOR: Dubikov, G.I., Belopukhova, E.B., Stremiakov, A.I.A., Sukhov, A.G.
 NAME: Sketch-map of Byngapur tectonic structure permafrost regionalization
 PUB: Moscow, Stroiizdat, 1984
 SCALE: Large
 SOURCE: Geocryological conditions and their changes forecast in the primary development of the North, p.122
 REGION: West Siberia
 LEGEND: Zoning referring to permafrost expansion, lithological composition, genesis and age of deposits
 LAT/LONG: 64°00-68°00/72°00-78°00
 INSTITUTE: PNIIS, Gosstroi USSR
 NUMBER: 00083

AUTHOR: Dubikov, G.I., Shmelev, L.M.
 NAME: Sketch-map of southern limit of mineral and organic perennial ground between the Urals and the Ob
 PUB: Moscow, Stroiizdat, 1976
 SCALE: 1:5 000 000
 SOURCE: "Transactions of Industrial and Research Institute for Engineering Investigations of Construction", issue 49, p.87
 REGION: West Siberia
 LEGEND: Boundaries by Kunitsin L.F., Belopukhova, E.B., Popov, A.I., Shpolianskaia, N.A., authors
 LAT/LONG: 60°00-64°00/60°00-66°00
 INSTITUTE: PNIIS, Gosstroi, USSR
 NUMBER: 00084

AUTHOR: Dubikov, G.I.
 NAME: Sketch-map of lithological composition and permafrost ice content to the depth 10 m in Yamburg structural high area
 PUB: Moscow, Stroiizdat, 1984
 SCALE: Large
 SOURCE: Geocryological conditions and their change forecast in the primary development regions of the North, p.108
 REGION: West Siberia
 LEGEND: Types of freezing, lithological composition, genetic and ground ice content
 LAT/LONG: 68°00-69°00/74°00-77°00
 INSTITUTE: PNIIS, Gosstroi USSR
 NUMBER: 00085

AUTHOR: Dunaeva, E.N., Koreisha, M.M.
 NAME: Geocryological sketch-map of the Caucasus
 PUB: Moscow, Nedra, 1989
 SCALE: 1:10 000 000
 SOURCE: Geocryology of the USSR. Mountain countries of the southern USSR / edited by E.E. Ershov, p.335

REGION: Caucasus
 LEGEND Expansion, thickness, temperature, types of seasonal freezing of soils
 LAT/LONG: 38°00-45°00/40°00-50°00
 INSTITUTE: Moscow State University, PNIIS Gosstroj USSR
 NUMBER 00086

AUTHOR Ershov, E.D., Danilov, I.D.
 NAME: Map of permafrost expansion types and large concentration of surface and ground ice
 PUB: Moscow, Nauka, 1988
 SCALE: 1:50 000 000
 SOURCE: Geocryology of the USSR. European territory of the USSR / edited by E.D. Ershov, p.111
 REGION: USSR
 LEGEND Cryogenic ground types, injected ice, sheet ice, polyzonal wedge ice, buried ice, glaciers, permafrost expansion
 LAT/LONG: 38°00-82°00/30°00-170°00
 INSTITUTE: Moscow State University
 NUMBER 00087

AUTHOR Ershov, E.D., Dunaeva, E.N., Parmuzin, S.IU.
 NAME: Sketch-map of seasonal freezing and thawing of soil types in the USSR
 PUB: Moscow, Nauka, 1988
 SCALE: 1:50 000 000
 SOURCE: Geocryology of the USSR. European territory of the USSR / edited by E.D. Ershov, p.141
 REGION: USSR
 LEGEND Thawing of soil types referring to average annual ground temperature and temperature fluctuations amplitude on surface
 LAT/LONG: 35°00-82°00/30°00-170°00
 INSTITUTE: Moscow State University
 NUMBER 00088

AUTHOR Ershov, E.D.
 NAME: Sketch-map of contemporary permafrost expansion on the Earth
 PUB: Moscow, Nauka, 1988
 SCALE: 1:100 000 000
 SOURCE: Geocryology of the USSR. European territory of the USSR / edited by E.D. Ershov, p.15
 REGION: The Globe
 LEGEND Seasonal frozen ground and permafrost expansion
 INSTITUTE: Moscow State University
 NUMBER 00089

AUTHOR Evseev, V.P.
 NAME: Scheme of palsas and flat-topped polygonal peatlands contemporary spreading
 PUB: Moscow, Moscow University Publishers, 1976
 SCALE: 1:10 000 000
 SOURCE: "Cryolithology problems", issue V / edited by A.I. Popov, p.108
 REGION: European North and West Siberia
 LEGEND Southern limit of palsas (southern limit of permafrost), northern limit of palsas, northern and southern limit of flat-topped polygonal peatlands
 LAT/LONG: 55°00-72°00/30°00-90°00
 INSTITUTE: Moscow State University
 NUMBER 00090

AUTHOR Fedorov, A.N.
 NAME: Sketch-map of physiographical districts in Leno-Amginsk interfluvial northern part
 PUB: Yakutsk, Permafrost Institute Publishers, 1983
 SCALE: 1:20 000 000

SOURCE: Geographical research in Yakutia, p.130
REGION: Southern Yakutia
LEGEND: Number of district (permafrost terrain, thickness, temperature, ice content)
LAT/LONG: 61°00-63°00/128°00-135°00
INSTITUTE: Permafrost Institute
NUMBER: 00091

AUTHOR: Fedorovich, D.I., Zhukov, V., Vialov, S.S.
NAME: Geocryological sketch-map
PUB: Moscow, Stroiizdat, 1980
SCALE: 1:25 000 000
SOURCE: "Textbook for engineering of bases and foundations on permafrost/Gersevanov Institute of Foundation and Underground Construction, Gosstroi, USSR, p.8-9
REGION: USSR
LEGEND: Permafrost spread, thickness, temperature of the ground to the depth 10m, southern limit of permafrost
LAT/LONG: 38°00-85°00/20°00-170°00
INSTITUTE: NIIOSP
NUMBER: 00092

AUTHOR: Feldman, G.M.
NAME: Atlas of forecast permafrost map of North-West Siberia
PUB: Yakutsk, Permafrost Institute Publishers, 1983
SCALE: 1:20 000 000
SOURCE: Feldman, G.M. Methodology book of permafrost temperature dynamic forecast (e.g., north West Siberia), p.7-40, 64 maps
REGION: West Siberia
LEGEND: Isotherms of average annual frozen ground temperature on active layer base and isolines of active layer base thickness (for 63 combinations)
LAT/LONG: 60°00-74°00/60°00-84°00
INSTITUTE: Permafrost Institute
NUMBER: 00093

AUTHOR: Fotiev, S.M.
NAME: Map of permafrost dynamics in Pleistocene and Holocene
PUB: Moscow, Moscow University Publishers, 1978
SCALE: 1:35 000 000
SOURCE: General permafrost(geocryology)/ edited by V.A. Kudriavtsev, p.386
REGION: USSR
LEGEND: Relict permafrost zone under Polar basin, partial and complete thawing of Pleistocene permafrost, Holocene frozen and thawing ground, thickness of late Holocene frozen and thawing ground
LAT/LONG: 38°00-82°00/30°00-170°00
INSTITUTE: PNIIS, Gosstroi USSR
NUMBER: 00094

AUTHOR: Fotiev, S.M.
NAME: Regionalization scheme of West Siberia referring to discontinuous and permafrost spread
PUB: Yakutsk, Geocryological Institute Publishers, 1986
SCALE: 1:10 000 000
SOURCE: "Questions of geocryological mapping", p.44. In the article by Fotiev, S.M. p.38-52
REGION: West Siberia
LEGEND: Permafrost spread, unfrozen ground, taliks, areas of types of cryolithogenesis
LAT/LONG: 59°30-74°00/69°00-84°00
INSTITUTE: PNIIS, Gosstroi, USSR
NUMBER: 00095

AUTHOR NAME: Fotiev, S.M.
 Regionalization scheme of West Siberia region referring to conditions of open-water absorption for talik formation
 PUB: Moscow, Nauka, 1991
 SCALE: 1:12 000 000
 SOURCE: "Permafrost and cryogenic processes: Collection of scientific articles" / Edited by G.I. Dubikov, p.74
 REGION: West Siberia
 LEGEND: Calculated boundary of the region where possible of open water absorption taliks were formed by favourable geological-geomorphological conditions, natural zones boundary
 LAT/LONG: 58°00-74°00/66°00-84°00
 INSTITUTE: PNIIS, Gosstroj, USSR
 NUMBER: 00096

AUTHOR NAME: Fotiev, S.M.
 Scheme of different types of permafrost extent in West Siberia
 PUB: Moscow, TSINIS, 1972
 SCALE: 1:12 000 000
 SOURCE: "Geocryological research for engineering investigations for construction" (Transactions of PNIIS, vol.XVIII), p.118
 REGION: West Siberia
 LEGEND: Permafrost stages, permafrost types and subtypes
 LAT/LONG: 60°00-72°00/60°00-86°00
 INSTITUTE: PNIIS, Gosstroj, U.S.S.R.
 NUMBER: 00097

AUTHOR NAME: Fotiev, S.M.
 Scheme of West Siberia regionalization referring to conditions of open talik formation
 PUB: Moscow, Nauka, 1991
 SCALE: 1:12 000 000
 SOURCE: "Permafrost and cryogenic processes: Collection of scientific articles" / Edited by G.I. Dubikov, p.77
 REGION: West Siberia
 LEGEND: Condition: very unfavourable - talik area less than 5%, unfavourable- talik area 5-25%, favourable - talik area (insolation-insolation-heat, water absorption and lake taliks)--25-50% and 50-75%, very favourable - 75-100% talik area
 LAT/LONG: 58°00-74°00/66°00-84°00
 INSTITUTE: PNIIS, Gosstroj, U.S.S.R.
 NUMBER: 00098

AUTHOR NAME: Fotiev, S.M., Danilova, N.S., Sheveleva, N.S.
 Sketch geocryological map of Central Siberia
 PUB: Moscow, Nauka, 1974
 SCALE: 1:7 500 000
 SOURCE: Fotiev, S.M., Danilova, N.S., Sheveleva, N.S. Permafrost conditions of Central Siberia, p.135
 REGION: Central Siberia
 LEGEND: Thickness, expansion, ground temperature, cryogenic processes, composition and genesis Quaternary deposits
 LAT/LONG: 52°00-80°00/78°00-138°00
 INSTITUTE: PNIIS, Gosstroj USSR
 NUMBER: 00099

AUTHOR NAME: Fotiev, S.M., Danilova, N.S., Sheveleva, N.S.
 Sketch-map of Quaternary deposits with their cryogenic structure and ice content characteristics (Northern Siberian lowland)

PUB: Moscow, Nauka, 1974
 SCALE: 1:7 500 000
 SOURCE: Fotiev, S.M., Danilova, N.S., Sheveleva, N.S. Permafrost conditions of Central Siberia, p.43
 REGION: Northern Siberian lowland
 LEGEND: Type of freezing, cryogenic structure, sheet ice, ice content, possible thaw settlements, age and genesis type of deposits
 LAT/LONG: 67°00-82°00/80°00-130°00
 INSTITUTE: PNIIS, Gosstroj USSR
 NUMBER: 00100

AUTHOR: Fotiev, S.M.
 NAME: Sketch-map of permafrost extent in Central Siberia
 PUB: Moscow, Nauka, 1975
 SCALE: 1:20 000 000
 SOURCE: Fotiev, S.M., Danilova, S.M., Sheveleva, N.S. Permafrost conditions of Central Siberia, p.105
 REGION: Central Siberia
 LEGEND: Expansion
 LAT/LONG: 52°00-80°00/78°00-138°00
 INSTITUTE: PNIIS, Gosstroj USSR
 NUMBER: 00101

AUTHOR: Fotiev, S.M.
 NAME: Sketch-map of permafrost thickness and spreading
 PUB: Moscow, Nedra, 1975
 SCALE: 1:20 000 000
 SOURCE: Fotiev, S.M., Danilova, N.S., Sheveleva, N.S. Permafrost conditions of Central Siberia, p.113
 REGION: Central Siberia
 LEGEND: Cryogenic structure (one and two stage), factual data about permafrost thickness
 LAT/LONG: 52°00-80°00/78°00-138°00
 INSTITUTE: PNIIS, Gosstroj USSR
 NUMBER: 00102

AUTHOR: Garagulia, L.S., Ershov, E.D., Kondratieva, K.A.
 NAME: Engineering-geological zoning map of the USSR
 PUB: Moscow, Nedra, 1988
 SCALE: 1:20 000 000
 SOURCE: Inset-map in monograph "Geocryology of the USSR. European territory of the USSR / edited by E.D. Ershov, E.D. "
 REGION: USSR
 LEGEND: Areas with technical changes of geocryological conditions and with activity and origin of cryogenic processes and frozen ground features, types of geocryological conditions technical changes, permafrost zones (expansion and temperature)
 LAT/LONG: 38°00-82°00/30°00-170°00
 INSTITUTE: Moscow State University
 NUMBER: 00103

AUTHOR: Garagulia, L.S., Trush, N.I., Bogoliubov, A.N.
 NAME: Permafrost map of Eruda valley
 PUB: Moscow, Nauka, 1989
 SCALE: Large
 SOURCE: Geocryology of the USSR. Mountain countries of the southern USSR / edited by E.D. Ershov, p.291
 REGION: Southern Siberia

LEGEND Temperature, lithological composition, total moisture content to the depth of 1,0 m
LAT/LONG: 56°00-62°00/89°00-96°00
INSTITUTE: Moscow State University
NUMBER 00104

AUTHOR Gasanov, Sh.Sh.
NAME: Interaction of atmosphere/cryosphere with the earth surface
PUB: Novosibirsk, Nauka, 1975
SCALE: 1:400 000 000
SOURCE: "Regional and special geocryological investigations", p.101
REGION: The Globe
LEGEND Area of atmosphere/cryosphere interaction
NUMBER 00105

AUTHOR Gasanov, Sh.Sh.
NAME: Map-scheme of general regularities of wedge and injective ice distributions
PUB: Moscow, Nauka, 1981
SCALE: 1:50 000 000
SOURCE: Gasanov, Sh.Sh. "Cryolithological analysis", p.113
REGION: The USSR
LEGEND Ground ice, wedge and injective ice, sea boundary in boreal transgression (by Lavrova, Troitskii)
LAT/LONG: 50°00-80°00/30°00-170°00
NUMBER 00106

AUTHOR Gavrilov, A.V.
NAME: Geocryology-engineering map of Aldan area
PUB: Moscow, Moscow University Publishers, 1975
SCALE: 1:1 000 000
SOURCE: Inset-map in monograph "South Yakutia"/ edited by Kudriavtsev
REGION: South Yakutia
LEGEND Expansion, temperature, thickness, seasonal freezing and thawing, geological and permafrost phenomena, taliks, genesis complexes of ground, water-bearing complexes, types of relief
LAT/LONG: 57°00-60°00/122°00-127°00
INSTITUTE: Moscow State University
NUMBER 00107

AUTHOR Gavrilova, M.K.
NAME: Reconnaissance map of permafrost zones of the world
PUB: Novosibirsk, Nauka, 1981
SCALE: 1:50 000 000
SOURCE: Inset-map in monograph "Contemporary climate and permafrost in continents"
REGION: The Globe
LEGEND Expansion of permafrost (island, continuous and discontinuous permafrost)
INSTITUTE: Permafrost Institute
NUMBER 00108

AUTHOR Gilichinski, D.A.
NAME: Depths of seasonal freezing for the southern part of West-Siberian plain (sketch-map)
PUB: Moscow, Nauka, 1986
SCALE: 1:7 500 000
SOURCE: Inset-map in monograph "Seasonal frozen ground zone in West Siberia" by Gilichinski, D.A.
REGION: West Siberia
LEGEND Count and natural depths of thaw

LAT/LONG: 50°00-61°00/62°00-94°00
 INSTITUTE: Institute of Soil Science and Photosynthesis
 NUMBER 00109

AUTHOR Gilichinski, D.A.
 NAME: Sketch-map (fragment) of ground water depth and mineralization (referring to their influence on seasonal freezing and ground temperature regime)
 PUB: Moscow, Nauka, 1986
 SCALE: 1:7 500 000
 SOURCE: Gilichinski, D.A. Seasonal frozen ground in West Siberia, p.42
 REGION: West Siberia
 LEGEND Bedding in layer of seasonal freezing ground water, which provide inflow moisture to freezing
 LAT/LONG: 50°00-61°00/62°00-94°00
 INSTITUTE: Institute of Soil Science and Photosynthesis
 NUMBER 00110

AUTHOR Gilichinski, D.A.
 NAME: Sketch-map fragment of the phase change of the quantity of heat in seasonal frozen layer
 PUB: Moscow, Nauka, 1986
 SCALE: 1:7 500 000
 SOURCE: Inset-map in monograph "Seasonal frozen ground in West Siberia" by Gilichinski, D.A.
 REGION: West Siberia
 LEGEND Quantity heat (13 ranges)
 LAT/LONG: 50°00-61°00/62°00-94°00
 INSTITUTE: Institute of Soil Science and Photosynthesis
 NUMBER 00111

AUTHOR Gilichinski, D.A.
 NAME: Sketch-map of expenditure of energy on permafrost conditions influencing compensation
 PUB: Moscow, Nauka, 1986
 SCALE: 1:7 500 000
 SOURCE: Inset-map in monograph "Seasonal frozen ground in West Siberia" by Gilichinski, D.A.
 REGION: West Siberia
 LEGEND Expenditure of energy on compensation of influence of permafrost conditions
 LAT/LONG: 50°00-61°00/62°00-94°00
 INSTITUTE: Institute of Soil Science and Photosynthesis
 NUMBER 00112

AUTHOR Gilichinski, D.A.
 NAME: Sketch-map of annual ground heat storage layer thickness, m
 PUB: Moscow, Nauka, 1986
 SCALE: 1:7 500 000
 SOURCE: Inset-map in monograph "Seasonal frozen ground in West Siberia" by Gilichinski, D.A.
 REGION: West Siberia
 LEGEND Annual ground heat storage layer thickness
 LAT/LONG: 50°00-61°00/62°00-94°00
 INSTITUTE: Institute of Soil Science and Photosynthesis
 NUMBER 00113

AUTHOR Gilichinski, D.A.
 NAME: Sketch-map of types of seasonal freezing of soil referring to the average annual ground temperature
 PUB: Moscow, Nauka, 1986
 SCALE: 1:7 500 000
 SOURCE: Inset-map in monograph D.Ç. Gilichinski, "Seasonal frozen ground zone in West Siberia"

REGION: West Siberia
LEGEND Type of seasonal freezing of soils
LAT/LONG: 50°00-61°00/62°00-94°00
INSTITUTE: Institute of Soil Science and Photosynthesis
NUMBER 00114

AUTHOR Gilichinski, D.A.
NAME: Sketch-map of snow cover influence on ground temperature
PUB: Moscow, Nauka, 1989
SCALE: 1:7 500 000
SOURCE: Inset map in monograph "Seasonal frozen ground in West Siberia" by Gilichinski, D.A.
REGION: West Siberia
LEGEND Heat/snow cover influence
LAT/LONG: 50°00-61°00/62°00-94°00
INSTITUTE: Institute of Soil Science and Photosynthesis
NUMBER 00115

AUTHOR Gilichinski, D.A.
NAME: Sketch-map of the depths of potential seasonal thawing
PUB: Moscow, Nauka, 1986
SCALE: 1:7 500 000
SOURCE: Insert-map in monograph "Seasonal frozen ground zone in West Siberia" by Gilichinski, D.A.
REGION: West Siberia
LEGEND Seasonal thawing (11 ranges)
LAT/LONG: 50°00-61°00/62°00-94°00
INSTITUTE: Institute of Soil Science and Photosynthesis
NUMBER 00116

AUTHOR Gilichinski, D.A.
NAME: Zoning referring to the development of frost heaving (fragment)
PUB: Moscow, Nauka, 1986
SCALE: 1:7 500 000
SOURCE: Gilichinski, D.A. Seasonal frozen ground in West Siberia, p.68
REGION: West Siberia
LEGEND Regions with the intensive and weak development of heave processes or absence of them, areas with few frost mounds
LAT/LONG: 50°00-61°00/62°00-94°00
INSTITUTE: Institute of Soil Science and Photosynthesis
NUMBER 00117

AUTHOR Glushkova, O.IU., Degtiarenko, IU.P., Prokhorova, T.P.
NAME: Aerophotogeomorphological map of eastern Verkhne-Khatyrskoi depression surrounding
PUB: Magadan, SVKNII DVO, 1987
SCALE: 1:1 000 000
SOURCE: "Quaternary period in North-East Asia", p.38. In the article by Glushkova, O.IU., Gegtyarenko IU.P., Prokhorov, T.P. p.33-55.
REGION: Kamchatka
LEGEND Gentle slopes of solifluction removal and accumulation; relief, working by cryogenic processes
LAT/LONG: 61°00-64°00/172°00-178°00
NUMBER 00118

AUTHOR Gogichishvili, V.V.
NAME: Sketch-map of cryological landscape indicator near settlement Azei (Irkutsk area)
PUB: Novosibirsk, Nauka, 1983

SCALE: Large
 SOURCE: Leshchikov, F.N., Shats, M.M. South Central Siberia permafrost, p.136-137
 REGION: Central Siberia
 LEGEND: Expansion permafrost ground and short-term permafrost, relief, vegetation associations
 LAT/LONG: 56°00-58°00/102°00-104°00
 INSTITUTE: Permafrost Institute
 NUMBER: 00119

AUTHOR: Gorbunov, A.P.
 NAME: Geocryological map of Dzhungarsk Alatau
 PUB: Moscow, Nauka, 1989
 SCALE: 1:1 500 000
 SOURCE: Geocryology of the USSR. Mountain countries of the southern of USSR / edited by E.E. Ershov, p.301
 REGION: Southern Siberia (Kazakhstan)
 LEGEND: Expansion, temperature, thickness, types of seasonal freezing of soils
 LAT/LONG: 40°00-47°00/75°00-83°00
 NUMBER: 00120

AUTHOR: Gorbunov, A.P.
 NAME: Geocryological map of Saur-Tarbagai
 PUB: Moscow, Nauka, 1989
 SCALE: 1:1 500 000
 SOURCE: Geocryology of the USSR. Mountain countries of the southern USSR / edited by E.D. Ershov, p.297
 REGION: Southern Siberia (Kazakhstan)
 LEGEND: Expansion, temperature, thickness, types of seasonal freezing of soils
 LAT/LONG: 47°00/83°00-85°00
 NUMBER: 00121

AUTHOR: Gorbunov, A.P.
 NAME: Permafrost under Fedchenko Glacier
 PUB: Moscow, Nauka, 1989
 SCALE: Large
 SOURCE: Geocryology of the USSR. Mountain countries of the southern USSR / edited by E.E. Ershov, p.327
 REGION: South-East Central Asia (Pamirs)
 LAT/LONG: 37°00-40°00/66°00-75°00
 NUMBER: 00122

AUTHOR: Gorbunov, A.P.
 NAME: Sketch-map of altitudinal geocryological zonality
 PUB: Moscow, Nauka, 1976
 SCALE: 1:200 000 000
 SOURCE: "Questions of global cryology", p.44
 REGION: Global
 LEGEND: Types of altitudinal zonality (8 ranges), tops with permafrost, absolute height of boundaries, permafrost belts, southern limit of permafrost in lowlands
 INSTITUTE: Permafrost Institute
 NUMBER: 00123

AUTHOR: Gorbunov, A.P.
 NAME: Spreading, thickness and average annual temperature of permafrost in Pamiro-Altai
 PUB: Moscow, Nauka, 1989
 SCALE: 1:4 000 000
 SOURCE: Geocryology of the USSR. Mountain countries of the southern USSR / edited by

REGION: E.E.Ershov, p.308
 LEGEND South-East Central Asia, Tien Shan
 LAT/LONG: Expansion, temperature, thickness
 40°00-43°00/69°00-80°00
 NUMBER 00124

AUTHOR Gorbunov, A.P.
 NAME: Spreading, thickness and average annual temperature of permafrost in Pamiro-Alai
 PUB: Moscow, Nauka, 1989
 SCALE: 1:2 500 000
 SOURCE: Geocryology of the USSR, Mountain countries of the southern USSR / edited by E.E. Ershov, p.325
 REGION: South-East Central Asia (Pamirs)
 LEGEND Expansion, temperature, thickness
 LAT/LONG: 37°00-40°00/66°00-75°00
 NUMBER 00125

AUTHOR Gotovtsev, S.P.
 NAME: Sketch-map of cryogenic processes zoning in Charo-Tokinsky interfluve
 PUB: Yakutsk, Permafrost Institute Publishers, 1983
 SCALE: Large
 SOURCE: Geographical research in Yakutia, p.91
 REGION: Southern Yakutiia
 LEGEND Area of cryogenic processes expansion and topography formation
 LAT/LONG: 58°00-60°00/118°00-120°00
 INSTITUTE: Permafrost Institute
 NUMBER 00126

AUTHOR Gravis, G.F., Drozdov, D.S., Stashenko, A.I.
 NAME: Fragment of engineering-geological map
 PUB: Yakutsk, Geocryological Institute Publishers, 1986
 SCALE: Large
 SOURCE: "Questions of geocryological mapping", p.89. In the article by Gravis, G.F., Drozdov, D.S., Stashenko, A.I. p.85-96.
 REGION: Central Yakutia
 LEGEND Thermokarst, rock stream
 LAT/LONG: 61°30/127°00
 INSTITUTE: VSEGINGEO
 NUMBER 00127

AUTHOR Grigorev, N.F.
 NAME: Climatic regionalization of cryolithozone
 PUB: Moscow, Nauka, 1981
 SCALE: 1:50 000 000
 SOURCE: Gasanov, Sh.Sh. "Cryolithological analysis", p.23
 REGION: The USSR
 LEGEND Permafrost boundary
 LAT/LONG: 50°00-80°00/30°00-70°00
 NUMBER 00128

AUTHOR Gruzlov, A.V., Badu, I.U.B., Lobov, A.P.
 NAME: Map of permafrost average annual temperature distribution
 PUB: Moscow, Moscow University Publishers, 1980
 SCALE: 1:7 500 000
 SOURCE: Trofimov, V.T., Badu I.U.B., Dubikov, G.I. Cryogenic structure and ice content of permafrost in West-Siberian platform, p.34-35

REGION: West Siberia
LEGEND Prevailing temperatures in 8 ranges, places of untouched permafrost
LAT/LONG: 64°00'-74°00'/65°00'-85°00'
INSTITUTE: Moscow State University
NUMBER 00129

AUTHOR Gruzlov, A.V., Badu, I.U.B., Varenyshev, V.B. et.al.
NAME: Map of permafrost thickness in West Siberia platform
PUB: Moscow, Moscow University Publishers, 1980
SCALE: 1:7 500 000
SOURCE: Trofimov, V.T., Badu, I.U.B., Dubikov, G.I. Cryogenic structure and ice content of permafrost in West-Siberian platform, p.46-47

REGION: West Siberia
LEGEND Prevailing permafrost thickness (5 ranges) and their expansion, boundary of territory where wide extent cooling grounds are found or assumed, untouched permafrost places
LAT/LONG: 64°00'-74°00'/65°00'-85°00'
INSTITUTE: Moscow State University
NUMBER 00130

AUTHOR Gruzlov, A.V., Badu, I.U.B., Varenyshev, V.B., Trofimov, V.T., Firsov, N.G.
NAME: Map of permafrost thickness in West Siberia platform
PUB: Novosibirsk, Nauka, 1990
SCALE: 1:3 000 000
SOURCE: Inset-map in monograph "Geocryological investigation history in West Siberia" / edited by Nekrasov

REGION: West Siberia
LEGEND Thickness, areas of discontinuous permafrost, boundaries of areas with cooling ground with cryopegs
LAT/LONG: 62°00'-74°00'/60°00'-90°00'
INSTITUTE: Moscow State University
NUMBER

AUTHOR Gruzlov, A.V., Trofimov, V.T., Filkin, N.A.
NAME: Scheme of distribution of permafrost thickness in Nadym, Taz, Pur river-systems and Tazovski peninsula
PUB: Moscow, Moscow University Publishers, 1977
SCALE: 1:5 000 000
SOURCE: Trofimov, V.T. "Regularities of spatial variability for engineering-geological conditions in West Siberian platform", p.77

REGION: West Siberia
LEGEND Expansion, thickness
LAT/LONG: 64°00'-69°00'/68°00'-88°00'
INSTITUTE: Moscow State University
NUMBER 00132

AUTHOR Gruzlov, A.V.
NAME: Scheme of permafrost extent at the Kola Peninsula
PUB: Moscow, Nauka, 1988
SCALE: 1: 7 500 000
SOURCE: Geocryology of the USSR. European territory of the USSR / edited by E.D. Ershov, p.272
REGION: Kola Peninsula
LEGEND Southern limit of permafrost
LAT/LONG: 66°00'-70°00'/32°00'-42°00'
INSTITUTE: Moscow State University
NUMBER 00133

AUTHOR Ivanova, T.F., Lynov, V.A., Menshikov, L.A. et.al.
NAME: Scheme of Vozei tectonic structure disposition
PUB: Moscow, Stroiizdat, 1984
SCALE: 1:10 000 000
SOURCE: Geocryological conditions and their change forecast in primary development regions in the North, p.135
REGION: European North of the USSR
LEGEND Expansion
LAT/LONG: 64°00-70°00/55°00-65°00
INSTITUTE: PNIIS, Gosstroi USSR
NUMBER 00134

AUTHOR Kachiurin, S.P.
NAME: Map of thermokarst extent in the USSR
PUB: Moscow, Academy of Sciences Publishers, 1961
SCALE: 1:15 000 000
SOURCE: Inset-map in monograph by Kachiurin, S.P. "Thermokarst in the USSR"
REGION: USSR
LEGEND Thermokarst on : wedge and segregation ice (contemporary and relict), hillocky terrain (contemporary and relict), thermokarst terraces, thermoabrasion, palsas, frost mounds
LAT/LONG: 38°00-82°00/20°00-170°00
INSTITUTE: Permafrost Institute
NUMBER 00135

AUTHOR Kaplin, T.N., Leibman, N.O.
NAME: Sketch - map of ice content in permafrost
PUB: Moscow, Stroyizdat, 1990
SCALE: 1:20 000 000
SOURCE: Climate for construction, reference book SNIP, p.51
REGION: USSR
LEGEND Lithological types of ground, ice content, macro-ice content, expansion types of ground ice
LAT/LONG: 38°00-80°00/20°00-170°00
INSTITUTE: Gosstroi USSR
NUMBER 00136

AUTHOR Kaplina, T.N.
NAME: Occurrence and age of syngeneses and epigenesis permafrost in northern Yakutia lowland
PUB: Moscow, Nauka, 1986
SCALE: Large
SOURCE: "Permafrost formation and cryogenic processes forecast", p.10. In article by Kaplina, T.N. p.3-14
REGION: Northern Yakutia
LEGEND Syngeneses permafrost deposits (thick to 40m of Middle Pleistocene age, thick to 5m Holocene), syngeneses and epigenesis permafrost strata
LAT/LONG: 68°00-73°00/110°00-160°00
INSTITUTE: PNIIS, Gosstroi, USSR
NUMBER 00137

AUTHOR Kaplina, T.N., Kostolyndina, N.K., Leibman, M.O.
NAME: Ruggedness of edoma relief by alases in Cukochya-Konkovskii bar region
PUB: Moscow, Nauka, 1986
SCALE: Large
SOURCE: "Permafrost formation and cryogenic processes forecast", p.52. In the article Kaplina, T.N., Kostolyndina N.K., Leibman, M.O. p.51-60
REGION: Northern Yakutia
LEGEND Alas depths (3 ranges), dome-shaped edoma of different altitude (3 ranges)

LAT/LONG: 68°00-69°30/156°00-157°00
INSTITUTE: PNIIS, Gosstroj, USSR
NUMBER 00138

AUTHOR: Kaplina, T.N., Kostalyndina N.K., Leibman, M.O.
NAME: Ruggedness of edoma relief by alases in Duvannyi dome region
PUB: Moscow, Nauka, 1986
SCALE: Large
SOURCE: "Permafrost formation and cryogenic processes forecast", p.54. In the article by Kaplina, T.N., Kostalyndina N.K., Leibman, M.O. p.51-60
REGION: Northern Yakutia
LEGEND: Alas depths (3 ranges), dome-shaped edoma of different altitude (3 ranges)
LAT/LONG: 168°00-169°00/158°00-160°00
INSTITUTE: PNIIS, Gosstroj, USSR
NUMBER 00139

AUTHOR: Kaplina, T.N., Kostalyndina N.K., Leibman, M.O.
NAME: Ruggedness of terrace edoma relief by alases of Chukochya and Bolshoi Konkovo interfluve
PUB: Moscow, Nauka, 1986
SCALE: Large
SOURCE: "Permafrost formation and cryogenic processes forecast", p.56. In the article by Kaplina, T.N., Kostalyndina H.K., Leibman, M.O. p.51-60
REGION: Northern Yakutia
LEGEND: Alas depths (3 ranges), altitude of terrace edoma
LAT/LONG: 69°00-71°00/156°00-161°00
INSTITUTE: PNIIS, Gosstroj, USSR
NUMBER 00140

AUTHOR: Kaplina, T.N., Znamenskii, E.N.
NAME: Schematic map of cryogenic processes and frozen features
PUB: Moscow, Stroiizdat, 1990
SCALE: 1:20 000 000
SOURCE: Climate for construction, reference book SNIP p.50
REGION: USSR
LEGEND: Cryogenic processes and frozen ground features
LAT/LONG: 38°00-80°00/20°00-170°00
INSTITUTE: Gostroi USSR
NUMBER 00141

AUTHOR: Kaplina, T.N., Kostalyndina N.K., Leibman M.O.
NAME: Sketch-map of relief levels in Kolyma lower course
PUB: Moscow, Nauka, 1986
SCALE: 1:5 000 000
SOURCE: "Permafrost formation and cryogenic processes forecast, p.52. In the article by Kaplina, T.N., Kostalyndina N.K., Leibman M.O. p.51-60
REGION: Northern Yakutia
LEGEND: Terrace and dome-shaped edoma of different altitudes, bar and dome names
LAT/LONG: 68°00-71°00/155°00-162°00
INSTITUTE: PNIIS, Gosstroj, USSR
NUMBER 00142

AUTHOR: Kashperuk, P.I., Trofimov, V.T., Firsov, N.G.
NAME: Depths of seasonal ground freezing and thawing
PUB: Moscow, Nedra, 1989
SCALE: 1:20 000 000

- SOURCE:** Geocryology of the USSR. West Siberia / edited by E.D. Ershov, p.82
REGION: West Siberia
LEGEND: Maximum natural depths of seasonal freezing and thawing
LAT/LONG: 50°00'-70°30'/60°00'-90°30'
INSTITUTE: Moscow State University
NUMBER: 00143
- AUTHOR:** Kashperuk, P.I., Trofimov, V.T., Firsov, N.G.
NAME: Seasonal freezing and thawing of ground types referring to lithological and moisture content
PUB: Moscow, Nedra, 1989
SCALE: 1:20 000 000
SOURCE: Geocryology of the USSR. West Siberia / edited by E.D. Ershov, p.79
REGION: West Siberia
LEGEND: Lithological composition, types of depth of thawing and their boundaries
LAT/LONG: 50°00'-70°30'/60°00'-90°30'
INSTITUTE: Moscow State University
NUMBER: 00144
- AUTHOR:** Kashperuk, P.I., Trofimov, V.T., Firsov, N.G.
NAME: Seasonal freezing and thawing of ground types referring to average annual temperature and amplitudes on surface groundmass
PUB: Moscow, Nedra, 1989
SCALE: 1:20 000 000
SOURCE: Geocryology of the USSR. West Siberia / edited by E.D. Ershov, p.77
REGION: West Siberia
LEGEND: Average annual temperature ground types and temperature amplitudes
LAT/LONG: 50°00'-70°30'/60°00'-90°30'
INSTITUTE: Moscow State University
NUMBER: 00145
- AUTHOR:** Katasonov, E.M.
NAME: Geomorphological map of Tumara drainage basin
PUB: Moscow, Academy of Sciences Publishers, 1963
SCALE: 1:300 000 000
SOURCE: "Conditions and permafrost development features in Siberia and North-East", p.8. In the article by Katasonov, E.M. p.5-24
REGION: Central Yakutia
LEGEND: Surface with thermokarst lakes and depression
LAT/LONG: 63°45'-64°30'/129°00'-130°00'
INSTITUTE: Geocryological Institute
NUMBER: 00146
- AUTHOR:** Kaznacheeva, I.A.
NAME: Scheme of exogenesis processes and phenomena in Malo-Bolshezemel region
PUB: Moscow, Nauka, 1988
SCALE: 1:7 500 000
SOURCE: Geocryology of the USSR. European territory of the USSR / edited by E.D. Ershov, p.295
REGION: European North
LEGEND: Cryogenic processes and phenomena (hillocky peatland, thermokarst, thermal erosion, polygonal wedge ice, neogenesis, syngeneses and epigenesis permafrost)
LAT/LONG: 65°00'-69°00'/52°00'-64°00'
INSTITUTE: SO NIIOSP
NUMBER: 00147
- AUTHOR:** Kaznacheeva, I.A.

NAME: Scheme of polygonal ground types extent associated with their relief forms in Malo-Boshezemesky region
 PUB: Moscow, Nedra, 1988
 SCALE: 1:5 000 000
 SOURCE: Geocryology of the USSR. European territory of the USSR / edited by E.D. Ershov, p.297
 REGION: European North
 LEGEND: Contemporary syngensis and epigenesis wedge ice, buried wedge ice, pseudomorphs, microrelief in growth and stabilization phases
 LAT/LONG: 66°00-69°00/48°00-64°00
 INSTITUTE: SO NIIOSP
 NUMBER: 00148

AUTHOR: Kaznacheeva, I.A., Sukhodolskii, S.E.
 NAME: Zoning scheme of Malo-Bolshezemel'sky region referring to discontinuous permafrost
 PUB: Moscow, Nauka, 1988
 SCALE: 1:5 000 000
 SOURCE: Geocryology of the USSR. European territory of the USSR / edited by E.D. Ershov, p.278
 REGION: European North
 LEGEND: Permafrost zone referring to expansion
 LAT/LONG: 48°00-66°00/64°00-72°00
 INSTITUTE: SO NIIOSP
 NUMBER: 00149

AUTHOR: Khrustalev, L.H., Novikov, F.I.A., Nadessdin, A.V., Maksimenko, A.S.
 NAME: Sketch-map of natural complexes resistivity impact
 PUB: Moscow, Nedra, 1988
 SCALE: 1:15 000 000
 SOURCE: Geocryology of the USSR. European territory of the USSR / edited by E.D. Ershov, p.343
 REGION: European North
 LEGEND: Temperature, expansion, ice content, permafrost and unfrozen ground thermal characteristics, characteristics of resistance referring to the potential exhibit of thermokarst and heave
 LAT/LONG: 65°00-74°00/32°00-68°00
 INSTITUTE: SO NIIOSP
 NUMBER: 00150

AUTHOR: Khrutskii, S.F., Afanasenko, V.E., Kondratieva, K.A.
 NAME: Hydrogeological zoning scheme
 PUB: Moscow, Moscow University Publishers, 1972
 SCALE: 1:1 000 000
 SOURCE: Permafrost Studies, XII, p.60
 REGION: Central Siberia
 LEGEND: Hydrogeological structures (cryogenic massif), taliks, watery characteristics, geological structure
 LAT/LONG: 69°00-71°00/140°00-143°00
 INSTITUTE: Moscow State University
 NUMBER: 00151

AUTHOR: Khrutskii, S.F., Afanasenko V.E., Kondratieva, K.A.
 NAME: Permafrost sketch-map
 PUB: Moscow, Moscow University Published, 1972
 SCALE: 1:1 000 000
 SOURCE: Permafrost Studies, XII, p.55
 REGION: Central Siberia
 LEGEND: Permafrost thickness, average annual temperature of ground, morphostructures, genesis and lithological content

LAT/LONG: 69°00-71°00/140°00-143°00
 INSTITUTE: Moscow State University
 NUMBER 00152

AUTHOR Koldysheva, R.IA.
 NAME: Geocryological sketch-map of Tunkinskaya depression
 PUB: Yakutsk, Yakutskoe knizh. izd-vo, 1966
 SCALE: 1:1 500 000
 SOURCE: "Reports of VIII All-Union Geocryological Conference", issue 3. Regional geocryology, p.140. In the report by Koldysheva, R.IA. p.137-142
 REGION: Transbaikal
 LEGEND Permafrost thickness, the depths of permafrost table in wells, boundaries of zone (real, assumed): zones without permafrost, permafrost thickness 4-40 m, 40-215 m with taliks islands
 LAT/LONG: 51°00-52°00/101°00-104°00
 NUMBER 00153

AUTHOR Koldysheva, R.IA.
 NAME: Scheme (fragment) of aeration zone for permafrost
 PUB: Moscow, Nauka, 1977
 SCALE: 1:7 500 000
 SOURCE: "Frozen ground and snow cover", p.169
 REGION: Transbaikal
 LEGEND Boundaries of continuous, discontinuous and island permafrost
 LAT/LONG: 52°00-54°00/95°00-115°00
 NUMBER 00154

AUTHOR Koldysheva, R.IA.
 NAME: Scheme of general types of water exchange in permafrost aeration zone
 PUB: Moscow, Nauka, 1977
 SCALE: 1:7 500 000
 SOURCE: "Frozen ground and snow cover", p.178
 REGION: Transbaikal
 LEGEND Infiltration type with rather high energy exchange in system "atmosphere-ground aeration zone-suprapermafrost waters-permafrost"
 LAT/LONG: 53°00-54°00/95°00-115°00
 NUMBER 00155

AUTHOR Kondrateva, K.A., Kudriavtsev, V.A.
 NAME: Cryolithozone map of the USSR
 PUB: Moscow, Enlightenment, 1980
 SCALE: 1:35 000 000
 SOURCE: Romanovskii, N.N. "Cold of the Earth: Textbook for student", p.18-19
 REGION: The USSR
 LEGEND Thickness, temperature, spreading. Ocean, shelf, northern, southern, subglacial cryolithozone. Relict cryolithozone, the boundary of syngenesisal freezing ground of "ice complex"
 LAT/LONG: 38°00-82°00/20°00-170°00
 INSTITUTE: Moscow State University
 NUMBER 00156

AUTHOR Kondratieva, K.A., Khrutskii, S.F.
 NAME: Air temperature amplitudes map for the Central Siberia
 PUB: Moscow, Nedra, 1989
 SCALE: 1:20 000 000
 SOURCE: Geocryology of the USSR. Central Siberia / edited by E.D. Ershov, p.41

REGION: Central Siberia
 LEGEND: Southern and northern limit of permafrost zones, contemporary permafrost expansion
 LAT/LONG: 50°00-75°00/80°00-135°00
 INSTITUTE: Moscow State University
 NUMBER: 00157

AUTHOR: Kondratieva, K.A.
 NAME: Average annual ground temperature inversion distribution in river valley
 PUB: Moscow, Nedra, 1989
 SCALE: Large
 SOURCE: Geocryology of the USSR. Central Siberia / edited by E.D. Ershov, p.66
 REGION: Central Siberia (Vilyui River)
 LEGEND: Average annual ground temperatures, lithological composition
 LAT/LONG: 62°00-64°00/110°00-130°00
 INSTITUTE: Moscow State University
 NUMBER: 00158

AUTHOR: Kondratieva, K.A.
 NAME: Central Siberia cryolithology map
 PUB: Moscow, Nedra, 1989
 SCALE: 1:20 000 000
 SOURCE: Geocryology of the USSR. Central Siberia / edited by E.D. Ershov, p.88
 REGION: Central Siberia
 LEGEND: Type of freezing, lithological composition, cryogenic structure, macro ice inclusions, volumetric ice content, expansion, southern limit of permafrost
 LAT/LONG: 50°00-75°00/80°00-135°00
 INSTITUTE: Moscow State University
 NUMBER: 00159

AUTHOR: Kondratieva, K.A.
 NAME: Central Siberia engineering-geocryological regionalization map
 PUB: Moscow, Nedra, 1989
 SCALE: 1:20 000 000
 SOURCE: Geocryology of the USSR. Central Siberia / edited by E.D. Ershov, p.391
 REGION: Central Siberia
 LEGEND: Permafrost zone boundaries, region numbers (the table gives the characteristics of engineering-geocryology), the tendency for development of cryogenic processes as a result of construction
 LAT/LONG: 50°00-75°00/80°00-135°00
 INSTITUTE: Moscow State University
 NUMBER: 00160

AUTHOR: Kondratieva, K.A., Fotiev, S.M.
 NAME: Central Siberia permafrost regionalization referring to types of permafrost zone structures
 PUB: Moscow, Nedra, 1989
 SCALE: 1:35 000 000
 SOURCE: Geocryology of the USSR. Central Siberia / edited by E.D. Ershov, p.97
 REGION: Central Siberia
 LEGEND: Stages of frozen, frost, cool, with saline ground water and practically anhydrous ground, boundaries of districts with different permafrost structure, geocryological zones, contemporary permafrost expansion
 LAT/LONG: 50°00-75°00/80°00-135°00
 INSTITUTE: Moscow State University, PNIIS, Gosstroj USSR
 NUMBER: 00161

AUTHOR Kondratieva, K.A.
 NAME: Central Siberia regionalization referring to permafrost
 PUB: Moscow, Nedra, 1989
 SCALE: 1:20 000 000
 SOURCE: Geocryology of the USSR. Central Siberia / edited by E.D. Ershov, p.130
 REGION: Central Siberia
 LEGEND Morphostructure and geocryological features, southern limit of contemporary permafrost
 LAT/LONG: 50°00-75°00/80°00-135°00
 INSTITUTE: Moscow State University
 NUMBER 00162

AUTHOR Kondratieva, K.A., Danilova, N.S.
 NAME: Cryogenic age map (onset of ground freezing) in Central Siberia
 PUB: Moscow, Nedra, 1989
 SCALE: 1:20 000 000
 SOURCE: Geocryology of the USSR. Central Siberia / edited by E.D. Ershov, p.30
 REGION: Central Siberia
 LEGEND Pleistocene, Holocene, recent (seasonal freezing of soil), age of ground, southern limit of permafrost in Pliocene and in present
 LAT/LONG: 50°00-75°00/80°00-135°00
 INSTITUTE: Moscow State University, PNIIS, Gosstroj USSR
 NUMBER 00163

AUTHOR Kondratieva, K.A., Oberman, N.G., Sukhodolskii, S.E.
 NAME: European North of the USSR zoning scheme referring to subareal permafrost thickness and vertical structure
 PUB: Moscow, Nauka, 1988
 SCALE: 1:10 000 000
 SOURCE: Geocryology of the USSR. European territory of the USSR / edited by E.D. Ershov, p.248
 REGION: European North of the USSR
 LEGEND Shallow down permafrost thickness; relict permafrost under unfrozen ground; permafrost thickness with cryopegs occurring under permafrost
 LAT/LONG: 60°00-85°00/40°00-60°00
 INSTITUTE: Moscow State University, PUGRO, PNIIS, Gosstroj USSR
 NUMBER 00164

AUTHOR Kondratieva, K.A.
 NAME: Frozen and unfrozen ground expansion and their average annual temperature (°C)
 PUB: Moscow, Nedra, 1989
 SCALE: 1:5 000 000
 SOURCE: Geocryology of the USSR. Central Siberia / edited by E.D. Ershov, p.222
 REGION: Central Siberia
 LEGEND Permafrost expansion and temperature
 LAT/LONG: 60°00-68°00/90°00-101°00
 INSTITUTE: Moscow State University
 NUMBER 00165

AUTHOR Kondratieva, K.A.
 NAME: Geobotanical zones with dominant vegetation in Central Siberia
 PUB: Moscow, Nedra, 1989
 SCALE: 1:35 000 000
 SOURCE: Geocryology of the USSR. Central Siberia / edited by E.D. Ershov, p.47
 REGION: Central Siberia
 LEGEND Northern and southern geocryological zones boundary, southern limit of permafrost
 LAT/LONG: 50°00-75°00/80°00-135°00
 INSTITUTE: Moscow State University

NUMBER 00166

AUTHOR NAME: Kondratieva, K.A., Kudriavtsev, V.A., Khrutskii, S.F.
PUB: Geocryological- hydrogeological map of southern Yakutia
Moscow, Moscow University Publishers, 1967
SCALE: 1:2 500 000
SOURCE: Permafrost Studies, VII, p.100-101
REGION: Southern Yakutia
LEGEND: Types of underground water referring to permafrost, permafrost expansion and thickness, water-bearing complexes and special forms of underground water accumulation and display
LAT/LONG: 55°00-60°00/120°-135°00
INSTITUTE: Moscow State University
NUMBER 00167

AUTHOR NAME: Kondratieva, K.A., Maksimova, L.N.
PUB: Geocryological map of Baikal-Patom region
Moscow, Nedra, 1989
SCALE: 1:5 000 000
SOURCE: Geocryology of the USSR. Mountain territories of south USSR / edited by E.D. Ershov, p.68
REGION: Transbaikal
LEGEND: Expansion, temperature, thickness
LAT/LONG: 57°00-61°00/110°00-116°00
INSTITUTE: Moscow State University
NUMBER 00168

AUTHOR NAME: Kondratieva, K.A.
PUB: Geotemperature map of Malaya and Bolshaya Botuobiya interfluve
Moscow, Nedra, 1989
SCALE: Large
SOURCE: Geocryology of the USSR. Central Siberia / edited by E.D. Ershov, p.237
REGION: Central Siberia
LEGEND: Temperature, age of deposits
LAT/LONG: 62°00-63°00/114°00-115°00
INSTITUTE: Moscow State University
NUMBER 00169

AUTHOR NAME: Kondratieva, K.A.
PUB: Map of average annual ground temperatures and ice sheets in Central Siberia
Moscow, Nedra, 1989
SCALE: 1:20 000 000
SOURCE: Geocryology of the USSR. Central Siberia / edited by E.D. Ershov, p.61
REGION: Central Siberia
LEGEND: Average annual temperature of ground, geocryological zones boundary, southern limit of permafrost
LAT/LONG: 50°00-75°00/80°00-135°00
INSTITUTE: Moscow State University
NUMBER 00170

AUTHOR NAME: Kondratieva, K.A., Dunaeva E.N.
PUB: Map of average annual permafrost temperature in the USSR
Moscow, Nedra, 1988
SCALE: 1:20 000 000
SOURCE: Inset-map in monograph "Geocryology of the USSR. European territory of the USSR / edited by E.D. Ershov"

REGION: USSR
LEGEND Seasonal frozen ground and permafrost expansion, temperature, boundaries of temperature and permafrost zone
LAT/LONG: 38°00-82°00/30°00-170°00
INSTITUTE: Moscow State University
NUMBER 00171

AUTHOR Kondratieva, K.A., Kudriavtsev, V.A., Khrutskii, S.F.
NAME: Map of average annual ground temperature zoning of Yano-Indigirskii interfluvial
PUB: Moscow, Moscow University Publishers, 1972
SCALE: 1:2 500 000
SOURCE: Permafrost Studies, XII, p.70
REGION: Central Siberia
LEGEND Average annual ground temperature zone, geological-genesis complexes
LAT/LONG: 67°30-73°00/138°00-144°00
INSTITUTE: Moscow State University
NUMBER 00172

AUTHOR Kondratieva, K.A.
NAME: Map of frozen ground features expansion in geology-genesis types and formation
PUB: Moscow, Nedra, 1989
SCALE: 1:20 000 000
SOURCE: Geocryology of the USSR. Central Siberia / edited by E.D. Ershov, p.124
REGION: Central Siberia
LEGEND Soil wedges, wedge ice, frost mounds, hillocky peatland, sheet ice, hillocky terrain, thermokarst, frost fracturing, cave ice, glaciers
LAT/LONG: 50°00-75°00/80°00-135°00
INSTITUTE: Moscow State University
NUMBER 00173

AUTHOR Kondratieva, K.A.
NAME: Map of generalized permafrost zones thickness in Central Siberia
PUB: Moscow, Nedra, 1989
SCALE: 1:20 000 000
SOURCE: Geocryology of the USSR. Central Siberia \ edited by E.D. Ershov, p.105
REGION: Central Siberia
LEGEND Permafrost thickness in geocryological zones and shelf zone
LAT/LONG: 50°00-75°00/80°00-135°00
INSTITUTE: Moscow State University
NUMBER 00174

AUTHOR Kondratieva, K.A.
NAME: Map of North-Siberia lowland permafrost zone thickness (m)
PUB: Moscow, Nedra, 1989
SCALE: 1:15 000 000
SOURCE: Geocryology of the USSR. Central Siberia / edited by E.D. Ershov, p.156
REGION: Central Siberia
LEGEND Permafrost thickness (8 ranges), permafrost with cryopegs boundary (permafrost thickness 100-300 and 0-100 m)
LAT/LONG: 70°00-75°00/85°00-130°00
INSTITUTE: Moscow State University
NUMBER 00175

AUTHOR Kondratieva, K.A.
NAME: Map of permafrost thickness in the USSR
PUB: Moscow, Nedra, 1988

SCALE: 1:20 000 000
 SOURCE: Inset-map in monograph "Geocryology of the USSR. European territory of the USSR" / edited by E.D. Ershov
 REGION: USSR
 LEGEND: Thickness, permafrost zones boundaries : subareal, subglacial, southern limit of relict and contemporary permafrost
 LAT/LONG: 38°00-82°00/30°00-170°00
 INSTITUTE: Moscow State University
 NUMBER: 00176

AUTHOR: Kondratieva, K.A.
 NAME: Map of permafrost expansion and thickness (m) of upper stage in Central Siberia
 PUB: Moscow, Nedra, 1989
 SCALE: 1:20 000 000
 SOURCE: Geocryology of the USSR. Central Siberia / edited by E.D. Ershov, p.104
 REGION: Central Siberia
 LEGEND: Permafrost thickness in discontinuous and continuous permafrost zones (in valleys and on divide), permafrost zones boundaries
 LAT/LONG: 50°00-75°00/80°00-135°00
 INSTITUTE: Moscow State University
 NUMBER: 00177

AUTHOR: Kondratieva, K.A., Khrutskii, S.F.
 NAME: Map of permafrost expansion and thickness (m) and cooling below 0° ground with cryopegs, bedding under permafrost layer
 PUB: Moscow, Nedra, 1989
 SCALE: 1:35 000 000
 SOURCE: Geocryology of the USSR. Central Siberia / edited by E.D. Ershov, p.107
 REGION: Central Siberia
 LEGEND: Thickness
 LAT/LONG: 50°00-75°00/80°00-135°00
 INSTITUTE: Moscow State University
 NUMBER: 00178

AUTHOR: Kondratieva, K.A.
 NAME: Map of permafrost in the USSR
 PUB: Moscow, Moscow University Publishers, 1978
 SCALE: 1:35 000 000
 SOURCE: General Permafrost Studies (Geocryology) / edited by Kudriavtsev, V.A., p.14
 REGION: USSR
 LEGEND: Expansion, temperature, thickness
 LAT/LONG: 38°00-82°00/30°00-170°00°
 INSTITUTE: Moscow State University
 NUMBER: 00179

AUTHOR: Kondratieva, K.A., Kudriavtsev, V.A.
 NAME: Map of permafrost zoning of the USSR
 PUB: Moscow, Moscow University Publishers, 1979
 SCALE: 1:40 000 000
 SOURCE: Inset-map in monograph "Geocryological survey methods" / edited by V.A. Kudriavtsev
 REGION: USSR
 LEGEND: Northern and Southern permafrost zones (expansion, temperature, thickness), cryopeg zones, two layer stratas, the regular repetition of seasonal freezing soil zones
 LAT/LONG: 38°00-82°00/30°00-170°00
 INSTITUTE: Moscow State University
 NUMBER: 00180

AUTHOR Kondratieva, K.A., Danilova, N.S., Baulin, V.V.
 NAME: Map of permafrost age
 PUB: Moscow, Nedra, 1988
 SCALE: 1:20 000 000
 SOURCE: Inset-map in monograph "Geocryology of the USSR. European territory of the USSR / edited by E.D. Ershov"
 REGION: USSR
 LEGEND: Expansion and geological age, southern limit of permafrost in Pliocene, Pleistocene, present
 LAT/LONG: 38°00-82°00/30°00-170800
 INSTITUTE: Moscow State University, PNIIS, Gosstrois USSR
 NUMBER 00181

AUTHOR Kondratieva, K.A.
 NAME: Map of Severnaya Zemlya average annual ground and glacial cover temperatures
 PUB: Moscow, Nedra, 1989
 SCALE: 1:2 000 000
 SOURCE: Geocryology of the USSR. Central Siberia / edited by E.D. Ershov, p.136
 REGION: The North Polar region
 LEGEND: Temperature in 6 ranges
 LAT/LONG: 82°00-87°00/90°00-110°00
 INSTITUTE: Moscow State University
 NUMBER 00182

AUTHOR Kondratieva, K.A.
 NAME: Map of Severnaya Zemlya glacial cover thickness
 PUB: Moscow, Nedra, 1989
 SCALE: 1:2 000 000
 SOURCE: Geocryology of the USSR. Central Siberia / edited by E.D. Ershov, p.135
 REGION: Central Siberia
 LEGEND: Thickness
 LAT/LONG: 82°00-87°00/90°00-110°00
 INSTITUTE: Moscow State University
 NUMBER 00183

AUTHOR Kondratieva, K.A., Khrutskii S.F.
 NAME: Map of snow cover thickness in Central Siberia
 PUB: Moscow, Nedra, 1989
 SCALE: 1:35 000 000
 SOURCE: Geocryology of the USSR. Central Siberia / edited by E. D. Ershov, p.45
 REGION: Central Siberia
 LEGEND: Southern and northern limit of permafrost zones, temporary permafrost expansion, snow cover thickness
 LAT/LONG: 50°00-75°00/80°00-135°00
 INSTITUTE: Moscow State University
 NUMBER 00184

AUTHOR Kondratieva, K.A.
 NAME: Map of Taimyr (northern part) permafrost zone thickness (m)
 PUB: Moscow, Nedra, 1989
 SCALE: 1:10 000 000
 SOURCE: Geocryology of the USSR. Central Siberia / edited by E.D. Ershov, p.144
 REGION: Central Siberia
 LEGEND: Permafrost thickness (7 ranges)
 LAT/LONG: 74°00-78°00/80°00-115°00
 INSTITUTE: Moscow State University

NUMBER 00185
 AUTHOR Kondratieva, K.A.
 NAME: Map of Taimyr (northern part) permafrost zone average annual ground temperatures
 PUB: Moscow, Nedra, 1989
 SCALE: 1:10 000 000
 SOURCE: Geocryology of the USSR. Central Siberia / edited by E.D. Ershov, p.144
 REGION: Central Siberia
 LEGEND Temperature (4 ranges)
 LAT/LONG: 74°00-78°00/80°00-115°00
 INSTITUTE: Moscow State University
 NUMBER 00186
 AUTHOR Kondratieva, K.A.
 NAME: Map of temperature macrozones in Central Siberia
 PUB: Moscow, Nedra, 1989
 SCALE: 1:35 000 000
 SOURCE: Geocryology of the USSR. Central Siberia / edited by E.D. Ershov, p.67
 REGION: Central Siberia
 LEGEND Prevalent average annual temperatures of ground, permafrost zone boundaries, southern limit of permafrost
 LAT/LONG: 50°00-75°00/80°00-135°00
 INSTITUTE: Moscow State University
 NUMBER 00187
 AUTHOR Kondratieva, K.A.
 NAME: Map of Tungus-Vilyui region permafrost thickness (m)
 PUB: Moscow, Nedra, 1989
 SCALE: 1:5 000 000
 SOURCE: Geocryology of the USSR. Central Siberia / edited by E.D. Ershov, p.222
 REGION: Central Siberia
 LEGEND Thickness in southern (4 ranges) and northern (5 ranges) permafrost zones
 LAT/LONG: 60°00-68°00/90°00-101°00
 INSTITUTE: Moscow State University
 NUMBER 00188
 AUTHOR Kondratieva, K.A.
 NAME: North-Siberia lowland average annual ground temperature (°C)
 PUB: Moscow, Nedra, 1989
 SCALE: 1:15 000 000
 SOURCE: Geocryology of the USSR. Central Siberia / edited by E.D. Ershov, p.151
 REGION: Central Siberia
 LEGEND Temperature (5 ranges)
 LAT/LONG: 70°00-75°00/85°00-130°00
 INSTITUTE: Moscow State University
 NUMBER 00189
 AUTHOR Kondratieva, K.A.
 NAME: Permafrost and glaciers average annual temperatures (°C) map in Novaya Zemlya archipelago
 PUB: Moscow, Nauka, 1988
 SCALE: 1:3 500 000
 SOURCE: Geocryology of the USSR. European territory of USSR / edited by E.D. Ershov, p.266
 REGION: Novaya Zemlya
 LEGEND Temperature spans in cold and warm years
 LAT/LONG: 70°00-77°00/51°00-72°00

INSTITUTE: Moscow State University
 NUMBER 00190

AUTHOR Kondratieva, K.A.
 NAME: Permafrost and glaciers average annual temperatures (°C) map in Zemlya Frantsa Iosifa
 PUB: Moscow, Nauka, 1988
 SCALE: 1:20 000 000
 SOURCE: Geocryology of the USSR. European territory of the USSR / edited by E.D. Ershov, p.263
 REGION: Franz Jozef Land
 LEGEND Temperature
 LAT/LONG: 76°00-84°00/42°00-66°00
 INSTITUTE: Moscow State University
 NUMBER 00191

AUTHOR Kondratieva, K.A.
 NAME: Permafrost and seasonal frozen ground expansion map
 PUB: Moscow, Nauka, 1988
 SCALE: 1:50 000 000
 SOURCE: Geocryology of the USSR. European territory of the USSR / edited by E.D. Ershov, p.100
 REGION: USSR
 LEGEND Expansion
 LAT/LONG: 38°00-82°00/20°00-170°00
 INSTITUTE: Moscow State University
 NUMBER 00192

AUTHOR Kondratieva, K.A., Gavrilov, A.V.
 NAME: Permafrost zoning scheme of the USSR
 PUB: Moscow, Nauka, 1988
 SCALE: 1:40 000 000
 SOURCE: Geocryology of the USSR. European territory of the USSR / edited by E.D. Ershov, p.159
 REGION: USSR
 LEGEND Figures are shown for the first and second order regions
 LAT/LONG: 35°00-82°00/30°00-170°00
 INSTITUTE: Moscow State University
 NUMBER 00193

AUTHOR Kondratieva, K.A., Khrutskii, S.F.
 NAME: Sketch-map of the permafrost zone base depth
 PUB: Moscow, Nedra, 1989
 SCALE: 1:20 000 000
 SOURCE: Geocryology of the USSR. Central Siberia / edited by E.D. Ershov, p.102
 REGION: Central Siberia
 LEGEND Permafrost zones base bedding below or above sea level line
 LAT/LONG: 50°00-75°00/80°00-135°00
 INSTITUTE: Moscow State University
 NUMBER 00194

AUTHOR Koniakhin, M.A.
 NAME: Air temperature: air, mean winter day near surface distribution and oxygen-isotope content of contemporary wedge ice in territory of former USSR
 PUB: Moscow, Moscow University Publisher, 1992
 SCALE: 1:40 000 000
 SOURCE: "Geoecology of the North (Introduction to Geocryoeology)"/ Edited by V.I. Solomatin, p.45
 REGION: The U.S.S.R.
 LEGEND Isotope-oxygenous content of recent wedge ice, mean temperature of mean winter air
 LAT/LONG: 50°00-80°00/40°00-170°00

INSTITUTE: Moscow State University
 NUMBER 00195

AUTHOR Koniakhin, M.A.
 NAME: Sheet ice bedding conditions (territory of Bovanenkovo GKM)
 PUB: Moscow, Moscow State University, 1992
 SCALE: Large
 SOURCE: "Geocology of the North (Introduction to Geocryocology)"/ Edited by V.I. Solomatin, p.47
 REGION: West Siberia, Yamal Peninsula
 LEGEND Contour lines of absolute height of sheet ice table. Absolute points of sheet ice base. Stripping sheet ice wells.
 LAT/LONG: 70°00/70°00
 INSTITUTE: Moscow State University
 NUMBER 00196

AUTHOR Kononova R.S., Neizvestnov, I.A.V., Tolstikhin, N.I.
 NAME: Boundaries of seasonal cryopeg expansion in winter (World Ocean)
 PUB: Moscow, Moscow University Publishers, 1971
 SCALE: 1:200 000 000
 SOURCE: Permafrost Studies, XI, p.78
 REGION: The Globe
 LEGEND Seasonal cryopegs
 NUMBER 00197

AUTHOR Kostiaev, A.G.
 NAME: Scheme of permafrost limit in Eurasia in maximum cold period
 PUB: Moscow, Moscow University Publishers, 1965
 SCALE: 1:80 000 000
 SOURCE: "Ground ice", issue II / edited by A.I. Popov, p.12
 REGION: Eurasia
 LEGEND Southern limit of permafrost, southern limit of permafrost in Europe if glacial cover was absent, area of sedimentation with syngeneses freezing and formation of thick wedge ice, glaciers, periglacial zone of loess surface formations
 LAT/LONG: 20°00-85°00/20°00-170°00
 INSTITUTE: Moscow State University
 NUMBER 00198

AUTHOR Kozlova, A.E.
 NAME: Geomorphological sketch-map of Taz drainage basin
 PUB: Moscow, Nauka, 1972
 SCALE: 1:2 500 000
 SOURCE: Inset-map in monograph "Natural conditions of the Tazovskii oil-gas containing region industrialization"
 REGION: West Siberia
 LEGEND Polygonal relief, frost mounds, thermokarst depressions
 LAT/LONG: 62°30-68°00/78°00-84°00
 INSTITUTE: Geographical Institute
 NUMBER 00199

AUTHOR Kritsuk, L.N., Mel'nikov, E.S., Moskalenko, N.G.
 NAME: Permafrost regionalization scheme for West Siberia referring to permafrost resistance to technological exposure (on landscape base)
 PUB: Yakutsk, Geocryological Institute Publishers, 1986
 SCALE: 1:15 000 000
 SOURCE: "Questions of geocryological mapping", p.65. In the article by Kritsuk, L.N., Mel'nikov, E.S., Moskalenko, N.G. p.53-67

REGION: West Siberia
 LEGEND: Permafrost resistance at the expense of ice content, temperature and sheet ice
 LAT/LONG: 60°00-74°00/60°00-85°00
 INSTITUTE: VSEGINGEO
 NUMBER: 00200

AUTHOR: Kudriavtsev, V.A., Kondratieva, K.A.
 NAME: Geocryological map of USSR
 SCALE: 1:2 500 000
 REGION: USSR
 LEGEND: Complex of geological conditions, lithological composition, cryogenic structure, thickness, expansion, ice content, temperature, cryogenic processes and frozen ground features, depth of bedding and thickness of deposits with cryopegs
 LAT/LONG: 40°00-82°00/30°00-170°00
 INSTITUTE: Moscow State University
 NUMBER: 00201

AUTHOR: Kudriavtsev, V.A., Baulin, V.V., Gruzlov, A.V. /edited by E.M.Sergeev
 NAME: Map of permafrost conditions of Russian non-chernozem zone (except Ural, Zaurale, Kaliningrad region)
 PUB: Moscow, GUGK, 1984
 SCALE: 1:1 500 000
 REGION: Non-chernozem zone, Russia
 LEGEND: Genesis complexes, lithological composition, temperature, annual amplitude ground temperature fluctuations
 LAT/LONG: 52°00-69°00/28°00-64°00
 INSTITUTE: Moscow State University, Geological Ministry, Ministry of Education (Russia)
 NUMBER: 00202

AUTHOR: Kudriavtsev, V.N., Kondratieva, K.A., Vitkina, N.Kh.
 NAME: Seasonal freezing of soils type map
 PUB: Moscow, Moscow University Publishers, 1962
 SCALE: Large
 SOURCE: Inset-map in symposium articles "Permafrost Studies, II"
 REGION: Central Siberia
 LEGEND: Average annual ground temperature, the depth of seasonal freezing (natural and settlement), temperature amplitudes above soils, prevalent lithological composition of winter freezing layer
 LAT/LONG: 52°00-58°00/95°00-104°00
 INSTITUTE: Moscow State University
 NUMBER: 00203

AUTHOR: Kudryashov, V.G., Trofimov, V.T.
 NAME: Permafrost base in West Siberian platform
 PUB: Moscow, Nedra, 1989
 SCALE: 1:20 000 000
 SOURCE: Geocryology of the USSR. West Siberia / edited by E.D. Ershov, p.123
 REGION: West Siberia
 LEGEND: Depth of permafrost base
 LAT/LONG: 50°00-70°30/60°00-90°30
 INSTITUTE: Moscow State University
 NUMBER: 00204

AUTHOR: Kuznetsova, I.L., Parmuzin, S.IU., Rogatina, N.P.
 NAME: Critical depth of lakes with different thermal resistance to snow cover (2 maps)
 PUB: Moscow, Stroizdat, 1987

SCALE: 1:15 000 000
 SOURCE: "Recommendations to estimate the change of permafrost conditions in industrialization territories of West Siberia / PNIIS, Gosstro, USSR", p.22-23
 REGION: West Siberia
 LEGEND: The depth of lakes with different thermal resistance (5 range)
 LAT/LONG: 62°00-75°00/64°00-86°00
 INSTITUTE: PNIIS, Gosstro, USSR
 NUMBER: 00205

AUTHOR: Kuznetsova, I.L., Parmuzin, S.IU., Rogatina N.P.
 NAME: Critical snow cover thickness with different thermal resistance of vegetation in winter and summer (3 maps)
 PUB: Moscow, Stroiizdat, 1987
 SCALE: 1:15 000 000
 SOURCE: "Recommendations to estimate the change of permafrost conditions in industrialization territories of West Siberia" / PNIIS, Gosstro, USSR, p.18-20
 REGION: West Siberia
 LEGEND: Critical snow cover thickness on places with clayey, sand soil and peat for different thermal resistance of vegetation cover
 LAT/LONG: 60°00-80°00/60°00-85°00
 INSTITUTE: PNIIS, Gosstro, USSR
 NUMBER: 00206

AUTHOR: Kuznetsova, I.L.
 NAME: Engineering-geocryological sketch-map (fragments) of Yakutia region littoral lowlands
 PUB: Moscow, Stroiizdat, 1984
 SCALE: Large
 SOURCE: "Cryogenic processes and features". Collection scientific articles / edited by E.A. Vtyurina, p.52-53
 REGION: North-East
 LEGEND: Ice content in the upper and lower horizon of sediments, wedge ice, thermokarst, thermoerosion, ice content (macro), cryopegs, geological genesis complexes
 LAT/LONG: 64°00-74°00/130°00-162°00
 INSTITUTE: PNIIS, Gosstro, USSR
 NUMBER: 00207

AUTHOR: Kuznetsova, I.L., Parmuzin, S.IU., Rogatina, N.P.
 NAME: Regionalization map of the north of West Siberia referring to potential perennial freezing ground with removal of vegetation cover
 PUB: Moscow, Stroiizdat, 1987
 SCALE: 1:12 000 000
 SOURCE: "Recommendation to estimate the change of permafrost conditions in industrialization territories of West Siberia/PNIIS, Gosstro, USSR, p.21
 REGION: North of West Siberia
 LEGEND: Number of zones (explanation in text p.17-18)
 LAT/LONG: 62°00-75°00/64°00-86°00
 INSTITUTE: PNIIS, Gosstro, USSR
 NUMBER: 00208

AUTHOR: Kuznetsova, I.L.
 NAME: Sketch-map of engineering-geocryological zonality in littoral lowlands
 PUB: Moscow, Stroiizdat, 1984
 SCALE: 1:10 000 000
 SOURCE: "Cryogenic processes and features". Collection scientific articles / edited by Vtiurina, E.A., p.58
 REGION: North-East

LEGEND Boundaries and numbers of engineering-geocryological zones, temperature zones
 LAT/LONG: 67°00-74°00/130°00-162°00
 INSTITUTE: PNIIS, Gosstro, USSR
 NUMBER 00209

AUTHOR Lakhtina, O.V., Sukhodolskaia, L.A.
 NAME: Sketch-map of permafrost extent in Holocene climatic optimum
 PUB: Moscow, Nauka, 1981
 SCALE: 1:12 000 000
 SOURCE: "History of permafrost development in Eurasia", p.123. In the article by Lakhtina, O.V., Sukhodolskaia, L.A. p.113-125
 REGION: Transbaikal
 LEGEND Permafrost spread
 LAT/LONG: 50°00-58°00/102°45-120°00
 INSTITUTE: PNIIS, Gosstro, USSR
 NUMBER 00210

AUTHOR Lakhtina, O.V., Sukhodolskaia, L.A.
 NAME: Sketch-map of permafrost extent in Sartan stage
 PUB: Moscow, Nauka, 1981
 SCALE: 1:12 000 000
 SOURCE: "History of permafrost development in Eurasia", p.120. In the article by Lakhtina, O.V., Sukhodolskoi L.A. p.113-125
 REGION: Transbaikal
 LEGEND Permafrost spread
 LAT/LONG: 50°00-58°00/102°45-120°00
 INSTITUTE: PNIIS, Gosstro, USSR
 NUMBER 00211

AUTHOR Leshchikov, F.N.
 NAME: Geocryological map of Angaro-Ilimsk interfluve
 PUB: Moscow, Nedra, 1989
 SCALE: 1:1 500 000
 SOURCE: Geocryology of the USSR. Central Siberia / edited by E.D. Ershov, p.327
 REGION: Central Siberia
 LEGEND Expansion, thickness, the depth of seasonal freezing and thawing, permafrost terrain: frost mounds, thermokarst pits and lakes, solifluction features, polygonal wedge ice and hillocky terrain, icings, content of deposits
 LAT/LONG: 56°00-58°00/102°00-105°00
 INSTITUTE: Permafrost Institute
 NUMBER 00212

AUTHOR Leshchikov, F.N.
 NAME: Hillocky terrain expansion in Priangare
 PUB: Moscow, Nedra, 1989
 SCALE: 1:10 000 000
 SOURCE: Geocryology of the USSR. Central Siberia / edited by E.D. Ershov, p.331
 REGION: Central Siberia
 LEGEND Places with hillocky terrain, northern limit of permafrost terrain
 LAT/LONG: 51°00-59°00/92°00-110°00
 INSTITUTE: Permafrost Institute
 NUMBER 00213

AUTHOR Leshchikov, F.N.
 NAME: Map of seasonal freezing soil calculation of depths in the impact zone of the Boguchan reservoir

PUB: Moscow, Nedra, 1989
 SCALE: 1:5 000 000
 SOURCE: Geocryology of the USSR. Central Siberia / edited by E.D. Ershov, p.370
 REGION: Central Siberia
 LEGEND: The depth of freezing in natural conditions and standards for silty sands and clay, permafrost expansion, content and genesis of deposits
 LAT/LONG: 95°00-103°00/58°00-60°00
 INSTITUTE: Permafrost Institute
 NUMBER: 00214

AUTHOR: Leshchikov, F.N.
 NAME: Scheme in permafrost conditions in Muisko-Kaundinskoi and Charscoi depressions and their mountain surroundings
 PUB: Moscow, Nedra, 1989
 SCALE: 1: 5 000 000
 SOURCE: Geocryology of the USSR. Mountain countries of the southern USSR / edited by E.D. Ershov, p.93
 REGION: Transbaikial
 LEGEND: Thickness, temperature, expansion, cryogenic processes and frozen ground features
 LAT/LONG: 56°00-58°00/110°00-119°00
 INSTITUTE: Permafrost Institute
 NUMBER: 00215

AUTHOR: Leshchikov, F.N.
 NAME: Scheme of Angaro-Lensk permafrost region referring to the types of technogenic impact
 PUB: Moscow, Nedra, 1989
 SCALE: 1:10 000 000
 SOURCE: Geocryology of the USSR. Central Siberia / edited by E.D. Ershov, p.373
 REGION: Central Siberia
 LEGEND: Regions: without permafrost, pervasive and local permafrost degradation, short-term permafrost and ice lens formation, cryogenic processes activation under technogenic impact
 LAT/LONG: 51°00-62°00/95°00-110°00
 INSTITUTE: Permafrost Institute
 NUMBER: 00216

AUTHOR: Leshchikov, F.N.
 NAME: Scheme of permafrost conditions in Verhne-Angarscoi depression
 PUB: Moscow, Nedra, 1989
 SCALE: 1:1 500 000
 SOURCE: Geocryology of the USSR. Mountain countries of the south USSR / edited by E.D. Ershov, p.90
 REGION: Transbaikial
 LEGEND: Expansion, thickness, temperature, cryogenic processes and frozen ground features
 LAT/LONG: 55°00-57°00/109°00-116°00
 INSTITUTE: Permafrost Institute
 NUMBER: 00217

AUTHOR: Leshchikov, F.N., Serov, A.G.
 NAME: Sketch-map of calculated depths of seasonal freezing
 PUB: Novosibirsk, Nauka, 1983
 SCALE: 1:1 000 000
 SOURCE: Leshchikov, F.N., Shats M.M., South Central Siberia permafrost, p.148
 REGION: Central Siberia
 LEGEND: Expansion, genesis and content ground, calculation and natural depth of freezing
 LAT/LONG: 57°00-60°00/97°00-103°00

INSTITUTE: Permafrost Institute
 NUMBER 00218

AUTHOR Leshchikov, F.N.
 NAME: Sketch-map of Barguzin lowland and surrounding permafrost
 PUB: Moscow, Nedra, 1989
 SCALE: 1:1 500 000
 SOURCE: Geocryology of the USSR. Mountain contries of the south USSR / edited by E.D. Ershov, p.88
 REGION: Transbaikal
 LEGEND Expansion, thickness, temperature, depth of freezing and thawing cryogenic processes and frozen ground features, content of deposits
 LAT/LONG: 53°00-55°00/108°00-112°00
 INSTITUTE: Permafrost Institute
 NUMBER 00219

AUTHOR Leshchikov, F.N.
 NAME: Sketch-map of Srednego-Priangariya permafrost
 PUB: Novosibirsk, Nauka, 1983
 SCALE: 1:2 500 000
 SOURCE: Leshchikov, F.N., Shats, M.M. South Central Siberia permafrost, p.135
 REGION: Central Siberia
 LEGEND Expansion, thickness, temperature, frozen ground features and permafrost terrain
 LAT/LONG: 56°00-60°00/100°00-104°00
 INSTITUTE: Permafrost Institute
 NUMBER 00220

AUTHOR Leshchikov, F.N.
 NAME: The depths (m) of seasonal freezing map in Angaro-Lenski region
 PUB: Moscow, Nedra, 1989
 SCALE: 1:10 000 000
 SOURCE: Geocryology of the USSR. Central Siberia / edited by E.D. Ershov, p.326
 REGION: Central Siberia
 LEGEND Seasonal freezing of soils (5 ranges of the depth)
 LAT/LONG: 52°00-60°00/105°00-107°00
 INSTITUTE: Permafrost Institute
 NUMBER 00221

AUTHOR Likhanov, B.N.
 NAME: Sketch-map of physiographic regionalization
 PUB: Moscow, Nauka, 1972
 SCALE: 1:2 500 000
 SOURCE: Inset-map in monograph "Natural conditions of the Tazovskii oil-gas containing region industrialization"
 REGION: West Siberia
 LEGEND Flat-topped bogs and palsa peatlands
 LAT/LONG: 62°30-68°00/78°00-84°00
 INSTITUTE: Geographical Institute
 NUMBER 00222

AUTHOR Lurie, I.S.
 NAME: Sketch-map of upper level permafrost ice content
 PUB: Moscow, Moscow University Publishers, 1972
 SCALE: 1:3 000 000
 SOURCE: Permafrost Studies, XII, p.170
 REGION: West Siberia (Tazovski peninsula)

LEGEND Lithology and ice content, ground ice genesis types, average annual ground temperature, permafrost expansion, genesis types and age deposits
 LAT/LONG: 67°00-69°00/74°00-77°30
 INSTITUTE: Moscow State University
 NUMBER 00223

AUTHOR Mel'nikov, E.S.
 NAME: Landscape Regionalization scheme of northwest Siberia
 PUB: Yakutsk, Geocryological Institute Publishers, 1986
 SCALE: 1:15 000 000
 SOURCE: "Questions of geocryological mapping", p.56. In the article by Kritsuk, L.N., Mel'nikov, E.S., Moskalenko, N.G., p.53-67
 REGION: West Siberia
 LEGEND Boundaries and indexed landscape provinces, subprovinces, districts. Geocryological conditions and their changes (tabl. 1, p.62-63)
 LAT/LONG: 60°00-74°00/60°00-85°00
 INSTITUTE: VSEGINGEO
 NUMBER 00224

AUTHOR Mel'nikov, E.S., Moskalenko, N.G.
 NAME: Map of natural complexes in northwest Siberia for forecasting and planning of environment protection referring to construction
 PUB: Moscow, GUGK, 1991
 SCALE: 1:1 000 000
 REGION: West Siberia
 LEGEND Geomorphology, age of deposits, lithological composition (upper horizon), temperature, expansion, cryogenic processes, characteristics of seasonal frozen and thawing layers, permafrost terrain, tolerance for technological effects
 LAT/LONG: 50°00-70°30/60°00-90°30
 INSTITUTE: VSEGINGEO, Geological Ministry, USSR
 NUMBER 00225

AUTHOR Mel'nikov, E.S.
 NAME: Scheme of engineering-geological zoning of northern West Siberia
 PUB: Moscow, Moscow State University, 1977
 SCALE: 1:7 500 000
 SOURCE: Trofimov, V.T. "Regularities of spatial variability for engineering-geological conditions in West Siberia platform", p.147
 REGION: West Siberia
 LEGEND Engineering-geological provinces, genesis of deposits
 LAT/LONG: 62°00-74°00/60°00-90°00
 INSTITUTE: VSEGINGEO
 NUMBER 00226

AUTHOR Mel'nikov, P.I.
 NAME: Geocryological sketch-map of Yakutia
 PUB: Novosibirsk, Nauka, 1974
 SCALE: 1:25 000 000
 SOURCE: "General geocryology"/ Edited by Mel'nikov, P.I., Tolstikhin, N.I., p.99
 REGION: Central Siberia
 LEGEND Permafrost thickness, temperature, wedge ice, injected ice, injected-segregated ice, depths of seasonal thaw, continuous and discontinuous permafrost zones
 LAT/LONG: 55°00-77°00/105°00-160°00
 INSTITUTE: Permafrost Institute
 NUMBER 00227

AUTHOR Mel'nikov, P.I.
 NAME: Permafrost and landscapes map of Yakutsk Republic
 PUB: Moscow, GUGK, 1991
 SCALE: 1:2 500 000
 REGION: Yakutia
 LEGEND Expansion, temperature, ice content, cryogenic structure, thickness of seasonal frozen and thawed layers, lithogenic complexes, landscapes, permafrost and landscape
 LAT/LONG: 56°30 - 77°00/106°00 - 163°00
 INSTITUTE: Permafrost Insitute
 NUMBER 00228

AUTHOR Mimeev, V.S., Ryashin, V.A.
 NAME: Zoning of Transbaikal referring to depth of seasonal thawing and freezing
 PUB: Moscow, Nedra, 1989
 SCALE: 1:12 000 000
 SOURCE: Geocryology of the USSR. Mountain territories south USSR / edited by E.D. Ershov, p.44
 REGION: Transbaikal
 LEGEND Lithological composition, depths of freezing and thawing
 LAT/LONG: 51°00-58°00/103°00-120°00
 NUMBER 00229

AUTHOR Molodykh, I.I.
 NAME: Hillocky terrain extent and its engineering-geological parameters scheme
 PUB: Moscow, Nauka, 1988
 SCALE: 1:10 000 000
 SOURCE: Geocryology of the USSR. European territory of the USSR / edited by E.D. Ershov, p.255
 REGION: South of European part of the USSR
 LEGEND Hillocky terrain in recent and ancient landscapes, forms, parameters and relative positions in plan
 LAT/LONG: 45°00-52°00/25°00-37°00
 NUMBER 00230

AUTHOR Neizvestnov, IA.V.
 NAME: Hydrogeological basins of the Eurasia Arctic shelf in the Late Cenozoic (19-18 thousand years ago)
 PUB: Leningrad, Nedra, 1983
 SCALE: 1:150 000 000
 SOURCE: "The primary problems of paleogeography in Late Cenozoic of the Arctic", p.183
 REGION: Arctic shelf
 LEGEND Hydrogeological artesian freezing basin
 LAT/LONG: 68°00-90°00/20°00-160°00
 NUMBER 00231

AUTHOR Nekrasov, I.A.
 NAME: Baikalo-Amur main line. Geocryological map
 PUB: Moscow, GUGK, 1979
 SCALE: 1:2 500 000
 REGION: Transbaikal
 LEGEND Expansion, thickness, temperature, bedding of ground ice, type of permafrost, surface cover influence on temperature
 LAT/LONG: 52°00-58°00/105°00-141°00
 INSTITUTE: Permafrost Institute
 NUMBER 00232

AUTHOR Nekrasov, I.A., Zabolotnik, S.I.
 NAME: Geocryological map of Amur area
 PUB: Yakutsk, Permafrost Institute Publishers, 1983
 SCALE: 1:2 500 000
 SOURCE: Regional geocryological research in East Asia / edited by I.A. Nekrasov, p.112
 REGION: Far East of USSR
 LEGEND Expansion, temperature, thickness, cryogenic structure
 LAT/LONG: 48°00-56°00/120°00-135°00
 INSTITUTE: Permafrost Institute
 NUMBER 00233

AUTHOR Nekrasov, I.A., Mikova, A.I.
 NAME: Morphology and temperature at the Kolyma River headstream
 PUB: Novosibirsk, Nauka, 1975
 SCALE: 1:3 000 000
 SOURCE: Inset-map in monograph "Regional and special geocryological investigations " / Edited by V.S. Iakupov, I.V. Klimovskii
 REGION: East Siberia
 LEGEND Permafrost thickness, temperature, taliks
 LAT/LONG: 59°00-64°00/145°00-157°00
 INSTITUTE: Permafrost Institute
 NUMBER 00234

AUTHOR Nekrasov, I.A., Mikova, A.I.
 NAME: Morphology and temperature of permafrost at the Kolyma River
 PUB: Novosibirsk, Nauka, 1975
 SCALE: 1:2 000 000
 SOURCE: Regional subject and geocryological investigations, p.20
 REGION: East Siberia
 LEGEND Thickness, expansion and temperature
 LAT/LONG: 60°00-69°00/145°00-160°00
 NUMBER 00235

AUTHOR Nekrasov, I.A., Mikova, A.I.
 NAME: Permafrost zone thickness and temperature of Arkagala River system
 PUB: Novosibirsk, Nauka, 1975
 SCALE: Large
 SOURCE: "Regional and special geocryological investigations " / Edited by V.S. Iakupov, I.V. Klimovskii, p.7
 REGION: East Siberia
 LEGEND Permafrost thickness and temperature
 LAT/LONG: 62°00-70°00/145°00-150°00
 INSTITUTE: Permafrost Institute
 NUMBER 00236

AUTHOR Nekrasov, I.A., Mikova, A.I.
 NAME: Permafrost zone thickness and temperature in northern part of Seimchano- Buyundinskoi depression and its moutain surroundings
 PUB: Novosibirsk, Nauka, 1975
 SCALE: Large
 SOURCE: "Regional and special geocryological investigations " / Edited by V.S. Iakupov, I.V. Klimovskii, p.11
 REGION: East Siberia
 LAT/LONG: 62°00-64°00/150°00-155°00
 INSTITUTE: Permafrost Insitute
 NUMBER 00237

- AUTHOR Nekrasov, I.A., Zabolotnik, S.I.
NAME: Sketch-map of seasonal thaw depths zoning in Amur area
PUB: Yakutsk, Permafrost Institute Publishers, 1983
SCALE: 1:2 500 000
SOURCE: Regional geocryological reseach in East Asia / edited by I.A. Nekrasov, p.122
REGION: Far East of USSR
LEGEND Range of variations of depth of seasonal thawing sandy and clayey soils
LAT/LONG: 48°00-56°00/120°00-135°00
INSTITUTE: Permafrost Institute
NUMBER 00238
- AUTHOR Nekrasov, I.A.
NAME: Sketch-map of permafrost zone in Northern Hemisphere
PUB: Novosibirsk, Nauka, 1974
SCALE: 1:50 000 000
SOURCE: Inset-map in monograph "The general geocryology"/ Edited by Mel'nikov, P.I., Tolstikhin, N.I.
REGION: Northern Hemisphere
LEGEND Continuous, discontinuous permafrost and permafrost islands
LAT/LONG: 25°00-90°00/
INSTITUTE: Permafrost Institute
NUMBER 00239
- AUTHOR Nevecheria, V.L.
NAME: Regionalization map of southwestern Siberian South referring to rate of frost heaving processes in ground
PUB: Yakutsk, Yakutskoe knizh. izd-vo, 1966
SCALE: 1:15 000 000
SOURCE: "Reports of VIII All-Union Geocryological Conference", no.3. Regional geocryology, p.56. In the report by Nevecheria, V.L. p.53-59
REGION: West Siberia
LEGEND Districts: a) with high rate, b) with average rate, c.) with low rate of frost heaving processes in ground; boundaries of districts
LAT/LONG: 50°00-58°00/72°00-92°00
INSTITUTE: Siberian Institute of Power Engineering
NUMBER 00240
- AUTHOR Nevecheria, V.L.
NAME: The distribution of the frost-susceptible ground near the West Siberian railways
PUB: Yakutsk, Yakutskoe knizh. izd-vo, 1966
SCALE: 1:7 500 000
SOURCE: "Reports of VIII All-Union Geocryological Conference", no. 3. Regional geocryology, p.58. The report by Nevecheria, V.L. pp .53-59.
REGION: West Siberia
LEGEND Frost-susceptible ground
LAT/LONG: 52°00-58°00/78°00-90°00
INSTITUTE: Siberian Institute of Power Engineering
NUMBER 00241
- AUTHOR Nikitenko, F.A.
NAME: Scheme of engineering-geological zoning of West Siberia platform
PUB: Moscow, Moscow University Publishers, 1977
SCALE: 1:10 000 000
SOURCE: Trofimov, V.T. "Regularities of spatial variability for engineering-geological conditions in West Siberia", p.246
REGION: West Siberia

- LEGEND** Expansion, provinces of structure denudation plains
LAT/LONG: 51°00-74°00/60°00-90°00
NUMBER 00242
- AUTHOR** Novikov, V.P., Svitoch, A.A.
NAME: Geomorphological scheme of Aion Island
PUB: Moscow, Nauka, 1980
SCALE: 1:1 500 000
SOURCE: "Recent deposits and paleogeography of Pleistocene in Chukchi", p.178
REGION: Chuckchee
LEGEND Fluvio-lacustrine plain, active reworking by thermokarst processes, thermokarst depression
LAT/LONG: 69°00-70°00/167°00-170°00
INSTITUTE: Pacific Ocean Geographical Institute
NUMBER 00243
- AUTHOR** Oberman, N.G.
NAME: Hydrogeology zoning scheme of European North of the USSR
PUB: Moscow, Nauka, 1988
SCALE: 1:20 000 000
SOURCE: Geocrylogy of the USSR. European territory of the USSR / edited by E.D. Ershov, p.221
REGION: European North and Northern Urals
LEGEND Hydrogeological structures referring to permafrost expansion
LAT/LONG: 60°00-85°00/40°00-60°00
NUMBER 00244
- AUTHOR** Oberman, N.G.
NAME: Permafrost scheme of Trans-Urals
PUB: Moscow, Nauka, 1981
SCALE: 1:20 000 000
SOURCE: "History of permafrost development in Eurasia", p.62. In the article by Oberman, N.G. p.60-73
REGION: European North
LEGEND Permafrost zones: thick, two-layer and relict permafrost; thick permafrost with sheet ice and basal cryopegs; boundaries: zone, permafrost spread, thickness; sheet ice, frost mounds
LAT/LONG: 66°00-70°00/58°00-66°00
NUMBER 00245
- AUTHOR** Oberman, N.G.
NAME: Scheme of Ural permafrost
PUB: Moscow, Nedra, 1988
SCALE: 1:5 000 000
SOURCE: Geocrylogy of the USSR. European territory of the USSR / edited by E.D. Ershov, p.303
REGION: Northern and Polar Ural
LEGEND Seasonal frozen ground and permafrost zones, permafrost-temperature zonations (expansion, thickness, temperature)
LAT/LONG: 61°00-70°00/56°00-68°00
NUMBER 00246
- AUTHOR** Oberman, N.G.
NAME: Zoning scheme of Malo-Bolshezemelsky region referring to permafrost structure and thickness
PUB: Moscow, Nauka, 1988
SCALE: 1:3 500 000
SOURCE: Geocrylogy of the USSR. European territory of the USSR / edited by E.D. Ershov, p.284

REGION: European North
LEGEND: Territory with prevailing relict permafrost, unfrozen ground, places with cryopeg, depth of permafrost base isolines
LAT/LONG: 65°00-69°00/52°00-64°00
NUMBER: 00247

AUTHOR: Ospennikov, E.N., Chizhova, N.E.
NAME: Map of areal extent for geological processes and phenomena and forecasting their development for construction
PUB: Moscow, Moscow University Publishers, 1980
SCALE: 1:5 000 000
SOURCE: Ospennikov, E.N., Trush, N.I., Chizhov, A.B., Chizhova, A.B. Exogenesis geological processes and phenomena (South Yakutia), p.198-199
REGION: South Yakutia
LEGEND: Cryogenic processes and features
LAT/LONG: 57°00-69°00/123°00-128°00
INSTITUTE: Moscow State University
NUMBER: 00248

AUTHOR: Ospennikov, E.N.
NAME: Map of rock stream expansion in Aldano-Timpton interfluvial territory
PUB: Moscow, Nedra, 1989
SCALE: 1:2 000 000
SOURCE: Geocryology of the USSR. Central Siberia / edited by E.D. Ershov, p.305
REGION: Central Siberia
LEGEND: Rock streams areal extent
LAT/LONG: 55°00-60°00/122°00-130°00
NUMBER: 00249

AUTHOR: Parmuzin, S.IU.
NAME: Map of permafrost resistance referring to potential thermokarst development
PUB: Yakutsk, Geocryological Institute Publishers, 1986
SCALE: 1:10 000 000
SOURCE: "Questions of geocryological mapping", p.83. In the article by Parmuzin, S.IU. p.78-85
REGION: West Siberia
LEGEND: Places with potential thermokarst development. Southern limits : wedge ice and sheet ice spread
LAT/LONG: 60°00-74°00/64°00-88°00
INSTITUTE: Moscow State University
NUMBER: 00250

AUTHOR: Pavlova, O.P.
NAME: Geocryological sketch-map of eastern part of Baikal mountain country
PUB: Moscow, Nauka, 1975
SCALE: 1:2 500 000
SOURCE: Inset-map in monograph "Neotectonic, permafrost and ground water formation"
REGION: Transbaikalia
LEGEND: Scheme of geocryological regionalization, content and structure of permafrost, spreading, temperature, thickness, icings
LAT/LONG: 55°00-60°00/110°00-130°00
INSTITUTE: PNIIS, Gosstroiz, USSR
NUMBER: 00251

AUTHOR: Pavlova, O.P.
NAME: Scheme of icings and springs locations on alluvial fan of Verkhniy Sakukan River, April 1971

PUB: Moscow, Nauka, 1975
 SCALE: Large
 SOURCE: "Neotectonic, permafrost and ground water", p.45
 REGION: Transbaikal
 LEGEND: Icings, springs, southern limit of permafrost
 LAT/LONG: 56°00-60°00/118°00-120°00
 INSTITUTE: PNIIS, Gosstro, USSR
 NUMBER: 00252

AUTHOR: Pizhankova, A.I., Chizhov, A.B.
 NAME: Map of Southern Yakutia permafrost regionalization
 PUB: Moscow, Nedra, 1989
 SCALE: 1:10 000 000
 SOURCE: Geocryology of the USSR. Central Siberia / edited by E.D. Ershov, p.196
 REGION: Central Siberia
 LEGEND: Temperature, expansion, genesis and age of deposits
 LAT/LONG: 55°00-60°00/120°00-135°00
 INSTITUTE: Moscow State University
 NUMBER: 00253

AUTHOR: Popov, A.G., Kostiaev, A.G.
 NAME: Sketch-map of Mesopleistocene periglacial features in Asia
 PUB: Moscow, Moscow University Publishers, 1962
 SCALE: 1:40 000 000
 SOURCE: Inset-map in monograph "Questions of geographical geocryology and periglacial morphology" / edited by Popov, A.I.
 REGION: Asia
 LEGEND: Type of cryolithogenesis, area of sedimentation, southern limit of permafrost, seasonal freezing, permafrost neogenesis, ice-wedge polygons, flat-topped polygonal peatland, ground wedges, alasses, hillocky terrain
 LAT/LONG: 40°00-80°00/40°00-170°00
 INSTITUTE: Moscow State University
 NUMBER: 00254

AUTHOR: Popov, A.I., Rozenbaum, G.E., Tumel, N.V.
 NAME: Cryolithogenesis processes zoning scheme of the USSR
 PUB: Moscow, Moscow University Publishers, 1985
 SCALE: 1:60 000 000
 SOURCE: Popov, A.I., Rozenbaum, G.E., Tumel, N.V., Cryolithology, p.215
 REGION: USSR
 LEGEND: Ocean and continental sectors of polar, subpolar, boreal and subboreal zones
 LAT/LONG: 40°00-80°00/30°00-170°00
 INSTITUTE: Moscow State University
 NUMBER: 00255

AUTHOR: Popov, A.I.
 NAME: Cryolithology map (permafrost region)
 PUB: Moscow, GUGK, 1985
 SCALE: 1:4 000 000
 SOURCE: 1982
 REGION: USSR
 LEGEND: Cryogenic structure, types of cryogenesis, type of ground
 LAT/LONG: 38°00-82°00/30°00-170°00
 INSTITUTE: Moscow State University
 NUMBER: 00256

- AUTHOR** Popov, A.I.
NAME: Geocryological-geological zoning map of the USSR
PUB: Moscow, Moscow University Publishers, 1958
SCALE: 1:15 000 000
SOURCE: Inset-map in "Geographical Faculty scientific works in the International Geophysical Year", N 1. In report by Popov, A.I., p.239-264
REGION: USSR
LEGEND Numbers and regions names (geocryological information in text)
LAT/LONG: 43°00-32°00/30°00-170°00
INSTITUTE: Moscow State University
NUMBER 00257
- AUTHOR** Popov, A.I.
NAME: Geocryological sketch-map of West Siberia
PUB: Novosibirsk, Nauka, 1990
SCALE: 1:8 000 000
SOURCE: Nekrasov, and others "Geocryological investigations history in West Siberia", p.25
REGION: West Siberia
LEGEND Expansion (zones and subzones), azonal and abnormal permafrost conditions, short-term and sporadic permafrost zone
LAT/LONG: 62°00-74°00/60°00-90°00
INSTITUTE: Moscow State University
NUMBER 00258
- AUTHOR** Popov, A.I.
NAME: Global frost areas map
PUB: Moscow, Higher Institutes, 1973
SCALE: 1:200 000 000
SOURCE: Popov, A.I., Tushinskii, G.K. "Geocryology and glaciology. Teaching aid for university of the USSR", p.47
REGION: Global
LEGEND Permafrost and glacial covers, areas of systematic seasonal freezing, short-term and nonsystematic freezing
INSTITUTE: Moscow State University
NUMBER 00259
- AUTHOR** Popov, A.I.
NAME: Ground ice
PUB: Moscow, GUGK, 1971
SCALE: 1:6 000 000
SOURCE: Tyumen region atlas, vol.1, p.15
REGION: West Siberia
LEGEND Type of freezing, frozen earth material, pure ice in the ground, lithological composition
LAT/LONG: 56°00-73°00/60°00-87°00
INSTITUTE: Moscow State University
NUMBER 00260
- AUTHOR** Popov, A.I.
NAME: Map of ground ice in the northern European part of the USSR and Siberia
PUB: Moscow, Higher Institutes, 1973
SCALE: 1:10 000 000
SOURCE: Inset-map in monograph "Geocryology and glaciology. Teaching aid for universities of the USSR"
REGION: USSR
LEGEND Genesis types of ground ice in sediment deposits and original rocks (migration, wedge ice, hydrolaccoliths, ice cores in peat mounds), limits of temperature zone by Kudriavtsev,

- LAT/LONG: V.A., southern limit of permafrost
 50°00-85°00/35°00-170°00
 INSTITUTE: Moscow State University
 NUMBER 00261
- AUTHOR Popov, A.I.
 NAME: Map of permafrost on European territory of the USSR and Siberia
 PUB: Moscow, Moscow University Publishers, 1962
 SCALE: 1:2 500 000
 SOURCE: Inset-map in monograph "Questions of geographical geocryology and periglacial morphology"/ edited by A.I. Popov
 REGION: The USSR
 LEGEND Genesis types of ground ice in unconsolidated sediments and indigenous ground, wedge ice, migration ice, ice cores of pingo, frost blisters, ice in cracks of crystalline and metamorphic rocks, age of permafrost
- LAT/LONG: 40°00-80°00/40°00-170°00
 INSTITUTE: Moscow State University
 NUMBER 00262
- AUTHOR Popov, A.I.
 NAME: Permafrost terrain
 PUB: Moscow, GUGK, 1971
 SCALE: 1:6 000 000
 SOURCE: Tyumen region atlas, vol.1, p.15
 REGION: West Siberia
 LEGEND Polygonal ground (rising and descending development), frost mounds, nivation relief, solifluction features, grounds
- LAT/LONG: 56°00-73°00/60°00-87°00
 INSTITUTE: Moscow State University
 NUMBER 00263
- AUTHOR Popov, A.I.
 NAME: Sketch-map of cryological-geomorphological phenomena in the USSR
 PUB: Moscow, Academy of Sciences Publishers, 1963
 SCALE: 1:25 000 000
 SOURCE: Inset-map in proceedings "Reports on International Permafrost Conference"
 REGION: USSR
 LEGEND Ice-wedge polygons. Polygonal-wedge ice with thermokarst, flat-topped polygonal peatlands, alasses, ground wedge polygons, etc.
- LAT/LONG: 50°00-80°00/30°00-170°00
 INSTITUTE: Moscow State University
 NUMBER 00264
- AUTHOR Popov, A.I., Kostiaev, A.G.
 NAME: Sketch-map of Neopleistocene and contemporary periglacial features in Asia
 PUB: Moscow, Moscow University Publishers, 1962
 SCALE: 1:40 000 000
 SOURCE: Inset-map in monograph "Questions of geographical geocryology and periglacial morphology"/ edited by A.I. Popov
 REGION: Asia
 LEGEND Type of cryolithogenesis, southern limit of permafrost, seasonal freezing, permafrost neogenesis, ice-wedge polygons, flat-topped polygonal peatlands, ground wedges, alasses
- LAT/LONG: 40°00-82°00/30°00-170°00
 INSTITUTE: Moscow State University
 NUMBER 00265

AUTHOR Popov, A.I.
 NAME: Sketch-map of cryological geomorphological features in the USSR
 PUB: Moscow, Academy of Sciences Publishers, 1963
 SCALE: 1:2 500 000
 SOURCE: Inset-map in Collection scientific articles "Permafrost International Conference reports
 "/ edited by Tsitovich, N.A.
 REGION: The USSR
 LEGEND Type of cryolithogenesis, ice wedge polygons, ground wedges, hillocky terrain, flat-topped
 polygonal peatlands, stone sorted polygons, solifluction features, frost mounds
 LAT/LONG: 38°00-82°00/20°00-170°00
 INSTITUTE: Moscow State University
 NUMBER 00266

AUTHOR Popov, A.I.
 NAME: Sketch-map of permafrost in West Siberia
 PUB: Moscow, Moscow University Publishers, 1969
 SCALE: 1:5 000 000
 SOURCE: Permafrost Studies, IX, p.10-11
 REGION: West Siberia
 LEGEND Pure ice in the ground (content, genesis, cryogenic structure, freezing types), frozen earth
 material (content, genesis, age, cryogenic structure, freezing types, ice content)
 LAT/LONG: 64°00-72°00/60°00-83°00
 INSTITUTE: Moscow State University
 NUMBER 00267

AUTHOR Popov, A.I.
 NAME: Sketch-map of the Northern Eurasia in Mesopleistocene
 PUB: Moscow, Moscow University Publishers, 1965
 SCALE: 1:40 000 000
 SOURCE: "Ground ice", issue II / edited by A.I. Popov, p.37
 REGION: Eurasia
 LEGEND Area of maximum permafrost extent in Northern Eurasia in the Pleistocene, permafrost
 boundaries on continental areas, relict shore line
 LAT/LONG: 40°00-80°00/70°00-170°00
 INSTITUTE: Moscow State University
 NUMBER 00268

AUTHOR Ravdonekas, O.V.
 NAME: Map-scheme of permafrost base near Ust-Port settlement
 PUB: Moscow, Nedra, 1985
 SCALE: Large
 SOURCE: Baulin, V.V. "Permafrost in oil-gas containing regions of the USSR", p.99
 REGION: The Yenisei North
 LEGEND Isolines of permafrost base
 LAT/LONG: 69°30/84°30
 NUMBER 00269

AUTHOR Romanovskii, N.N., Kondratieva, K.A., Kudriavtsev, V.A.
 NAME: Chemical hydrological zoning scheme
 PUB: Moscow, Moscow University Publishers, 1970
 SCALE: 1:1 000 000
 SOURCE: Permafrost Studies, X, p.42
 REGION: Central Siberia
 LEGEND Places of fresh water and brackish water expansion and ground ice with sulphate
 magnesium-calcium content
 LAT/LONG: 68°00-70°00/140°00-144°00

- INSTITUTE: Moscow State University
NUMBER 00270
- AUTHOR Romanovskii, N.N., Kondratieva, K.A., Kudriavtsev, V.A.
NAME: Scheme of Cenozoic deposits extent in Uyandinskaya superimposed basin
PUB: Moscow, Moscow University Publishers, 1970
SCALE: 1:500 000 000
SOURCE: Permafrost Studies, X, p.82
REGION: Central Siberia
LEGEND Thermokarst lake, alasses recent beds, solifluction deposits, thermokarst lakes, genesis and age deposits
LAT/LONG: 68°00-70°00/140°00-142°00
INSTITUTE: Moscow State University
NUMBER 00271
- AUTHOR Romanovskii, N.N., Kondratieva, K.A., Kudriavtsev, V.A.
NAME: Scheme of hydrogeological zoning and icings location: Selenyakhski hydrogeological cryogenic massif, Polousnensko-Tuostachcki hydrogeological cryogenic massif
PUB: Moscow, Moscow University Publishers, 1970
SCALE: Large
SOURCE: Permafrost Studies, X, p.52
REGION: Central Siberia
LEGEND The icings can be distinguished by special feature of ground-water recharge, individual hydrolaccoliths, places with open talik, geological structure
LAT/LONG: 68°00-70°00/140°00-144°00
INSTITUTE: Moscow State University
NUMBER 00272
- AUTHOR Romanovskii, N.N., Kondratieva, K.A., Kudriavtsev, V.A.
NAME: Scheme of permafrost thickness spreading
PUB: Moscow, Moscow University Publishers, 1970
SCALE: Large
SOURCE: Permafrost Studies, X, p.36
REGION: Central Siberia
LEGEND Permafrost thickness, open taliks, boundaries of different permafrost thickness expansion, geological-tectonic structure
LAT/LONG: 68°00-70°00/140°00-144°00
INSTITUTE: Moscow State University
NUMBER 00273
- AUTHOR Rozenberg, L.I.
NAME: Scheme of Kuraiskoi lowland permafrost
PUB: Moscow, Nauka, 1989
SCALE: Large
SOURCE: Geocryology of the USSR. Mountain countries of the southern USSR / edited by E.D. Ershov, p.233
REGION: Southern Siberia (Altai)
LEGEND Expansion, genesis of deposits
LAT/LONG: 49°30-50°30/87°00-89°00
NUMBER 00274
- AUTHOR Rozenberg, L.I. (using maps by N.I. Trush and K.A. Kondrateva)
NAME: Scheme of Kuznetsk highland permafrost spreading
PUB: Moscow, Nauka, 1989
SCALE: Large
SOURCE: Geocryology of the USSR. Mountain countries of the southern USSR / edited by E.D.

REGION: Ershov, p.252
 LEGEND: Southern Siberia
 LAT/LONG: Expansion, temperature, thickness
 NUMBER: 50°00-55°00/85°00-90°00
 00275

AUTHOR: Rozenberg, L.I. (using map by N.I. Trush)
 NAME: Scheme of permafrost of Tuva Central area
 PUB: Moscow, Nauka, 1989
 SCALE: 1:7 500 000
 SOURCE: Geocryology of the USSR. Mountain countries of the southern USSR / edited by E.D. Ershov, p.280

REGION: Southern Siberia (Tuva)
 LEGEND: Expansion, temperature, thickness
 LAT/LONG: 50°00-55°00/90°00-100°00
 NUMBER: 00276

AUTHOR: Rozenberg, L.I. (using maps by M.M. Shats and N.I. Trush)
 NAME: Scheme of permafrost in Altai
 PUB: Moscow, Nauka, 1989
 SCALE: 1:3 000 000
 SOURCE: Geocryology of the USSR. Mountain countries of the southern USSR / edited by E.D. Ershov, p.231

REGION: Southern Siberia (Altai)
 LEGEND: Expansion, temperature, thickness
 LAT/LONG: 48°00-53°30/81°00-89°00
 NUMBER: 00277

AUTHOR: Rozenberg, L.I.
 NAME: Scheme of seasonal frozen ground and permafrost in Enisey ridge
 PUB: Moscow, Nauka, 1989
 SCALE: 1:3 000 000
 SOURCE: Geocryology of the USSR. Mountain countries of the southern USSR / edited by E.D. Ershov, P.290

REGION: Southern Siberia
 LEGEND: Expansion, temperature, thickness
 LAT/LONG: 55°00-62°00/93°00-100°00
 NUMBER: 00278

AUTHOR: Rozenberg, L.I. (using map by N.I. Trush)
 NAME: Scheme of seasonal frozen ground and permafrost extent in East Sayan
 PUB: Moscow, Nauka, 1989
 SCALE: 1:7 500 000
 SOURCE: Geocryology of the USSR. Mountain countries of the southern USSR / edited by E.D. Ershov, p.267

REGION: Southern Siberia (Easten Sayan)
 LEGEND: Expansion, temperature, thickness, dynamic short-term permafrost, seasonal freezing of ground
 LAT/LONG: 51°00-54°00/92°00-99°00
 NUMBER: 00279

AUTHOR: Rozenberg, L.I. (using map by N.I.Trush)
 NAME: Scheme of West Sayan permafrost extent
 PUB: Moscow, Nauka, 1989
 SCALE: 1:7 500 000
 SOURCE: Geocryology of the USSR. Mountain countries of the southern USSR / edited by E.D.

- Ershov, p.261
 REGION: Southern Siberia (Western Sayan)
 LEGEND: Expansion, temperature, thickness
 LAT/LONG: 53°00-54°00/91°00-93°00
 NUMBER: 00280
- AUTHOR: Sendek, S.V., Barbashinov, G.L.
 NAME: Okhotsk Sea shore geocryology
 PUB: Novosibirsk, Nauka, 1975
 SCALE: 1:1 000 000
 SOURCE: "Regional and special geocryological investigations " / Edited by V.S. Iakupov, I.V. Klimovskii, p.19
 REGION: East Siberia
 LEGEND: Permafrost expansion, thickness, taliks
 LAT/LONG: 59°00-61°00/151°00-155°00
 NUMBER: 00281
- AUTHOR: Sergeev, E.M.
 NAME: Engineering- geological map of West Siberian platform
 PUB: Moscow, GUGK, 1972
 SCALE: 1:1 500 000
 REGION: West Siberia
 LEGEND: Formations, genesis complexes, lithological composition of upper complex, permafrost conditions, expansion, temperature, thickness, ice content, cryogenic processes, permafrost terrain
 LAT/LONG: 50°00-70°30/60°00-90°30
 INSTITUTE: Moscow State University, Geological Ministry, the Second Hydrogeological Department
 NUMBER: 00282
- AUTHOR: Severskii, I.V., Severskii, E.V.
 NAME: Scheme of deep seasonal freezing of soil area in Central Asia mountains and south-eastern Kazakhstan
 PUB: Yakutsk, Geocryological Institute Publishers, 1986
 SCALE: 1:7 500 000
 SOURCE: "Questions of geocryological mapping", p.36. In the article by Severskii, I.V., Severskii, E.V. p.29-38
 REGION: Central Asia, southeastern Kazakhstan
 LEGEND: Deep seasonal freezing of soil areas and their boundaries
 LAT/LONG: 37°00-44°00/70°00-83°00
 NUMBER: 00283
- AUTHOR: Shats, M.M., Leshchikov, F.N.
 NAME: Map of permafrost in the southern part of Central Siberia
 PUB: Novosibirsk, Nauka, 1983
 SCALE: 1:2 500 000
 SOURCE: Insert-map in monograph "South Central Siberia permafrost" by Leshchikov, F.N., Shats, M.M.
 REGION: Central Siberia
 LEGEND: Seasonal frozen and permafrost grounds
 LAT/LONG: 53°00-61°00/90°00-110°00
 INSTITUTE: Permafrost Institute
 NUMBER: 00284
- AUTHOR: Shats, M.M.
 NAME: Sketch-map of permafrost expansion in the Krasnoyarsk region south
 PUB: Novosibirsk, Nauka, 1980

SCALE: 1:5 000 000
 SOURCE: Permafrost studies in development of regions of the USSR, p.122
 REGION: Central Siberia
 LEGEND Seasonal frozen ground, permafrost (islands, discontinuous, sporadic)
 LAT/LONG: 52°00-56°00/89°00-95°00
 INSTITUTE: Permafrost Institute
 NUMBER 00285

AUTHOR Sheko, A.I., Fotiev, S.M.
 NAME: Map of permafrost zoning of Baikalo-Amur main line zone
 PUB: Moscow, GUGK, 1988
 SCALE: 1:3 000 000
 SOURCE: Atlas of geological maps of Baikalo-Amur main line zone
 REGION: Baikal, Transbaikal, Amur regions
 LEGEND Expansion, temperature, thickness, depth of seasonal freezing and thaw, cryogenic processes
 LAT/LONG: 52°00-58°00/105°00-141°00
 INSTITUTE: Tsentrogeologiya
 NUMBER 00286

AUTHOR Shevchenko, V.K.
 NAME: Map of permafrost zoning of Transbaikal
 PUB: Moscow, Nedra, 1989
 SCALE: 1:10 000 000
 SOURCE: Geocryology of the USSR. Mountain territories south USSR / edited by E.D. Ershov, p.63
 REGION: Transbaikal
 LEGEND Expansion, temperature, thickness
 LAT/LONG: 51°00-80°00/103°00-120°00
 NUMBER 00287

AUTHOR Sheveleva, N.S.
 NAME: Geocryological sketch-map of the Yenisei-North region
 PUB: Yakutsk, Yakutskoe knizh. izd-vo, 1966
 SCALE: 1:1 500 000
 SOURCE: Inset-map in monograph "Reports of VIII All-Union Geocryological Conference", issue3.Regional geocryology
 REGION: The Yenisei-North
 LEGEND Type of freezing, ice content, cryogenic structure, thaw settlement, permafrost thickness, temperature, spreading; age, lithological composition and genesis of deposits; regionalization
 LAT/LONG: 67°20-70°00/84°00-92°00
 INSTITUTE: PNIIS, Gosstroj, USSR
 NUMBER 00288

AUTHOR Sheveleva, N.S.
 NAME: Sketch-map of permafrost temperature in Central Siberia
 PUB: Moscow, Nedra, 1975
 SCALE: 1:20 000 000
 SOURCE: Fotiev, S. M., Danilova, N.S., Sheveleva, N.S. Permafrost conditions of Central Siberia, p.110
 REGION: Central Siberia
 LEGEND Temperature isolines of grounds on base of layer with annual temperature fluctuations
 LAT/LONG: 52°00-80°00/78°00-138°00
 INSTITUTE: PNIIS, Gosstroj USSR
 NUMBER 00289

AUTHOR Sheveleva, N.S.
 NAME: Temperature of permafrost, sketch-map of the Yenisei-North
 PUB: Yakutsk, Yakutskoe knizh. izd-vo, 1966
 SCALE: 1:5 000 000
 SOURCE: "Reports of VIII All-Union Geocryological Conference", no. 3. Regional geocryology, p.78.
 In the report by Sheveleva, N.S. p.71-80
 REGION: The Yenisei-North
 LEGEND Isotherms
 LAT/LONG: 67°20'-70°00'/84°00'-92°00'
 INSTITUTE: PNIIS, Gosstroj, USSR
 NUMBER 00290

AUTHOR Shpolianskaia, N.A.
 NAME: Permafrost (spreading, temperature)
 PUB: Moscow, GUGK, 1971
 SCALE: 1:4 000 000
 SOURCE: Tyumen region atlas, publish 1, p.14
 REGION: West Siberia
 LEGEND Expansion, temperature
 LAT/LONG: 56°00'-73°00'/60°00'-87°00'
 INSTITUTE: Moscow State University
 NUMBER 00291

AUTHOR Shpolianskaia, N.A.
 NAME: Permafrost in Kazan epoch
 PUB: Moscow, Moscow University Publishers, 1981
 SCALE: 1:25 000 000
 SOURCE: Shpolianskaia, N.V. "West Siberian permafrost region and the tendency for development",
 p.108
 REGION: West Siberia
 LEGEND Active freezing areas, initial permafrost degradation areas, southern limit of permafrost,
 permafrost thickness and temperature
 LAT/LONG: 55°00'-72°00'/60°00'-85°00'
 INSTITUTE: Moscow State University
 NUMBER 00292

AUTHOR Shpolianskaia, N.A.
 NAME: Permafrost in maximum glacial epoch
 PUB: Moscow, Moscow University Publishers, 1981
 SCALE: 1:25 000 000
 SOURCE: Shpolianskaia, N.V. "West Siberian permafrost region and the tendency for development",
 p.104
 REGION: West Siberia
 LEGEND Active freezing areas in the divides, areas of glaciation, areas of freezing in river valleys,
 permafrost thickness and temperature, southern limit of permafrost
 LAT/LONG: 50°00'-72°00'/50°00'-85°00'
 INSTITUTE: Moscow State University
 NUMBER 00293

AUTHOR Shpolianskaia, N.A.
 NAME: Permafrost in thermic maximum stage
 PUB: Moscow, Moscow University Publishers, 1981
 SCALE: 1:25 000 000
 SOURCE: Shpolianskaia, N.V. "West Siberian permafrost region and the tendency for development",
 p.109
 REGION: West Siberia

LEGEND Disequilibrium, equilibrium, relict permafrost, southern limit of permafrost, northern forest boundary, permafrost thickness and temperature on the depth of annual temperature fluctuations, the depth of relict permafrost table and base
LAT/LONG: 55°00-72°00/60°00-85°00
INSTITUTE: Moscow State University
NUMBER 00294

AUTHOR Shpolianskaia, N.A.
NAME: Permafrost in Zyriansk glaciation epoch
PUB: Moscow, Moscow University Publishers, 1981
SCALE: 1:25 000 000
SOURCE: Shpolianskaia, N.V. "West Siberian permafrost region and the tendency for development", p.108
REGION: West Siberia
LEGEND Glaciation areas, active freezing places on the divides and river valleys, disequilibrium permafrost, southern limit of permafrost, permafrost thickness and temperature
LAT/LONG: 55°00-72°00/60°00-85°00
INSTITUTE: Moscow State University
NUMBER 00295

AUTHOR Shpolianskaia, N.A.
NAME: Permafrost map
PUB: Yakutsk, Yakutskoe knizh. izd-vo, 1966
SCALE: 1:7 500 000
SOURCE: Inset-map in monograph "Reports of VIII All-Union Geocryological Conference", issue 3. Regional geocryology
REGION: Transbaikal
LEGEND Temperature, thickness, spreading
LAT/LONG: 49°00-53°00/103°00-118°00
INSTITUTE: Moscow State University
NUMBER 00296

AUTHOR Shpolianskaia, N.A.
NAME: Quaternary changes in permafrost
PUB: Moscow, GUGK, 1971
SCALE: 1:16 000 000
SOURCE: Tyumen region atlas, p.15
REGION: West Siberia
LEGEND Temperature, thickness, southern limit of permafrost
LAT/LONG: 56°00-73°00/60°00-87°00
INSTITUTE: Moscow State University
NUMBER 00297

AUTHOR Shpolianskaia, N.A.
NAME: Seasonal freezing and thawing
PUB: Moscow, GUGK, 1971
SCALE: 1:4 000 000
SOURCE: Tyumen region atlas, p.15
REGION: West Siberia
LEGEND Depths and dates freezing and thawing
LAT/LONG: 56°00-73°00/60°00-87°00
INSTITUTE: Moscow State University
NUMBER 00298

AUTHOR Shpolianskaia, N.A.
NAME: Seasonal freezing and thawing dynamics

PUB: Moscow, GUGK, 1971
 SCALE: 1:12 000 000
 SOURCE: Tyumen region atlas, publish 1, p.15
 REGION: West Siberia
 LEGEND: Depths and dates of freezing and thawing
 LAT/LONG: 56°00-73°00/60°00-87°00
 INSTITUTE: Moscow State University
 NUMBER: 00299

AUTHOR: Shpolianskaia, N.A.
 NAME: Seasonal freezing and thawing of soils in West Siberia
 PUB: Moscow, Moscow University Publishers, 1981
 SCALE: 1:12 000 000
 SOURCE: Shpolianskaia, N.A. "West Siberian permafrost region and the tendency for development", p.40
 REGION: West Siberia
 LEGEND: Depths of seasonal thaw and freezing in permafrost and unfrozen ground and organic soils
 LAT/LONG: 55°00-74°00/65°00-85°00
 INSTITUTE: Moscow State University
 NUMBER: 00300

AUTHOR: Shpolianskaia, N.A.
 NAME: Sketch-map of permafrost in West Siberia
 PUB: Moscow, Nauka, 1976
 SCALE: 1:12 000 000
 SOURCE: "Questions of global cryology", p.51
 REGION: West Siberia
 LEGEND: Areas of untouched and closed contemporary and relict permafrost, deep bedding of relict permafrost; zones: permafrost expansion, permafrost steady-state conditions, permafrost degradation, forecast to the end of century
 LAT/LONG: 60°00-74°00/55°00-90°00
 INSTITUTE: Moscow State University
 NUMBER: 00301

AUTHOR: Shpolianskaia, N.A.
 NAME: West Siberian North Regionalization scheme referring to the Ob River run off shortening influence under the permafrost
 PUB: Moscow, Moscow University Publishers, 1981
 SCALE: 1:10 000 000
 SOURCE: Shpolianskaia, N.V. "West Siberian permafrost region and the tendency for development", p.146
 REGION: West Siberia
 LEGEND: Southern limit of permafrost in ground and peatlands, southern limit of relict permafrost
 LAT/LONG: 60°00-72°00/60°00-80°00
 INSTITUTE: Moscow State University
 NUMBER: 00302

AUTHOR: Shpolianskaia, N.A.
 NAME: West Siberian permafrost spreading, temperature and thickness
 PUB: Moscow, Moscow University Publishers, 1981
 SCALE: 1:12 000 000
 SOURCE: Shpolianskaia, N.A. "West Siberian permafrost region and the tendency for development", p.8
 REGION: West Siberia
 LEGEND: Areas, zones, types referring to permafrost expansion, temperature and thickness,

single-layer and two-layer permafrost, southern limit of permafrost
 LAT/LONG: 58°00-74°00/65°00-85°00
 INSTITUTE: Moscow State University
 NUMBER 00303

AUTHOR Slavianskii, A.M., Shpakov, O.N.
 NAME: Okhotsk Sea shore geocryology
 PUB: Novosibirsk, Nauka, 1975
 SCALE: 1:1 000 000
 SOURCE: "Regional and special geocryological investigations"/ Edited by V.S. Iakupov, I.V. Klimovskii, p.19
 REGION: East Siberia
 LEGEND Permafrost expansion, thickness, taliks
 LAT/LONG: 59°00-61°00/150°00-151°30
 NUMBER 00304

AUTHOR Solomatin, V. I.
 NAME: Scheme of geocryological zonality
 PUB: Novosibirsk, Nauka, 1986
 SCALE: 1:50 000 000
 SOURCE: Solomatin, V.I. "Petrogenesis of ground ice", p.203
 REGION: USSR
 LEGEND Boundaries: geocryological zone, natural zonality
 LAT/LONG: 50°00-80°00/40°00-170°00
 INSTITUTE: Moscow State University
 NUMBER 00305

AUTHOR Solomatin, V.I.
 NAME: Cryolithozone landscape structure
 PUB: Moscow, Moscow University Publishers, 1992
 SCALE: 1:80 000 000
 SOURCE: "Geocology of the North (introduction in geocryocology)"/ Edited by V.I. Solomatin, p.19
 REGION: Northern Hemisphere
 LEGEND Boundary of cryolithozone, types and subtypes: subarctic, boreal, subboreal
 LAT/LONG: 40°00-90°00/
 INSTITUTE: Moscow State University
 NUMBER 00306

AUTHOR Solomatin, V.I.
 NAME: Scheme of regionalization of cryolithozone in Eurasia
 PUB: Novosibirsk, Nauka, 1986
 SCALE: 1:50 000 000
 SOURCE: Solomatin, V.I. "Petrogenesis of ground ice", p.206
 REGION: The USSR
 LEGEND Boundaries: geocryological zone, morphostucture, natural zonality
 LAT/LONG: 50°00-80°00/40°00-170°00
 INSTITUTE: Moscow State University
 NUMBER 00307

AUTHOR Solovev, P.A.
 NAME: Alas valley is in a primitive state of development on Lena River
 PUB: Moscow, Academy of Sciences Publishers, 1963
 SCALE: 1:10 000
 SOURCE: "Conditions and permafrost development features in Siberia and North-East", p.84. In the article by Solovev, P.A. p.80-90
 REGION: Central Yakutia

- LEGEND** Polygon wedge ice thawing, bed of thermokarst depression, limit of icy complex, polygonal system in depression, slope brow of alas valley
LAT/LONG: 60°00-68°00/120°00-140°00
INSTITUTE: Geocryological Institute
NUMBER 00308
- AUTHOR** Solovev, P.A.
NAME: Lithological composition of ground at a depth of 2 m in the Yakutsk district
PUB: Yakutsk, Yakutsk Republican Printing Plant, 1958
SCALE: Large
SOURCE: "Transactions of the North-East Permafrost Institute department", no. 1, p.182. In the article by Solovev, P.A. p.179-191
REGION: Yakutia
LEGEND Average annual temperature geoisotherms at the depth of 10 m, lithological composition
LAT/LONG: 62°30/129°30
INSTITUTE: Permafrost Institute
NUMBER 00309
- AUTHOR** Solovev, P.A.
NAME: Mature alas valley of Kokhara River
PUB: Moscow, Academy of Sciences Publishers, 1963
SCALE: 1:100 000 000
SOURCE: "Conditions and permafrost development features in Siberia and North-East", p.83. In the article by Solovev, P.A. p.80-90
REGION: Central Yakutia
LEGEND Thermokarst is in a primitive state of development, alas depression, lake depression on alas valley floor
LAT/LONG: 60°00-68°00/120°00-140°00
INSTITUTE: Geocryological Institute
NUMBER 00310
- AUTHOR** Solovev, P.A.
NAME: The geoisotherms arrangement dependent on the age of Yakutsk town construction
PUB: Yakutsk, Yakutsk Republican Printing Plant, 1958
SCALE: Large
SOURCE: "Transaction of the North-East Permafrost Institute department", no. 1, p.184. In the article by Solovev, P.A. p.179-191
REGION: Yakutia
LEGEND Geoisotherms, geomorphological levels, places of building
LAT/LONG: 62°30/129°30
INSTITUTE: Permafrost Institute
NUMBER 00311
- AUTHOR** Solovev, V.A., Telepnev, E.V.
NAME: Sketch-map of arctic shelf cryolithozone
PUB: Leningrad: Nedra, 1983
SCALE: 1:10 000 000
SOURCE: "The primary problems of paleogeography of the Late Cenozoic in the Arctic", p.188
REGION: Arctic shelf
LEGEND Predominant unfrozen cryolithozone with cryopegs. Relict permafrost islands in area of extent of above zero temperature ground on ocean bed, cryopegs, predominant continuous permafrost, island permafrost
LAT/LONG: 68°00-84°00/20°00-162°00
NUMBER 00312
- AUTHOR** Soloviev, V.A.

NAME: Forecast map of Laptev and East-Siberian seas Arctic shelf permafrost
 PUB: Yakutsk, Permafrost Institute Publishers, 1979
 SCALE: 1:10 000 000
 SOURCE: Permafrost zone of Arctic shelf, p.36
 REGION: Arctic shelf of the USSR
 LEGEND: Expansion and thickness of relict and contemporary permafrost
 LAT/LONG: 70°00-80°00/108°00-170°00
 INSTITUTE: Permafrost Institute
 NUMBER: 00313

AUTHOR: Soloviev, V.A.
 NAME: Ground with cryopegs expansion and permafrost in Arctic Ocean and Barents sea shelf zones
 PUB: Moscow, Nauka, 1988
 SCALE: 1:10 000 000
 SOURCE: Geocryology of the USSR. European territory of the USSR / edited by E.D. Ershov, p.260
 REGION: Barents Sea
 LEGEND: Ground thickness with seasonal and perennial cryopegs, permafrost and seasonal frozen ground with cryopegs, sporadic and continuous permafrost thickness less than 50 m, relict ground ice
 LAT/LONG: 64°00-82°00/30°00-75°00
 NUMBER: 00314

AUTHOR: Soloviev, V.A., Michaluk, U.N.
 NAME: Laptev Sea and East-Siberian Sea shelf in late Cenozoic
 PUB: Leningrad, PGO "Sevmorgeologia", 1982
 SCALE: 1:20 000 000
 SOURCE: Hydrogeological and permafrost conditions of Arctic and Continental shelf of Eurasia, p.25
 REGION: East Arctic shelf of the USSR
 LEGEND: Areas of the longest period of Pleistocene freezing, areas of short duration of freezing before Holocene, boundaries of the sea advance
 LAT/LONG: 70°00-80°00/108°00-180°00
 NUMBER: 00315

AUTHOR: Soloviev, V.A., Ginsburg, G.D.
 NAME: Map of research areas location in Enisei-Khatanga and Leno-Anabar depression
 PUB: Moscow, Nedra, 1989
 SCALE: 1:20 000 000
 SOURCE: Geocryology of the USSR. Central Siberia / edited by E.D. Ershov, p.156
 REGION: Central Siberia
 LEGEND: Permafrost zone thickness (established and assumed)
 LAT/LONG: 70°00-75°00/85°00-130°00
 INSTITUTE: Permafrost Institute
 NUMBER: 00316

AUTHOR: Spesivtsev, V.I., Leshchikov, F.N.
 NAME: Sketch-map of permafrost conditions in upper part of Kulenga River
 PUB: Novosibirsk, Nauka, 1983
 SCALE: Large
 SOURCE: Leshchikov, F.N., Shats, M.M. South Central Siberia permafrost, p.140
 REGION: Central Siberia
 LEGEND: Expansion, temperature, thickness, depth of seasonal freeze and thaw, cryogenic processes and permafrost terrain
 LAT/LONG: 56°00-59°00/105°00-107°00
 INSTITUTE: Permafrost Institute

NUMBER 00317
 AUTHOR NAME: Sukhodolskii, S.E.
 European North-East zoning scheme referring to conditions of paragenesis association between ground water and permafrost
 PUB: Moscow, Nauka, 1982
 SCALE: 1:7 500 000
 SOURCE: Sukhodolski S.E. "Underground water and permafrost paragenesis", p.135
 REGION: North-East European part of the USSR
 LEGEND Condition types: platform, submontane. Condition grades: northern, southern. Condition facies : water-chemical delay, water-thermal delay, water-thermal intensive. Index of hydrogeocryological district
 LAT/LONG: 65°00-67°00/56°00-58°00
 INSTITUTE: PNIIS, Gosstroj USSR
 NUMBER 00318
 AUTHOR NAME: Sukhodolskii, S.E., Kaznacheeva, I.A., Kondratieva, K.A., Oberman, N.G., Soloviev, V.A.
 European North of the USSR zoning scheme referring to discontinuous permafrost
 PUB: Moscow, Nauka, 1988
 SCALE: 1:10 000 000
 SOURCE: Geocryology of the USSR. European territory of the USSR / edited by E.D. Ershov, p.229
 REGION: European North of the USSR
 LEGEND Expansion
 LAT/LONG: 60°00-85°00/40°00-60°00
 INSTITUTE: PNIIS, Gosstroj USSR
 NUMBER 00319
 AUTHOR NAME: Sukhodolskii, S.E.
 European North of the USSR zoning referring to permafrost age
 PUB: Moscow, Nauka, 1988
 SCALE: 1:10 000 000
 SOURCE: Geocryology of the USSR. European territory of the USSR / edited by E.D. Ershov, p.212
 REGION: North of European part of the USSR
 LEGEND Age and genesis of permafrost, permafrost expansion, vertical structure
 LAT/LONG: 64°00-76°00/32°00-72°00
 INSTITUTE: PNIIS, Gosstroj USSR
 NUMBER 00320
 AUTHOR NAME: Sukhodolskii, S.E.
 Geocryological scheme of Yanei-ti-vis head stream
 PUB: Moscow, Nauka, 1982
 SCALE: Large
 SOURCE: Sukhodolski S.E. "Underground water and permafrost paragenesis", p.117
 REGION: North-East European part of the USSR
 LEGEND Permafrost thickness, open talik, thermokarst lakes, surface springs
 LAT/LONG: 65°00-67°00/56°00-58°00
 INSTITUTE: PNIIS, Gosstroj USSR
 NUMBER 00321
 AUTHOR NAME: Sukhodolskii, S.E.
 Geocryological zones European North- East
 PUB: Moscow, Nedra, 1985
 SCALE: 1:5 000 000
 SOURCE: Baulin, V.V. "Permafrost in oil-gas containing regions of the USSR", p.86
 REGION: European North-East
 LEGEND Permafrost zones, of continuous and discontinuous permafrost

LAT/LONG: 66°00-70°00/52°00-66°00
INSTITUTE: PNIIS, Gosstro, USSR
NUMBER 00322

AUTHOR Sukhodolskii, S.E.
NAME: Map of native taliks in North-East European part of the USSR
PUB: Moscow, Nauka, 1982
SCALE: 1:5 000 000
SOURCE: Sukhodolski S.E. "Underground water and permafrost paragenesis", p.110
REGION: North-East European part of the USSR
LEGEND Under interblock depression taliks, under drainage line taliks, under temporary stream valley taliks: closed, open, saline water interapermafrost taliks, intrapermafrost taliks in Cenozoic deposits, permafrost zones boundaries

LAT/LONG: 48°00-66°00/64°00-70°00
INSTITUTE: PNIIS, Gosstro USSR
NUMBER 00323

AUTHOR Sukhodolskii, S.E., Kondratieva, K.A.
NAME: Map of permafrost expansion and average annual temperature in North-East European part of the USSR
PUB: Moscow, Nauka, 1982
SCALE: 1:5 000 000
SOURCE: Sukhodolski S.E. "Underground water and permafrost paragenesis", p.30
REGION: North-East European part of the USSR
LEGEND Permafrost average annual temperature and taliks, permafrost expansion, permafrost zones and zonal boundaries, index geocryological zones

LAT/LONG: 48°00-66°00/64°00-70°00
INSTITUTE: Moscow State University, PNIIS, Gosstro USSR
NUMBER 00324

AUTHOR Sukhodolskii, S.E.
NAME: North-East European part of the USSR permafrost thickness map
PUB: Moscow, Nauka, 1982
SCALE: 1:5 000 000
SOURCE: Sukhodolski S.E. "Underground water and permafrost paragenesis", p.42
REGION: North-East European part of the USSR
LEGEND Permafrost thickness (14 ranges), unfrozen ground, depth of relict permafrost table
LAT/LONG: 48°00-66°00/64°00-70°00
INSTITUTE: PNIIS, Gosstro USSR
NUMBER 00325

AUTHOR Sukhodolskii, S.E., Parmuzin, S.IU., Streletskaya, I.D.
NAME: Sketch-map of geocryological zoning of Bovanenkovo tectonic structure territory
PUB: Moscow, Stroiizdat, 1984
SCALE: Large
SOURCE: Geocryological conditions and their change forecast in the primary development regions of the North, p.69
REGION: West Siberia
LEGEND Geocryological regions and subregions (types of freezing, genesis, age, ice content, lithological composition, temperature, depth of thaw, depth of sheet ice table, cryogenic processes and frozen ground features)
LAT/LONG: 70°00-71°00/68°00-69°00
INSTITUTE: PNIIS, Gosstro USSR
NUMBER 00326

AUTHOR Sumgin, M.I.

NAME: Southern limit of permafrost in West Siberia plain on evidence from some scientists
PUB: Novosibirsk, Nauka, 1990
SCALE: 1:8 000 000
SOURCE: Nekrasov, I.A., et. al. "Geocryological investigations history in West Siberia", p.12
REGION: West Siberia
LEGEND Boundaries by Vild (1882), Yachevskomu (1989), Shtini and Mushketov (1925), Shostakovich (1928), Berg (1931), Evdokimov-Rokotovskii (1931), Sumginu (1931)
LAT/LONG: 62°00-74°00/60°00-90°00
NUMBER 00327

AUTHOR Svitoch, A.A., Khorev, V.S.
NAME: Geomorphological scheme of Anadyr lowland southern shore
PUB: Moscow, Nauka, 1980
SCALE: Large
SOURCE: "Recent deposits and paleogeography of Pleistocene in the Chukchi Sea", p.47
REGION: Chukchi Sea
LEGEND Solifluction slopes
LAT/LONG: 63°00-65°00/175°00-180°00
INSTITUTE: Pacific Ocean Geographical Institute
NUMBER 00328

AUTHOR Svitoch, A.A.
NAME: Geomorphological scheme of the shore (northern Konergino settlement)
PUB: Moscow, Nauka, 1980
SCALE: Large
SOURCE: "Recent deposits and paleogeography of Pleistocene in the Chukchi Sea", p.9
REGION: Chukchi Sea
LEGEND Places of active wedge ice thawing
LAT/LONG: 65°00-68°00/180°00-175°00
NUMBER 00329

AUTHOR Tolstikhin, O.N.
NAME: Hydrogeological scheme
PUB: Novosibirsk, Nauka, 1974
SCALE: 1:12 000 000
SOURCE: Inset-map in monograph "Icing and underground water of the North-East USSR" by Tolstikhin, O.N.
REGION: North-East of USSR
LEGEND Cryogenic basins, hydrogeological massifs and structures
LAT/LONG: 60°00-75°00/125°00-170°00
INSTITUTE: Permafrost Institute
NUMBER 00330

AUTHOR Tomirdiario, S.V.
NAME: Contemporary extent of Neopleistocene icy complex (edoma) and Holocene lake-alas plains (East-Siberian lowland)
PUB: Moscow, Nauka, 1980
SCALE: 1:20 000 000
SOURCE: Tomirdiario, S.V. "Ice complex of East-Siberia in Neopleistocene and Holocene", p.26
REGION: North-East
LEGEND Holocene thermokarst lake-alas plain, large icy complexes and relict Neopleistocene plain
LAT/LONG: 62°00-74°00/130°00-170°00
NUMBER 00331

AUTHOR Tomirdiario, S.V., in base of map by Velichko A.A.
NAME: General permafrost-glacial condition in Neopleistocene

PUB: Moscow, Nedra, 1978
SCALE: 1:400 000 000
SOURCE: Tomirdiario, S.V. "Natural processes and industrialization of territory in permafrost zone", p.8
REGION: Global
LEGEND Limits of permafrost in Neopleistocene and at present, limits of glaciations, sea ice, Neopleistocene sea ice cover
NUMBER 00332

AUTHOR Tomirdiario, S.V., Shilo, N.A.
NAME: Geocryological-geomorphological sketch-map of North-East of the USSR
PUB: Moscow, Nedra, 1978
SCALE: 1:15 000 000
SOURCE: Tomirdiario, S.V. "Natural processes and industrialization of territory of permafrost zone", p.4
REGION: North-East
LEGEND Polygonal wedge ice (active, degradation, stable), degradation island permafrost, solifluction lobes, larger icings, annual temperature of ground isotherms in upper part of permafrost
LAT/LONG: 55°00-80°00/125°00-170°00
NUMBER 00333

AUTHOR Tomirdiario, S.V.
NAME: Permafrost-landscape sketch-map of Anadyr lowland
PUB: Magadan, Knizh. izd-vo, 1972
SCALE: 1:5 000 000
SOURCE: Tomirdiario, S.V. "Perennial frost and industrialization of the mountain countries and lowlands, the Magadan area and Yakutia", p.64
REGION: North-East
LEGEND The landscapes: alas, undulating-morainic, kettlehole-lake on different geomorphological levels
LAT/LONG: 62°00-67°00/168°-178°00
NUMBER 00334

AUTHOR Tomirdiario, S.V.
NAME: Scheme of glacier, loesses and loessial-glacial plains in the Northern Hemisphere
PUB: Magadan, Knizh. izd-vo, 1972
SCALE: 1:200 000 000
SOURCE: Tomirdiario, S.V. "Perennial frost and industrialization in mountain countries and lowlands, the Magadan area and Yakutia", p.6
REGION: Northern Hemisphere
LEGEND Contemporary limit of permafrost in Eurasia and North America, loessial-glacial plains, loessial surface formations, glaciers, assumed zone of marine glaciation
NUMBER 00335

AUTHOR Tomirdiario, S.V.
NAME: Sketch-map of frost-glacial condition in the Northern Hemisphere
PUB: Magadan, Academy of Sciences Publishers, 1971
SCALE: 1:300 000 000
SOURCE: "Periglacial processes", no.38, p.139
REGION: Global
LEGEND Contemporary limit of permafrost zone
NUMBER 00336

AUTHOR Trofimov, V.T., Badu, I.U.B, Vasilchuk, I.U.K.
NAME: A change of macro-ice content in deposits of Mamont and Olenii peninsula (active growth

of polygon wedge ice)
PUB: Moscow, Moscow University Publishers, 1986
SCALE: 1:1 000 000
SOURCE: "Engineering-geological conditions of Gydan Peninsula", p.184
REGION: West Siberia, Gydan Peninsula
LEGEND: Situation in different periods: on boundary of Pleistocene and Holocene, at present time, macro-ice content shown in scale map
LAT/LONG: 71°00-72°30/76°30-78°30
INSTITUTE: Moscow State University
NUMBER: 00337

AUTHOR: Trofimov, V.T., Badu, IU.B., Vasilchuk, IU.K.
NAME: Permafrost extent and contemporary trend of polygonal ice wedge development in territory of Gydan Peninsula
PUB: Moscow, Moscow University Publishers, 1986
SCALE: 1:3 000 000
SOURCE: "Engineering-geological conditions of Gydan Peninsula", p.183
REGION: West Siberia, Gydan Peninsula
LEGEND: Zones of combined extent of relict and contemporary wedge ice, subzones of contemporary syngeneses and epigeneses polygonal wedge ice, degradation of wedge ice as a result of activity of erosion and temperature rise
LAT/LONG: 67°30-74°00/73°00-84°00
INSTITUTE: Moscow State University
NUMBER: 00338

AUTHOR: Trofimov, V.T., Gruzlov, I.V., Tyrtikov, A.P.
NAME: Permafrost thickness (m)
PUB: Moscow, Nauka, 1972
SCALE: 1:2 500 000
SOURCE: "Natural conditions of the Tazovskii oil-gas containing region industrialization", p.86
REGION: West Siberia
LEGEND: Permafrost thickness (5 ranges)
LAT/LONG: 63°00-68°00/78°00-87°00
INSTITUTE: Moscow State University
NUMBER: 00339

AUTHOR: Trofimov, V.T., Gruzlov, I.V., Tyrtikov A.P.
NAME: Scheme of distribution of average annual temperature of perennial frozen and unfrozen ground
PUB: Moscow, Nauka, 1972
SCALE: 1:2 500 000
SOURCE: "Natural conditions of the Tazovskii oil-gas containing region industrialization", p.89
REGION: West Siberia
LEGEND: Average annual temperature of ground (4 ranges)
LAT/LONG: 63°00-68°00/78°00-87°00
INSTITUTE: Moscow State University
NUMBER: 00340

AUTHOR: Trofimov, V.T., Badu, IU.B., Vasilchuk, IU.K.
NAME: Scheme of engineering-geological regionalization of Gydan Peninsula
PUB: Moscow, Moscow University Publishers, 1986
SCALE: 1:3 000 000
SOURCE: "Engineering-geological conditions of Gydan Peninsula", p.197
REGION: West Siberia, Gydan Peninsula
LEGEND: Engineering-geological subzones, areas (description of permafrost conditions in monograph)

LAT/LONG: 67°30'-74°00'/73°00'-84°00'
INSTITUTE: Moscow State University
NUMBER 00341

AUTHOR Trofimov, V.T.
NAME: Scheme of engineering-geological regionalization of Taz drainage basin
PUB: Moscow, Nauka, 1972
SCALE: 1:2 500 000
SOURCE: "Natural conditions of the Tazovskii oil-gas containing region industrialization", p.36
REGION: West Siberia
LEGEND Engineering-geological regions and their indexes (explanations in text p.35-37: permafrost spread)

LAT/LONG: 63°00'-68°00'/78°00'-87°00'
INSTITUTE: Moscow State University
NUMBER 00342

AUTHOR Trofimov, V.T.
NAME: Scheme of engineering-geological zoning of West Siberia platform
PUB: Moscow, Moscow University Publishers, 1977
SCALE: 1:10 000 000
SOURCE: Trofimov, V.T. "Regularities of spatial variability for engineering-geological conditions in West Siberia platform", p.248
REGION: West Siberia
LEGEND Expansion, geological-genesis complexes
LAT/LONG: 51°00'-74°00'/60°00'-90°00'
INSTITUTE: Moscow State University
NUMBER 00343

AUTHOR Trofimov, V.T., Datsko, L.T.
NAME: Scheme of ground type in West Siberian platform
PUB: Moscow, Moscow University Publishers, 1977
SCALE: 1:10 000 000
SOURCE: Trofimov, V.T. "Regularities of spatial variability for engineering-geological conditions in West Siberia platform", p.184
REGION: West Siberia
LEGEND Unfrozen ground, permafrost, permafrost in the lowermost layer of ground and unfrozen ground in upper part of section, ice content
LAT/LONG: 62°00'-74°00'/60°00'-90°00'
INSTITUTE: Moscow State University
NUMBER 00344

AUTHOR Trofimov, V.T., Gruzdrov, I.V., Tyrtikov, A.P.
NAME: Scheme of permafrost and unfrozen ground on different geomorphological levels
PUB: Moscow, Nauka, 1972
SCALE: 1:2 500 000
SOURCE: "Natural conditions of the Tazovskii oil-gas containing region industrialization", p.82
REGION: West Siberia
LEGEND Permafrost extent, geomorphological levels, lines of schematic geocryological sections
LAT/LONG: 63°00'-68°00'/78°00'-87°00'
INSTITUTE: Moscow State University
NUMBER 00345

AUTHOR Trofimov, V.T., Firsov, N.G.
NAME: West- Siberian platform zoning scheme referring to contemporary exogenesis geological processes and phenomena
PUB: Moscow, Moscow University Publishers, 1986

SCALE: 1:7 500 000
 SOURCE: West-Siberian platform exodynamic (spatial-temporal regularities) / edited by V.T. Trofimov, p.179
 REGION: West Siberia
 LEGEND: Contemporary processes and phenomena of extent zone in permafrost region (continuous and discontinuous permafrost expansion)
 LAT/LONG: 50°00-72°00/60°00-92°00
 INSTITUTE: Moscow State University
 NUMBER: 00346

AUTHOR NAME: Trush, N.I., Chizhova, N.E.
 NAME: Geocryological map of Aldano-Timpton interfluve
 PUB: Moscow, Moscow University Publishers, 1980
 SCALE: 1:2 000 000
 SOURCE: Ospennikov, E.N., Trush, N.I., Chizhov, A.B., Chizhova, N.I. Exogenesis geological processes and phenomena, p.24
 REGION: South Yakutia
 LEGEND: Expansion, temperature, thickness, ice content, cryogenic structure, cryogenic phenomena, lithological composition of grounds
 LAT/LONG: 57°00-69°00/123°00-128°00
 INSTITUTE: Moscow State University
 NUMBER: 00347

AUTHOR NAME: Tsitovich, N.A.
 NAME: Map of the depth of seasonal freezing of loam, loamy sand and clay in the USSR by normatives and specifications
 PUB: Moscow, Academy of Sciences Publishers, 1958
 SCALE: 1:30 000 000
 SOURCE: Tsitovich, N.A. "Foundations in permafrost", p.14
 REGION: European part of the USSR and West Siberia
 LEGEND: Isoline of normative depths of freezing loam, loamy sand and clay
 LAT/LONG: 40°00-68°00/24°00-84°00
 NUMBER: 00348

AUTHOR NAME: Tumel, V.F.
 NAME: Map of permafrost in the USSR
 PUB: Moscow, Academy of the USSR, 1946
 SCALE: 1:8 000 000
 SOURCE: Permafrost Studies, 1(1), p.5-11
 REGION: USSR
 LEGEND: Expansion, northern boundaries of area with temperature at the depth of 10 m above a -5, -3, -1°C, thickness
 LAT/LONG: 40°00-82°00/30°00-170°00
 NUMBER: 00349

AUTHOR NAME: Vasilchuk, I.U.K.
 NAME: A correlation between oxygen-18 content in contemporary wedge ice (formed in Eurasian permafrost over the last 100 years) and summary of winter temperature
 PUB: Moscow, Nauka, 1991
 SCALE: 1:35 000 000
 SOURCE: Vasilchuk, I.U.K. "Late Quaternary syngenes permafrost strata of the North of Eurasia: structure, oxygen-isotope content and conditions of formation", abstract thesis, p.16
 REGION: North of Eurasia
 LEGEND: Southern limit of contemporary active growth wedge ice
 LAT/LONG: 60°00-80°00/40°00-170°00
 INSTITUTE: PNIIS, Gosstrois USSR

NUMBER 00350

AUTHOR Vasilchuk, I.U.K., Badu, I.U.B., Trofimov, V.T.
NAME: Map of cryogenic formations and ice content of upper part (10 m) permafrost
PUB: Moscow, Moscow University Publishers, 1986
SCALE: 1:3 000 000
SOURCE: "Engineering-geological conditions of Gydan Peninsula", p.73
REGION: West Siberia, Gydan Peninsula
LEGEND Cryogenic formations, ice content, lithological composition, macro-ice content at the expense of polygonal-wedge ice, buried ice, injective-segregated ice, pingos, hydrolaccolith

LAT/LONG: 67°30'-74°00'/73°00'-84°00'
INSTITUTE: Moscow State University
NUMBER 00351

AUTHOR Vasilchuk, I.U.K., Kudriashov, V.G., Trofimov, V.T.
NAME: Schemes of forecast changes in engineering-geological conditions for Northwest Siberia (Yamal-Gydan Province) under the effect of natural climatic trend
PUB: Moscow, Moscow University Publishers, 1986
SCALE: 1:7 500 000
SOURCE: West-Siberian platform exodynamic (spatial-temporal regularities) / edited by V.T. Trofimov, p.218
REGION: Northwest Siberia
LEGEND Towards the end of the 19th century and by 2020 AD location of places with ground temperature above -3°, -5°, wedge ice growth-southern limit, frost mounds formation-northern limit

LAT/LONG: 65°00'-85°00'/68°00'-72°00'
INSTITUTE: Moscow State University
NUMBER 00352

AUTHOR Vasiliev, I.S.
NAME: Map of thawing ground in East Yakutia
PUB: Novosibirsk, Nauka, 1982
SCALE: 1:2 500 000
SOURCE: Inset-map in monograph by Vasiliev, I.S. "Conformity to natural laws seasonal freezing and thawing of ground in East Yakutia"
REGION: East Yakutia
LEGEND Depth of seasonal thaw, lithological composition of ground, genesis complexes, vegetation
LAT/LONG: 59°00'-76°00'/126°00'-162°00'
INSTITUTE: Permafrost Institute
NUMBER 00353

AUTHOR Vasiliev, I.S.
NAME: Sketch-map of ground thawing season duration
PUB: Novosibirsk, Nauka, 1982
SCALE: 1:7 500 000
SOURCE: Inset-map in monograph by Vasiliev, I.S. "Conformity to seasonal freezing and thawing of ground in East Yakutia"
REGION: East Yakutia
LEGEND Duration seasonal freezing (days), average long term dates of onset and end of thawing season
LAT/LONG: 59°00'-76°00'/126°00'-162°00'
INSTITUTE: Permafrost Institute
NUMBER 00354

AUTHOR Velichko, A.A.
NAME: Assumed limit of permafrost in North America in the third stage of Pleistocene

PUB: Moscow, Nauka, 1973
 SCALE: 1:80 000 000
 SOURCE: "Natural process in Pleistocene", p.174
 REGION: North America
 LEGEND: Assumed permafrost zones, contemporary permafrost zone
 LAT/LONG: 30°00-80°00/00°00-160°00
 INSTITUTE: Geographical Institute
 NUMBER: 00355

AUTHOR: Velichko, A.A.
 NAME: Cryogenic area of Northern Hemisphere in third stage of Pleistocene
 PUB: Moscow, Nauka, 1973
 SCALE: 1:150 000 000
 SOURCE: "Natural process in Pleistocene", p.114
 REGION: Northern Hemisphere
 LEGEND: Sea glaciation, permafrost, ice cover glaciation
 LAT/LONG: 40°00-90°00/
 INSTITUTE: Geographical Institute
 NUMBER: 00356

AUTHOR: Velichko, A.A.
 NAME: Degradation of sea ice and permafrost in passing from the third stage of Pleistocene to present time
 PUB: Moscow, Nauka, 1973
 SCALE: 1:200 000 000
 SOURCE: Inset-map in the monograph by Velichko, A.A. "Natural process in Pleistocene"
 REGION: Global
 LEGEND: Areas of degradation: permafrost, sea ice, limits of cover ice, permafrost, pack ice, sea ice, relict permafrost
 INSTITUTE: Geographical Institute
 NUMBER: 00357

AUTHOR: Velichko, A.A.
 NAME: Marine cover ice spread, permafrost and regression of the ocean in upper Pleistocene (the third stage)
 PUB: Moscow, Nauka, 1973
 SCALE: 1:200 000 000
 SOURCE: "Natural process in Pleistocene", p.173
 REGION: Global
 LEGEND: Limits: permafrost, sea ice, pack ice, area of ocean regression (northern permafrost zone)
 INSTITUTE: Geographical Institute
 NUMBER: 00358

AUTHOR: Velichko, A.A.
 NAME: Relict permafrost degradation in Eurasia
 PUB: Moscow, Nauka, 1973
 SCALE: 1:80 000 000
 SOURCE: "Natural process in Pleistocene", p.130
 REGION: USSR and Eastern Europe
 LEGEND: Zone of degradation on the land and on the sea, contemporary permafrost, contemporary sea ice spread, limit of permafrost
 LAT/LONG: 40°00-80°00/00°00-170°00
 INSTITUTE: Geographical Institute
 NUMBER: 00359

AUTHOR: Velichko, A.A.

NAME: Scheme of relict permafrost ("periglacial") morphosculpture on Russian plain
PUB: Moscow, Nauka, 1973
SCALE: 1:5 000 000
SOURCE: "Natural process in Pleistocene", p.111
REGION: European part of the USSR
LEGEND: Southern limits of contemporary and Pleistocene permafrost zones, cryogenic relief in permafrost zone: relict cryogenic and thermokarst relief, contemporary polygon relief, hillocky terrain, thermokarst, relict thermokarst
LAT/LONG: 40°00'-70°00'/30°00'-60°00'
INSTITUTE: Geographical Institute
NUMBER: 00360

AUTHOR: Vologodski, G.P., Gerakov, N.N., Doronina, M.A., Zaikova, Z.I., Zarubin, N.E., Palshin, G.B., Portnova, V.P.
NAME: Engineering-geological conditions of Transbaikal region
PUB: Moscow-Irkutsk, 1967
SCALE: 1:3 500 000
SOURCE: Atlas Zabaikale, p.16-17
REGION: Transbaikal
LEGEND: Cryogenic processes
LAT/LONG: 58°30'-49°50'/100°00'-140°00'
INSTITUTE: Siberia and Far East Institute Academy of Science of the USSR
NUMBER: 00361

AUTHOR: Vtiurin, B.I.
NAME: Sketch-map for total evident ice content of permafrost in the USSR
PUB: Moscow, Nauka, 1975
SCALE: 1:50 000 000
SOURCE: Vtiurin, B.I. Ground ice in the USSR, p.172
REGION: USSR
LEGEND: Ice content in percent to a depths of - 5m, 10m, 20m, 30m, 50m, districts of ice sheets expansion, frost mounds, wedge ice, glaciers
LAT/LONG: 38°00'-80°00'/30°00'-190°00'
INSTITUTE: Pacific Ocean Institute of Geography
NUMBER: 00362

AUTHOR: Vtiurin, B.I.
NAME: Sketch-map for total evident ground ice reserves in the USSR
PUB: Moscow, Nauka, 1975
SCALE: 1:50 000 000
SOURCE: Vtiurin, B.I. Ground ice in the USSR, p.174
REGION: USSR
LEGEND: Reserves of various types of ice
LAT/LONG: 38°00'-80°00'/30°00'-190°00'
INSTITUTE: Pacific Ocean Institute of Geography
NUMBER: 00363

AUTHOR: Vtiurin, B.I.
NAME: Sketch-map for ice wedges extent in USSR
PUB: Moscow, Nauka, 1975
SCALE: 1:50 000 000
SOURCE: Vtiurin, B.I. Ground ice in the USSR, p.142
REGION: USSR
LEGEND: Wedge ice expansion in percent of area, boundaries: contemporary permafrost, relict permafrost and expansion of wedge ice
LAT/LONG: 38°00'-82°00'/30°00'-170°00'

INSTITUTE: Pacific Ocean Institute of Geography
NUMBER 00364

AUTHOR Vtiurin, B.I., Shum'skii, P.A.
NAME: Sketch-map of injective ice extent
PUB: Moscow, Academy of Sciences Publishers, 1963
SCALE: 1:40 000 000
SOURCE: "Permafrost International Conference reports " / edited by Tsitovich, N.A., p.45
REGION: The USSR
LEGEND Southern limit of permafrost, frost mounds (seasonal, perennial) on lake taliks and springs, relict sheet ice and injective ice (existence and assumed)
LAT/LONG: 38°00-82°00/20°00-170°00
INSTITUTE: Geographical Institute
NUMBER 00365

AUTHOR Vtiurin, B.I.
NAME: Sketch-map of permafrost with various structures
PUB: Moscow, Nauka, 1975
SCALE: 1:50 000 000
SOURCE: Vtiurin, B.I. Ground ice in the USSR, p.166
REGION: USSR
LEGEND Permafrost (epigenesis and syngeneses types), polygenesis, two horizon permafrost
LAT/LONG: 38°00-82°00/30°00-170°00
INSTITUTE: Pacific Ocean Institute of Geography
NUMBER 00366

AUTHOR Vtiurin, B.I.
NAME: Sketch-map of massive ice extent in the USSR
PUB: Moscow, Nauka, 1975
SCALE: 1:25 000 000
SOURCE: Vtiurin, B.I. Ground ice in the USSR, p.136
REGION: USSR
LEGEND Frost mounds, sheet ice, wedge ice, cave ice, buried ice, unknown origin ice, glaciers
LAT/LONG: 38°00-82°00/30°00-190°00
INSTITUTE: Pacific Ocean Institute of Geography
NUMBER 00367

AUTHOR Vtiurin, B.I.
NAME: Sketch-map the extent of permafrost with various structures
PUB: Moscow, Nedra, 1975
SCALE: 1:50 000 000
SOURCE: Vtiurin, B.I. Ground ice in the USSR, p.102
REGION: USSR
LEGEND Syngeneses and epigenesis permafrost
LAT/LONG: 38°00-82°00/30°00-170°00
INSTITUTE: Pacific Ocean Institute of Geography
NUMBER 00368

AUTHOR Vtiurin, B.I.
NAME: Sketch-maps of area and volumetric macro-ice content at the expanse of wedge ice in the USSR
PUB: Moscow, Nauka, 1975
SCALE: 1:50 000 000
SOURCE: Vtiurin, B.I. Ground ice in the USSR, p.179-180
REGION: USSR
LEGEND Ice area in percent, volumetric macro-ice content on the different depths (to 30m)

LAT/LONG: 38°00-80°00/30°00-190°00
INSTITUTE: Pacific Ocean Institute of Geography
NUMBER 00369

AUTHOR Vtiurina, E.A.
NAME: General map of seasonal frozen ground
PUB: Moscow, Nauka, 1984
SCALE: 1:30 000 000
SOURCE: Inset-map in monograph "Seasonal frozen ground" by Vtiurina, E.A.
REGION: USSR
LEGEND Types of seasonal frozen ground of northern, intermediate and southern zones
LAT/LONG: 38°00-80°00/20°00-170°00
INSTITUTE: PNIIS, Gosstroj USSR
NUMBER 00370

AUTHOR Vtiurina, E.A.
NAME: General map of seasonal cryogenic ground
PUB: Moscow, Nauka, 1984
SCALE: 1:30 000 000
SOURCE: Inset-map in monograph "Seasonal frozen ground" by Vtiurina, E.A.
REGION: USSR
LEGEND Northern, transitional, intermediate, southern seasonal frozen ground and their combinations
LAT/LONG: 38°00-80°00/20°00-170°00
INSTITUTE: PNIIS, Gosstroj USSR
NUMBER 00371

AUTHOR Vtiurina, E.A.
NAME: Sketch-map of cryogenic structure of seasonal frozen ground
PUB: Moscow, Nauka, 1984
SCALE: 1:30 000 000
SOURCE: Inset-map in monograph "Seasonal frozen ground" by Vtiurina, E.A.
REGION: USSR
LEGEND Type of seasonal frozen ground, cryogenic structure, present wedge ice, icing, lake, glacial, river buried ice
LAT/LONG: 38°00-80°00/20°00-170°00
INSTITUTE: PNIIS, Gosstroj USSR
NUMBER 00372

AUTHOR Vtiurina, E.A.
NAME: Sketch-map of landscapes of Bolshezemel tundra
PUB: Moscow, TSINIS, 1971
SCALE: Large
SOURCE: "Geocryological and hydrogeological research for engineering investigations", Transactions of PNIIS, vol. 8, p.100
REGION: European North
LEGEND Peatlands, spot medallions, frost mounds
LAT/LONG: 65°00-70°00/45°00-65°00
INSTITUTE: PNIIS, Gosstroj USSR
NUMBER 00373

AUTHOR Zhigarev, L.A.
NAME: Map-scheme of permafrost of East-Siberian Sea and Chukchi Sea
PUB: Moscow, Nauka, 1981
SCALE: 1:12 000 000
SOURCE: Inset-map in Collection articles "History of permafrost development in Eurasia". In the

REGION: article by Zhigarev, L.A. p.181-191
 LEGEND: North-East USSR
 Permafrost and seasonal supercooled ground, relict permafrost, contemporary and relict permafrost extent in sea shore zone, isolines of the depth of zero annual amplitude, limit of permafrost complexes
 LAT/LONG: 65°00-76°00/135°00-170°00
 INSTITUTE: Moscow State University
 NUMBER: 00374

AUTHOR: Zhigarev, L.A., Parmuzin, O.IU.
 NAME: Thermal stability of permafrost for cold and warm climatic rhythms for different snow and vegetation cover thicknesses (18 maps)
 PUB: Moscow, Moscow University Publishers, 1992
 SCALE: 1:20 000 000
 SOURCE: "Geocology of the North (Introduction in geocryoeology)"/ Edited by V.I. Solomatin, p.190-192
 REGION: The North of West Siberia
 LEGEND: Thermal stability of permafrost for cold and warm climatic rhythms for different snow and vegetation cover thicknesses
 LAT/LONG: 66°00-73°00/60°00-84°00
 INSTITUTE: Moscow State University
 NUMBER: 00375

RUSSIAN PERMAFROST MAP INVENTORY INDEX BY SCALE

Numbers given refer to citations given in the inventory, p.33-113

SCALE: 1:1 000 000 — 1:1 500 00

00281, 00304, 00270, 00337, 00118, 00225,
00218, 00152, 00107, 00151, 00121, 00282,
00153, 00120, 00243, 00202, 00212, 00217,
00032, 00219, 00288

SCALE: 1:2 000 000 — 1:2 500 00

00249, 00235, 00048, 00183, 00182, 00347,
00070, 00071, 00228, 00232, 00222, 00220,
00033, 00233, 00062, 00266, 00061, 00060,
00284, 00262, 00339, 00340, 00125, 00238,
00201, 00199, 00342, 00353, 00005, 00172,
00013, 00167, 00012, 00251, 00345

SCALE: 1:3 000 000 — 1:4 000 000

00277, 00011, 00278, 00234, 00008, 00007,
00006, 00223, 00286, 00341, 00351, 00338,
00027, 00047, 00001, 00361, 00190, 00247,
00256, 00291, 00298, 00124, 00131

SCALE: 1:5 000 000 — 1:6 000 000

00168, 00165, 00188, 00084, 00360, 00014,
00019, 00334, 00214, 00215, 00325, 00132,
00078, 00324, 00323, 00322, 00025, 00246,
00248, 00053, 00290, 00142, 00285, 00267,
00049, 00148, 00149, 00263, 00260

SCALE: 1:7 500 000

00147, 00099, 00117, 00100, 00154, 00155,
00113, 00116, 00112, 00133, 00130, 00129,
00111, 00115, 00110, 00109, 00114, 00081,
00354, 00352, 00241, 00009, 00346, 00010,
00016, 00017, 00018, 00226, 00318, 00296,
00283, 00280, 00279, 00276, 00072

SCALE: 1:8 000 000 — 1:10 000 000

00258, 00327, 00349, 00063, 00164, 00186,
00185, 00095, 00036, 00035, 00055, 00090,
00287, 00058, 00250, 00313, 00319, 00015,
00086, 00044, 00253, 00312, 00314, 00065,
00209, 00242, 00068, 00021, 00213, 00343,
00320, 00216, 00134, 00230, 00057, 00302,
00221, 00074, 00308, 00344, 00261

SCALE: 1:12 000 000 — 1:16 000 000

00303, 00301, 00098, 00229, 00080, 00300,
00211, 00210, 00066, 00208, 00299, 00330,
00374, 00097, 00096, 00189, 00067, 00135,
00224, 00206, 00240, 00205, 00200, 00003,
00333, 00175, 00257, 00045, 00046, 00150,
00297

SCALE: 1:20 000 000 — 1:25 000 000

00069, 00163, 00004, 00244, 00191, 00073,
00075, 00159, 00160, 00145, 00144, 00162,
00245, 00157, 00170, 00204, 00056, 00375,
00194, 00091, 00039, 00176, 00171, 00181,
00093, 00173, 00174, 00315, 00143, 00316,
00103, 00331, 00289, 00141, 00101, 00136,
00102, 00177, 00295, 00294, 00043, 00293,
00292, 00042, 00264, 00034, 00041, 00038,
00037, 00092, 00367, 00227, 00028

SCALE: 1:30 000 000 — 1:40 000 000

00348, 00370, 00372, 00023, 00371, 00050,
00350, 00187, 00178, 00166, 00156, 00094,
00179, 00184, 00082, 00161, 00002, 00030,
00024, 00265, 00254, 00031, 00180, 00268,
00195, 00193, 00365

SCALE: 1:50 000 000 — 1:80 000 000

00087, 00369, 00128, 00108, 00020, 00106,
00368, 00026, 00366, 00307, 00362, 00305,
00088, 00192, 00364, 00363, 00239, 00059,
00255, 00355, 00306, 00198, 00359

SCALE: 1:100 000 000 — 1:500 000 000

00089, 00310, 00231, 00356, 00259, 00197,
00357, 00123, 00358, 00335, 00146, 00336,
00332, 00105, 00271

SCALE: LARGE (No scale given)

00051, 00373, 00119, 00122, 00126, 00127,
00104, 00137, 00138, 00139, 00140, 00158,
00169, 00196, 00203, 00085, 00207, 00083,
00269, 00329, 00328, 00022, 00326, 00321,
00317, 00311, 00079, 00309, 00029, 00272,
00077, 00076, 00064, 00236, 00237, 00054,
00273, 00252, 00052, 00040, 00275, 00274

APPENDIX 6

LIST OF INSTITUTIONS IN RUSSIA AND THE COMMONWEALTH OF INDEPENDENT STATES INVOLVED IN STUDIES OF PERMAFROST AND SEASONALLY-FROZEN GROUND

Compiled by M. Liebman with additional entries extracted from "Spravochnik Organizatsii Nauchno-Tekhnicheskoi Sfery", All-Russian Scientific and Technical Information Center, Moscow, 1994, translated by R.G. Barry. We have included telephone numbers where available. However they may be incomplete. Please contact an international operator before placing a call.

INSTITUTE: All-Russia Federal Institute for Planning and Research
CITY: 319922 Kharkhov
ADDRESS: Prosp. Pravdy, 10
COUNTRY: UKRAINE
PHONE: 94-0687

INSTITUTE: All-Russia Gold and Rare Metal Research Institute (VNII-1)
CITY: 685000, Magadan
ADDRESS: Gagarina str., 12
COUNTRY: RUSSIA
PHONE: 2-5739

INSTITUTE: All-Russia Institute for Natural Protection and Reservations (VNIIpriroda)
MINISTRY: Ministry for Environmental Protection and Natural Resources, RF
CONTACT: Peshkov, Andrei Sergeevich
CITY: 113628, Moscow
ADDRESS: Znamenskoe-Sadky
COUNTRY: RUSSIA
PHONE: (095)423-0311

INSTITUTE: All-Russia Oil Geological Prospecting Research Institute (VNIGRY)
CITY: 191104, St. Petersburg
ADDRESS: Litejny Prospekt, 39,
COUNTRY: RUSSIA

INSTITUTE: All-Russia Research and Planning-Surveying Institute for Pipeline Hydrotransport
(VNIPI Gidrotzuboprovod)
CITY: 125422, Moscow
ADDRESS: Solomennoi Storozhki Prosp., 12
COUNTRY: RUSSIA
PHONE: (095) 257-9852

INSTITUTE: All-Russia Research Institute for Hydrogeology and Engineering Geology
(VSEGINGEO)
CITY: 142452, Moscow Region
ADDRESS: Noginsky District, Pos Zeleny
COUNTRY: RUSSIA
PHONE: (095)521-1101

INSTITUTE: All-Russia Research Institute for Hydrometeorological Information
 World Data Center-B (VNIIGMI-MCD)
CITY: 249020, Moscow Region
ADDRESS: Obninsk, Koroleva str., 6
COUNTRY: RUSSIA
PHONE: (08439)255-2194

INSTITUTE: All-Russia Research Institute for Survey Methods and Techniques (VITR)
CITY: 199106, St. Petersburg
ADDRESS: Vesel'naya, 6
COUNTRY: RUSSIA
PHONE: (812) 217-5049

INSTITUTE: Arctic and Antarctic Research Institute
CONTACT: Ivanov, Vladimir
CITY: 199226, St. Petersburg
ADDRESS: Bering str., 38
COUNTRY: RUSSIA
PHONE: (812) 352-26-88

INSTITUTE: Bashkirskiy State Scientific and Planning Institute for the Oil Industry
 (Bashnipineft')
CITY: 450077, Ufa
ADDRESS: Lenina str., 86
COUNTRY: RUSSIA

INSTITUTE: Byelorussian Research Institute for Geological Survey (Byel NIERI)
CITY: 22060, Minsk GSP
ADDRESS: Staroborisovskiy Prop. 14
COUNTRY: BYELORUS

INSTITUTE: Center for International Projects (CMP)
MINISTRY: Ministry for Environmental Protection and Natural Resources, RF
CONTACT: Tikhanov, Sergei Eduardovich
CITY: 107078, Moscow
ADDRESS: Kedrov str., 8-1
COUNTRY: RUSSIA
PHONE: (095)207-4929

INSTITUTE: Chita Branch of the All-Russia Research, Planning and Construction Institute of
 Mining and Metallurgy of Non-ferrous Metals
CITY: 672078 Chita
ADDRESS: Lermontova str., 2
COUNTRY: RUSSIA

INSTITUTE: Chita Institute of Natural Resources
MINISTRY: Siberian Branch of the Russian Academy of Sciences
CONTACT: Malchikova, Irina
CITY: 672014, Chita
ADDRESS: Nedorezov str., 16, ChIPR
COUNTRY: RUSSIA
PHONE: (30222)6-2233

INSTITUTE: Chita Polytechnical Institute
MINISTRY: Ministry for Higher Education, RF
CITY: 672076, Chita
ADDRESS: Kalinin str., 17
COUNTRY: RUSSIA
PHONE: (30222)3-1825

INSTITUTE: Coal Institute
MINISTRY: Siberian Branch of the Russian Academy of Sciences
CITY: 650610, Kemerovo
ADDRESS: Rukavishnikova str., 21
COUNTRY: RUSSIA
PHONE: 28-1329

INSTITUTE: DalGiprotrans
MINISTRY: Ministry for Transport Construction
CONTACT: Solodovnikov, Boris Ivanovich
CITY: 680628, Khabarovsk
ADDRESS: Sheronov str., 56
COUNTRY: RUSSIA
PHONE: (4210)38-4860

INSTITUTE: Deep-Sea Oceanological Institute (TOL DVO RAN)
MINISTRY: Far Eastern Branch of the Russian Academy of Sciences
CITY: 690032 Vladivostok
ADDRESS: Radio str., 7
COUNTRY: RUSSIA
PHONE: 9-6500

INSTITUTE: Dokuchaev Soil Institute
MINISTRY: Academy for Agriculture
CONTACT: Naumov, Evgeny Mikhailovich
COUNTRY: RUSSIA
PHONE: (095) 230-8302

INSTITUTE: Far Eastern Research Institute for Planning-Surveying and Technological
Construction (Dal'NIIS)
CITY: 690106, Vladivostok
ADDRESS: Borodinskaya, 14
COUNTRY: RUSSIA
PHONE: 6-0077

INSTITUTE: Federal Center for Geoecological Systems (FCGS)
MINISTRY: Ministry for Environmental Protection and Natural Resources, RF
CONTACT: Gavrilov, Vsevolod Valerianovich
CITY: 101000 Moscow
ADDRESS: Central Post Office, P.O.Box 785
COUNTRY: RUSSIA
PHONE: (095)254-4933

INSTITUTE: Federal Scientific Center for Problems of Ecological Risk (Ecorisk)
MINISTRY: Ministry for Environmental Protection and Natural Resources, RF
CONTACT: Kharchenko, Sergei Grigorievich
CITY: 103064, Moscow
ADDRESS: Obukh str., 10, INTEKO
COUNTRY: RUSSIA

INSTITUTE: Fundamentproekt
MINISTRY: Ministry for Construction RF
CONTACT: Minkin, Mark Abramovich
CITY: 125843, Moscow
ADDRESS: Volokolamskoe Shosse, 1
COUNTRY: RUSSIA
PHONE: (095)158-9538

INSTITUTE: Geofond
MINISTRY: All-Russia Geological Archive
CITY: 123806, Moscow
ADDRESS: 3 Magistralnaya str, 38
COUNTRY: RUSSIA

INSTITUTE: Geographical Institute of the Kazakh Republic (IGAN RK)
CITY: 480100, Alma-Ata
ADDRESS: Pushkina str., 99
COUNTRY: KAZAKHSTAN
PHONE: 61-8869

INSTITUTE: Geographical Institute of the Russian Academy of Sciences (IGAN RF)
CITY: 109017 Moscow
ADDRESS: Staromonetny Per., 29
COUNTRY: RUSSIA
PHONE: (095) 128-2854

INSTITUTE: Geological Geophysical Institute of the Uzbek Academy of Sciences (IGIGAN Ruz)
CITY: 017, Tashkent
ADDRESS: Suleimanovoy str., 33
COUNTRY: UZBEKISTAN
PHONE: (3912) 33-7741

INSTITUTE: Geological Institute of the Estonia Academy of Sciences
CITY: 200101, Tallinn
ADDRESS: Estonia Boul., 7
COUNTRY: ESTONIA
PHONE: 60-5120

INSTITUTE: Gidroproekt
CITY: 125813, Moscow
ADDRESS: GSP, Volokolamskoe Shosse, 2
COUNTRY: RUSSIA

INSTITUTE: Gidroproekt. Bratsk Department
CITY: 665705, Irkutsk Region, Bratsk
ADDRESS: Gydrostroitelei str., 57
COUNTRY: RUSSIA

INSTITUTE: Gidroproekt. East-Siberian Department
CONTACT: Pohekutov, K.E.
CITY: 660607, Moscow
ADDRESS: Prospekt Mira, 94
COUNTRY: RUSSIA

INSTITUTE: Gidropetsgeologia
 CITY: 123436, Moscow
 ADDRESS: Marshal Rybalko str., 4
 COUNTRY: RUSSIA

INSTITUTE: Giprogor
 CITY: 125124, Moscow
 ADDRESS: 1st Yamskoe Pole, 15
 COUNTRY: RUSSIA

INSTITUTE: Giprotymenneftegaz
 CITY: 625000, Tyumen
 ADDRESS: Respublika str., 62
 COUNTRY: RUSSIA

INSTITUTE: Giprotymenneftegaz. Nizhnevartovsk Branch
 CITY: 626440, Nizhnevartovsk
 ADDRESS: Chapaev str., 36a
 COUNTRY: RUSSIA

INSTITUTE: Hydrometeorological Research Institute of the Russian Federation
 (Gidromet-tsentr RF)
 CITY: 234353, Moscow
 ADDRESS: Bol'shevistskaya, 9-13
 COUNTRY: RUSSIA
 PHONE: (095) 255-2222

INSTITUTE: Institute for Biological Problems of the North (IBPS DVO RAN)
 MINISTRY: Far Eastern Branch of the Russian Academy of Sciences
 CITY: 685000, Magadan
 ADDRESS: Prosp. Kal Marxa, 24
 COUNTRY: RUSSIA
 PHONE: 2-0166

INSTITUTE: Institute for Earth Cryosphere
 MINISTRY: Siberian Branch of the Russian Academy of Sciences
 CONTACT: Melnikov, Vladimir Pavlovich
 CITY: 625033, Tyumen
 ADDRESS: P.O. Box 2774
 COUNTRY: RUSSIA
 PHONE: (3452)24-5267

INSTITUTE: Institute for Ecological Problems of Nature Usage
 MINISTRY: Ministry for Environmental Protection and Natural Resources, RF
 CONTACT: Ignatiev, Aleksei Evgenievich
 CITY: 125319, Moscow
 ADDRESS: Kochkovsky proezd, 3
 COUNTRY: RUSSIA

INSTITUTE: Institute for Problems in Mechanics
 MINISTRY: Russian Academy of Sciences
 CITY: 117526, Moscow
 ADDRESS: Prosp. Vernadskogo, 101
 COUNTRY: RUSSIA
 PHONE: (095) 434-2210

INSTITUTE: Institute for Problems in the Use of Natural Resource and Ecology (IPIPRE AI RB)
MINISTRY: Russian Academy of Sciences
CITY: 220114, Minsk
ADDRESS: Staroborisovskiy Route, 10
COUNTRY: BYELORUS
PHONE: 64-2161

INSTITUTE: Institute for Problems of Industrial Ecology of the North
MINISTRY: Kola Scientific Branch of the Russian Academy of Sciences
CITY: 184200 Apatity
ADDRESS: Fersman str., 14
COUNTRY: RUSSIA

INSTITUTE: Institute for Problems of Northern Development (IPOS SO RAN)
MINISTRY: Siberian Branch of the Russian Academy of Sciences
CONTACT: Cibulsky, Vladimir Romanovich
CITY: 625033, Tyumen
ADDRESS: P.O. Box 2774
COUNTRY: RUSSIA

INSTITUTE: Institute for Soil Sciences and Photosynthesis (IPFS)
MINISTRY: Russian Academy of Sciences
CONTACT: Gilichinsky, David Abramovich
CITY: 142292, Moscow Region
ADDRESS: Pushchino
COUNTRY: RUSSIA
PHONE: (2777)3-2604

INSTITUTE: Institute of Geography
MINISTRY: Siberian Branch of the Russian Academy of Sciences
CONTACT: Alekseev, Vladimir Romanovich
CITY: 664033, Irkutsk
ADDRESS: Ulan-batorskaya str., 1
COUNTRY: RUSSIA
PHONE: (3952)46-2639

INSTITUTE: Institute of Oil and Gas Problems
MINISTRY: Russian Academy of Sciences
CITY: 117917, Moscow
ADDRESS: GSP-1, Leninsky Prospect, 65, IPNG
COUNTRY: RUSSIA
PHONE: (095)135-8286

INSTITUTE: Institute of the Earth's Crust (IK RAN)
MINISTRY: Siberian Branch of the Russian Academy of Sciences
CONTACT: Leshchikov, Fedor Nikolaevich
CITY: 664033, Irkutsk
ADDRESS: Lermontov str., 128
COUNTRY: RUSSIA
PHONE: (3952)46-5568

INSTITUTE: Institute YakutNIIPROalmaz
MINISTRY: Industrial and Research Joint Company "Yakutalmaz"
CONTACT: Novik, Pavel Evgenievich
CITY: 678170, Sakha (Yakutia) Republic
ADDRESS: Lenin str., 39
COUNTRY: RUSSIA
PHONE: (41136)2-2849

INSTITUTE: IrkutskGiprotrans
MINISTRY: Ministry for Transport Construction, RF
CITY: 664007, Irkutsk
ADDRESS: Dekabrskikh Sobytii str., 49
COUNTRY: RUSSIA
PHONE: (3952)4-1774

INSTITUTE: Kiev Construction Bureau. Main Office of Mechanized Construction
CITY: 252033, Kiev
ADDRESS: Tarasovskaya, 9
COUNTRY: UKRAINE
PHONE: 224-4930

INSTITUTE: LenGidroproekt
CITY: 197136, St. Petersburg
ADDRESS: prosp. Shchorsa, 77/79
COUNTRY: RUSSIA

INSTITUTE: LenGiprotrans
MINISTRY: Ministry for Transport Construction
CONTACT: Mamzelev, Anatoly Petrovich
CITY: 196105, St. Petersburg
ADDRESS: Moskovsky prospect, 143
COUNTRY: RUSSIA
PHONE: (812)298-8777; (812)298-9388

INSTITUTE: Main A.V.Voeikov Geophysical Observatory (GGO)
MINISTRY: Ministry for Environmental Protection and Natural Resources, RF
CITY: 194018, St. Petersburg
ADDRESS: Karbyshev str., 7
COUNTRY: RUSSIA

INSTITUTE: Melnikov's Permafrost Institute (IMZ SO RAN)
MINISTRY: Siberian Branch of the Russian Academy of Sciences
CONTACT: Kamenskii, Vyacheslav
CITY: Mikhailovich
ADDRESS: 677010 Yakutsk
COUNTRY: RUSSIA

INSTITUTE: Melnikov's Permafrost Institute. Chita Division
MINISTRY: Siberian Branch of the Russian Academy of Sciences
CONTACT: Salnikov, Pavel Ivanovich
CITY: 672010, Chita-10
ADDRESS: P.O.Box 539
COUNTRY: RUSSIA
PHONE: (30222)3-9270

INSTITUTE: Melnikov's Permafrost Institute. Igarka Permafrost Research

MINISTRY: Siberian Branch of the Russian Academy of Sciences
CONTACT: Karpov, Egor Gavrilovich
CITY: 663200, Igarka
ADDRESS: Bolshoi Theater str., 13, INIMS
COUNTRY: RUSSIA
PHONE: (operator)2-4170

INSTITUTE: Melnikov's Permafrost Institute. North-East Permafrost Research
MINISTRY: Siberian Branch of the Russian Academy of Sciences
CONTACT: Perlshtein, Georgy Zakharovich
CITY: 685000 Magadan
ADDRESS: Gorkogo str., d.20a
COUNTRY: RUSSIA

INSTITUTE: Melnikov's Permafrost Institute. Viluiskaya Permafrost Research Station
MINISTRY: Siberian Branch of the Russian Academy of Sciences
CONTACT: Velikin, Sergei Aleksandrovich
CITY: 678185, Sakha (Yakutia) Republic
ADDRESS: Pos.Chernyshevskii, VNIMS, 2-6
COUNTRY: RUSSIA
PHONE: (operator)7-12

INSTITUTE: Mining Institute for Northern Affairs of the Yakutsk Science Center (IGDS YANTs SO RAN)
MINISTRY: Siberian Branch of the Russian Academy of Sciences
CITY: 677891, Yakutsk, GSP
ADDRESS: 50 Lyet VLKSM, 26
COUNTRY: RUSSIA
PHONE: 3-5916

INSTITUTE: Moscow Engineering-Construction Institute
CONTACT: Kudoyarov, L.I.
CITY: 129337, Moscow
ADDRESS: Yaroslavskoe Shosse, 26
COUNTRY: RUSSIA

INSTITUTE: Moscow Institute of Railways
MINISTRY: Ministry of Railways
CONTACT: Ashpiz, Evgeny Samuilovich
CITY: Moscow
ADDRESS: Obrastsov str., 15, MIIT
COUNTRY: RUSSIA
PHONE: (095)284-2241

INSTITUTE: Moscow State University. Faculty of Geography. Cryolithological and Glaciological Department
MINISTRY: Ministry for Higher Education
CONTACT: Konishchev, Vyacheslav Nikolaevich
CITY: 119899, GSP-3, Moscow
ADDRESS: Leninskie Gory
COUNTRY: RUSSIA
PHONE: (095)939-3673

INSTITUTE: Moscow State University. Faculty of Geology. Geocryological Department
MINISTRY: Ministry for High Education
CONTACT: Ershov, Eduard Dmitrievich
CITY: 119899, GSP-3, Moscow
ADDRESS: Leninskie Gory
COUNTRY: RUSSIA
PHONE: (095)939-1281

INSTITUTE: MosGiprotrans
MINISTRY: Ministry for Transport Construction
CONTACT: Isakov, Yurii
CITY: Moscow
ADDRESS: Korchagin str., 2
COUNTRY: RUSSIA

INSTITUTE: Nizhny Novgorod Branch of the State Planning Institute "SANTEX (Sanitation) Project"
CITY: 603005, Nizhny Novgorod
ADDRESS: Dzerzhinskogo, 26
COUNTRY: RUSSIA

INSTITUTE: Northeastern Complex Research Institute (SVKNII DVO RAN)
MINISTRY: Far-Eastern Branch of the Russian Academy of Sciences
CITY: 685000 Magadan
ADDRESS: Portovaya, 16
COUNTRY: RUSSIA
PHONE: 3-0923

INSTITUTE: Novokuznetsk Complex Research and Planning-Surveying
MINISTRY: All-Russia Research and Planning-Surveying Institute for Pipeline Hydrotransport
CITY: 654000 Kemerovskaya Oblast
ADDRESS: Novokuznetsk, Prosp. Kommunarov, 2
COUNTRY: RUSSIA

INSTITUTE: Novosibirsk Branch. (SibCNIIS)
MINISTRY: All-Russian Research Institute for Transport Construction
CITY: 630056, Novosibirsk
ADDRESS: Sofilskaya, 18
COUNTRY: RUSSIA
PHONE: 45-0214

INSTITUTE: Pechora State Research and Planning Institute for the Oil Industry (Pechoripineft)
CITY: 410720 Saratov
ADDRESS: Revolyutsii, 7
COUNTRY: RUSSIA
PHONE: 24-1513

INSTITUTE: Polyarnouralgeologia (Polar Urals Geology)
MINISTRY: Geological Survey, RF
CONTACT: Oberman, Naum Grigirievich
COUNTRY: RUSSIA

INSTITUTE: Production and Research Institute for Engineering Survey and Construction (PNIIS)
MINISTRY: Ministry for Construction RF
CONTACT: Baulin, Vladimir
CITY: Moscow
ADDRESS: Okruznoy proezd, 18
COUNTRY: RUSSIA
PHONE: (095) 366-2554

INSTITUTE: Production and Research Institute for Engineering Survey and Construction (PNIIS)
MINISTRY: Ministry for Construction
CITY: 181201 Pskovskaya Oblast
ADDRESS: Postal Branch Rodina
COUNTRY: RUSSIA
PHONE: 6-7251

INSTITUTE: Research Institute of Foundations and Underground Structures (NIIOSP)
MINISTRY: Ministry for Construction RF
CONTACT: Sadovsky, Andrei Vladimirovich
CITY: Moscow
ADDRESS: 2 Institutskaya str., 6
COUNTRY: RUSSIA
PHONE: (095)174-8354

INSTITUTE: Research Institute of Foundations and Underground Structures (NIIOSP). Norilsk
Department
MINISTRY: Ministry for Construction, RF
CONTACT: Grebenets, Valerii Ivanovich
CITY: 663316, Norilsk
ADDRESS: Oktyabrskaya str., 28
COUNTRY: RUSSIA
PHONE: (operator)5-3201

INSTITUTE: Research Institute on Transport Construction
MINISTRY: Ministry of Transport Construction
CONTACT: Cernant, Aleksandr Alfredovich
CITY: 129329 Moscow
ADDRESS: Kolskaya str., 1, CNIIS
COUNTRY: RUSSIA
PHONE: (095)180-9398

INSTITUTE: St. Petersburg Branch of the State Republican Institute for Planning Municipal
Water Supply and Sewage
CITY: 198096, St. Petersburg
ADDRESS: Kronshtadtskaya, 8
COUNTRY: RUSSIA

INSTITUTE: St. Petersburg State University of Transportation
MINISTRY: Ministry of Railways
CONTACT: Are, Felix Ernestovich
CITY: 1990031, St. Petersburg
ADDRESS: Moskovsky Prospect, 9
COUNTRY: RUSSIA
PHONE: (812)168-8318

INSTITUTE: St. Petersburg Zonal Scientific Research and Design Institute for Residential and Public Buildings (LenZNIEP)
CITY: St. Petersburg
COUNTRY: RUSSIA

INSTITUTE: Scientific Center "Chukotka"
MINISTRY: Far Eastern Branch of the Russian Academy of Sciences
CONTACT: Kotov, Anatoly Nikolaevich
CITY: 686710, Magadan Oblast
ADDRESS: Anadyr, Energetikov str., 15
COUNTRY: RUSSIA
PHONE: (operator)4-4370

INSTITUTE: Scientific-Production Association "Soyuzgaztekhnologiya"
CITY: 142717, Moskovskaya Oblast
ADDRESS: Postal Branch Razvilka
COUNTRY: RUSSIA
PHONE: 399-92-06

INSTITUTE: SibGiprotrans
MINISTRY: Ministry for Transport Construction
CONTACT: Vakulenko, Aleksandr Aleksandrovich
CITY: 630099, Novosibirsk
ADDRESS: Vokzalnaya magistral, 15
COUNTRY: RUSSIA
PHONE: (3832)29-5631

INSTITUTE: State and Legal Institute of the Ukrainian Academy of Sciences (IGP ANU)
CITY: 25001 Kiev,
ADDRESS: GSP-1, Trekhsvyatitelskaya, 4
COUNTRY: UKRAINE
PHONE: 228-5155

INSTITUTE: State Research Institute for Natural Protection of the Arctic and the North (GosNII OPAS)
MINISTRY: Ministry for Environmental Protection and Natural Resources, RF
CONTACT: Makeev, Vyacheslav Mikhailovich
CITY: 193231, St.-Petersburg
COUNTRY: RUSSIA

INSTITUTE: Stock Company "Rosneftegazstroy"
CONTACT: Mazu, I.I.
CITY: 109172, Moscow
ADDRESS: Narodnaya str., 4
COUNTRY: RUSSIA

INSTITUTE: Transcaucasian Regional Hydrometeorological Research Institute (ZakNIGMI)
CITY: 380012, Tbilisi
ADDRESS: D. Agmashenebeli, 150
COUNTRY: GEORGIA
PHONE: 95-0311

INSTITUTE: Tula Branch of the Central Research Institute for Geological Exploration
(Tu Ts NIGRI)
CITY: 300904 Tula
ADDRESS: Pos. Mendelevskiy
Mendeleva str., 16
COUNTRY: RUSSIA
PHONE: 22-6500

INSTITUTE: Tynda Permafrost Research Station
MINISTRY: Ministry for Transport Construction, (CNIIS)
CONTACT: Minailov, Gavriil Pavlovich
CITY: 676080, Tynda-2
ADDRESS: Merzlotnaya str., 2
COUNTRY: RUSSIA
PHONE: (41656)3-2515

INSTITUTE: Tynda Permafrost Station of the All-Russia Research Institute for Transport
Construction
CITY: 676000 Amurskaya Oblast
ADDRESS: Tynda
COUNTRY: RUSSIA

INSTITUTE: Tyumen State Research and Planning Institute for Natural Gas
(Tyymen NINGiprogaz)
CITY: 625016 Tyumen
ADDRESS: Respubliki str., 213 b
COUNTRY: RUSSIA
PHONE: 21-0165

INSTITUTE: TyumenNIIGiprogaz
CITY: 625000, Tyumen
ADDRESS: Melnikajta str., 90a
COUNTRY: RUSSIA

INSTITUTE: Vedeneev Research Institute on Hydrotechnic
MINISTRY: Ministry of Energy
CONTACT: Krivinogova, Ninel Fedorovna
CITY: 195220, St. Petersburg
ADDRESS: Gzhatskaya str., 21
COUNTRY: RUSSIA
PHONE: (812) 535-8890

INSTITUTE: VNIPIgasdobycha
CITY: 410000, Saratov
ADDRESS: Sakko & Vancetti str., 4
COUNTRY: RUSSIA

INSTITUTE: VNIPIgasdobycha. Novosibirsk Branch
CITY: 630000, Novosibirsk
ADDRESS: Komsomolsky prospect, 24
COUNTRY: RUSSIA

INSTITUTE: West-Siberian Zone Research and Production Institute
for Investigations on Construction
MINISTRY: Ministry for Construction, RF
CONTACT: Bezhentsev, Andrei Anatolievich
CITY: 625019, Tyumen
ADDRESS: Respublika str., 207
COUNTRY: RUSSIA
PHONE: (3452)22-8767

INSTITUTE: Yakutia State University
CITY: 677620, Yakutsk, GSP
ADDRESS: Belinsky str., 58
COUNTRY: RUSSIA
PHONE: (41122)2-7101

INSTITUTE: ZapSibNIGNI
CONTACT: Belyaev, Valerii Alekseevich
CITY: 625000, Tyumen
ADDRESS: Volodarsky str., 56
COUNTRY: RUSSIA
PHONE: (3452)26-3080

INSTITUTE: ZapSibPNIIS
MINISTRY: Ministry for Construction, RF
CONTACT: Gorbylev, Mikhail Ivanovich
CITY: 625016, Tyumen
ADDRESS: Respublika str., 207
COUNTRY: RUSSIA
PHONE: (3452)22-9366

INSTITUTE: Zheldorproekt for Siberia and Far East
CITY: 630043, Novosibirsk
ADDRESS: Uritsky str., 39
COUNTRY: RUSSIA

APPENDIX 7

INTERNATIONAL ARCTIC ENVIRONMENTAL DATA DIRECTORY

C. Hanson
National Snow and Ice Data Center
University of Colorado
Boulder, Colorado

In 1986 NOAA sponsored a workshop, with federal and academic representatives for the U.S. and Canada, to identify and focus interest in the establishment of a system to identify the existence of and provide access to environmental data for the Arctic. The U.S. Geological Survey took the initiative to organize a representative group to develop a plan for the Arctic Environmental Data Directory, AEDD. This AEDD Working Group designed and implemented an Arctic-wide, on-line database of data set descriptions, based on and hosted by the USGS-Reston Environmental Science Data Directory, ESDD. The primary focus of the incorporation of directory entries into the AEDD was on U.S.-held data sets, and on improving the quality of the descriptions in the AEDD. During 1994 the AEDD was moved to a PC-based system hosted by the USGS-Anchorage. The "new" AEDD is fully searchable using WAIS and World Wide Web software. The next project identified by the AEDD Working Group is to convert the existing AEDD descriptions to the DIF (Directory Interchange Format) used by the Global Change Master Directory. Future efforts will focus on obtaining additional data descriptions for inclusion in the directory. The primary benefit of the AEDD to the Arctic research community is the ability to search for available data using keywords and obtain contact information to order copies of the data.

The International Arctic Environmental Data Directory (ADD) effort was initiated at a workshop, organized by the USGS-hosted AEDD and UNEP/GRID, in Arendal, Norway, in September 1993. The goal of ADD is to provide a comprehensive compilation of existing sources of Arctic environmental data. The objectives of ADD are:

1. To establish the International Arctic Environmental Data Directory (ADD) as an authoritative, high-quality, and user-friendly directory of environmental data sources covering the circum-polar Arctic.
2. To assess the quality and reliability of data-set descriptions by means of a set of internationally agreed-upon, well-defined criteria, specifically the DIF (Directory Interchange Format) data description format of the Global Change Master Directory.
3. To provide access to the ADD internationally using the Internet and appropriate international standards. This will link the directory components into a consistent, high quality data and information source for the international Arctic community.
4. To identify and form working relationships with institutions that hold Arctic environmental data, inform them about the ADD, and seek to reference their data in the ADD.
5. To seek advice and counsel from, and develop feedback mechanisms with, the international Arctic science and user community to establish and maintain relevance of the ADD to key environmental issues.
6. To develop and implement a process using agreed upon standards to identify, gather, and maintain data set descriptions in the ADD which are consistent, complete, accurate, and timely to meet the needs of the user community.
7. To publish and disseminate ADD to promote the preservation and use of Arctic environmental data and information.

The ADD user community encompasses researchers in governmental agencies and academia, public interest groups and the private sector, the interested public, educators at all levels, and decision and public policy makers.

The ADD Steering Committee has defined Arctic environmental sciences as ". . . the broad spectrum of disciplines investigating the physical, biological, and cultural resources and environments of the Arctic."

The Steering Committee has recommended the ADD Council consist of a Data Manager Group, with one representative from each country participating in ADD, and the Advisory and User Group, with representatives from major Arctic monitoring and research organizations. The ADD Council will appoint the ADD Executive Committee, a group of five persons providing circum-Arctic geographic representation. The main task of the Executive Committee is to plan and carry out activities of the ADD, with recommendations from the ADD Council.

ADD is seeking strong involvement with institutions in Russia. A workshop is planned (in Fall 1995) in Moscow, to focus on accessibility of Russian data holdings and development of data directories. The Ministry of Environment Protection, Moscow, has offered to host the workshop and the USGS and UNEP/GRID have offered support.

As an active member of the AEDD Working Group, WDC-A for Glaciology is maintaining a close interest and involvement in the development of ADD. It will seek to ensure that the IPA GGD is entered into the AEDD/ADD system and that the operating procedures for the GGD are commensurate with those of the ADD and other international information systems.

ACRONYMS

ACSYS	Arctic Climate System Study
ADD	International Arctic Environmental Data Directory
AEDD	Arctic Environmental Data Directory
AMIP	Atmospheric Model Intercomparison Project
ARCSS	Arctic System Science
CLIMEX	Climate Extremes of the Past
CNIIS	Ministry for Transport Construction (Russian)
COMNAP	Council of Managers of National Antarctic Programs
CRREL	Cold Regions Research and Engineering Laboratory
DIF	Directory Interchange Format
DZAA	Depth of Zero Annual Amplitude
ESDIM	Earth Science System Data and Information Management
GCM	General Circulation Model
GCMD	Global Change Master Directory
GD	Glaciological Data
GEWEX	Global Energy and Water Experiment
GGD	Global Geocryological Database
GISP	Greenland Ice Sheet Program
GLOCOPH	Global Palaeohydrology Database Project
GRID	Global Resources Information Database
IASC	International Arctic Science Committee
IGBP	International Geosphere Biosphere Program
INQUA	International Union for Quaternary Research
IPA	International Permafrost Association
IPFS	Institute for Soil Sciences and Photosynthesis (Russia)
IPNG	Institute of Oil and Gas Problems (Russia)
ITEX	International Tundra Experiment
IUGG	International Union of Geodesy and Geophysics
LAII	Land/Atmosphere/Ice Interactions
NGD	National Geocryological Database
NIIOSP	Research Institute of Foundations and Underground Structures (Russia)
NOAA	National Oceanic and Atmospheric Administration
NSF	National Science Foundation
NSIDC	National Snow and Ice Data Center
OAI	Ocean/Atmosphere/Ice Interactions
PAGES	Past Global Environmental Changes

PALE	Paleoenvironments of Arctic Lakes and Estuaries
PNIIS	Production and Research Institute for Engineering Survey and Construction (Russia)
RF	Russian Federation
RGD	Regional Geocryological Database
SCAR	Scientific Committee on Antarctic Research
SHEBA	Surface Heat Budget of the Arctic
UKMO	United Kingdom Meteorological Office
UNEP	United Nations Environment Program
VNIIGMI	All-Russia Research Institute for Hydrometeorological Information
WAIS	Wide Area Information Servers
WCMC	World Conservation Monitoring Center
WG	Working Group
WGMS	World Glacier Monitoring Service

BOOK NOTES

Geotermiya Merzloy Zony Litosfery Severa Azii. (The Geothermal Regime of the Frozen Zone of the Lithosphere in Northern Asia) by V.T. Balobayev, Nauka Press, Novosibirsk, 1991, 193 pp. ISBN-5-02-029996-0.

This monograph by Dr. Veniamin Balobayev, Deputy Director of the Permafrost Institute in Yakutsk, provides a detailed treatise on the thermophysical controls of temperature regime at the surface, in the upper lithosphere, and at depth in frozen rock strata. Throughout, the mathematical expressions of the physical relationships are fully presented, together with observational material. In the foreword, the author traces the twentieth century worldwide development of geophysics and study of ground temperature regimes. Nineteenth century observations on ground temperature in Siberia by A. Middendorf are noted, as well as technical advances in measurements at the Yakutsk Permafrost Institute in the 1960s to 1980s under I.V. Mel'nikov's leadership. The author himself published extensively during this period.

Chapter 1 treats the surface energy budget and atmospheric influences, illustrated by a range of climatic maps and diagrams for northeastern Siberia, before proceeding to describe ground temperature conditions. Chapter 2 discusses the principles of ground heat flow and the role of soil properties, snow cover, and vegetation. Mean ground temperature conditions along the Lena River and elsewhere are described and data tables are included for a selection of localities. Chapter 3 follows a similar treatment for deep ground temperatures in frozen and unfrozen material. Ground heat flux is characterized for the structural zones of northern Asia and western Siberia, including the influence of surface and structural inhomogenities, and the thickness distribution is mapped. The final chapter examines the non-stationary cryolithozone in the past, present and future. Temperature profiles and cross-sections are shown and methods for reconstructing past conditions are detailed. Pleistocene and Holocene geothermal parameters are calculated and there is a brief discussion on possible future conditions associated with anthropogenic influences. There is a 16 page bibliography of Russian and Western literature. The book contains much information on permafrost conditions and the geothermal regime in northern Asia that would be of great interest to western scientists.

Osnovy Kriogeneza Litofery (Principles of the Cryogenesis of the Lithosphere) by N.N. Romanovskiy, Moscow University Press, 1993, 336 pp. ISBN-5-211-02379-X

This text by Professor Nikolai Romanovskiy is developed from courses in geology and geocryology taught at Moscow State University. It is aimed at advanced students in those fields, as well as hydrogeologists, exploration geologists, geographers, mining and petroleum engineers, and construction workers. The author previously published monographs on patterned ground formation

and ground water in the permafrost zone. Following an outline of historical development of studies of ground freezing and thawing and the associated surface phenomena, or cryogenesis, there are chapters on the climate and geomorphic factors determining the zonal and altitudinal characteristics of permanently and seasonally frozen ground, and periglacial phenomena. Chapter 4 deals with the formation and composition of syngenetic and epigenetic frozen ground, thermokarst complexes, including tundra soils and alas depressions, and sediments deformed by cryoturbation and ice wedge growth. Chapter 5 addresses the processes of permafrost development and characteristics of permafrost thickness in relation to ground heat flow and thermal regime, drawing particularly on V.T. Balobayev's work. The influence of structural and hydrogeological conditions, glacial history, and Arctic marine transgressions and regressions is also treated and the relationship of these factors to gas hydrate occurrence is examined. The final three chapters describe the characteristics of cryogenetic processes and permafrost in the Eurasian platform area, in mountain areas, and offshore in the Arctic shelf seas, respectively. Here, specific regional information is presented and illustrated with maps and cross-sections.

The book is illustrated by some 89 figures, including a few half-tone photographs. Most of the diagrams are schematic, in keeping with the textbook character of the work. Surprisingly, it contains only seven tables of numerical information. There is a brief index and a list of 132 references, all but 13 of them to Russian sources. Focusing as it does primarily on northern and eastern Russia, the book provides an up-to-date and useful overview of Russian geocryological research. An English translation would be useful to western readers in the field.

Roger G. Barry
Professor of Geography and
Director, WDC-A for Glaciology/NSIDC
CIRES
University of Colorado
Boulder, Colorado

Other Activities

NSIDC Distributed Active Archive Center

NSIDC participates as one of the NASA-funded Distributed Active Archive Centers (DAACs) in the Earth Observing System Data and Information System Project (EOSDIS). The Earth Observing System (EOS) is a long-term interdisciplinary and multidisciplinary research effort to study global-scale processes that shape and influence the Earth as a system. EOSDIS will manage the data resulting from NASA's research satellites and field measurement programs, and other data essential for the interpretation of these measurements. It will also facilitate access to data held in the archives of other government agencies, organizations, and countries.

ARCSS

NSIDC is the Data Coordination Center for the Arctic System Science (ARCSS) Program, a long-term multidisciplinary research effort funded by the National Science Foundation in an attempt to understand the Arctic physical, chemical, biological, and social processes that interact with the total Earth system in order to advance the scientific basis for predicting environmental change.

How to Contact the Center

MAILING ADDRESS:

WDC-A for Glaciology
CIRES

Campus Box 449

University of Colorado

Boulder, Colorado 80309-0449 USA

Telephone: 1 (303) 492-6199

Telefax: 1 (303) 492-2468

e-mail: nsidc@kryos.colorado.edu



Printed on recycled paper.

7/95

World

Data

Center-A

for

Glaciology (Snow and Ice)

National Snow and Ice Data Center Boulder, Colorado USA


*"It's a capital mistake
to theorize
without data."*

—**Sherlock Holmes**
(Sir Arthur Conan Doyle)

Snow and ice on land and water are dominant components of the polar regions and seasonally cover large areas of the northern hemisphere. Seasonal snow cover and sea ice are key regulators of the climate system, while perennial ice in glaciers, in ice sheets, and underground (permafrost) provides records of past climatic conditions and also stores vast quantities of fresh water.

Data on snow and ice conditions are required in assessment of hazards (snowstorms, avalanches, icebergs, glacier outburst floods), prediction of water resources and winter recreation (mountain snow-packs), navigation in the polar seas (sea ice and icebergs), and shipping on rivers and lakes (lake and river ice).

The World Data Center-A (WDC-A) for Glaciology archives and distributes data on all forms of snow and ice. It collaborates with other WDCs for Glaciology in Moscow, Russia (B), Cambridge, England (C), and Lanzhou, China (D), and with the World Glacier Monitoring Service in Zürich, Switzerland, to facilitate international access to data and information. It also works with the International Commission on Snow and Ice (ICSI), the International Permafrost Association (IPA), and the World Climate Research Programme (WCRP).



Mean annual ice drift vectors, based on drifting buoys, manned and unmanned camps, and mean annual surface pressure (isolines in millibars); H is high pressure.

The International Permafrost Association is conducting a survey of available data on past and current investigations of permafrost, seasonally frozen ground and periglacial conditions and related laboratory studies. An electronic mail form is available (see reverse). You may submit more than one form if data types differ substantially. Results of this survey will be reported in *Frozen Ground* and other relevant publications. Please complete this form and return it to your IPA national representative with a copy to the World Data Center-A for Glaciology, Attn: Claire Hanson, Campus Box 449, University of Colorado, Boulder, Colorado 80309-0449, U.S.A.

Summary: Give brief details of site, collection technique, measurement frequency, and quality. Indicate what ancillary data on site, date, data processing and metadata are available or can be reliably obtained (with sources).

Current storage medium: CD-ROM / Spreadsheet / Word processor / Database / Data centre (give name) / Paper
 Is your data at risk of being lost? YES / NO
 Bibliography (Published and unpublished reports about this data set; attach additional sheets if necessary)

Words: (Maximum of 10 such as Active layer, Permafrost thickness, Temperature, Moisture content, Ice content, Chemistry, etc.)